

6 Neurochemistry of the Main Olfactory System

M. Ennis · K. A. Hamilton · A. Hayar

1	Introduction	139
2	The Olfactory Epithelium	139
2.1	Organization and Cell Types	139
2.1.1	Microvillar Cells and Sustentacular Cells	140
2.1.2	Basal Cells	140
2.1.3	Olfactory Receptor Neurons	141
2.2	Odor Receptors, Transduction, and Physiology of ORNs	141
2.3	Topography of ORN Projections to MOB	142
3	Main Olfactory Bulb	143
3.1	Olfactory Nerve Layer	143
3.2	Glomerular Layer	144
3.2.1	Neuron Types of the GL	144
3.2.2	Neurochemistry of JG Neurons	147
3.2.3	Synaptic and Bicompartamental Organization of the GL	148
3.3	External Plexiform Layer	154
3.3.1	Neuron Types of the EPL	154
3.3.2	Circuitry, Synaptology, and Sublaminar Organization of the EPL	156
3.4	Mitral Cell Layer, Mitral/Tufted Cell Synapses, and Neurochemistry	158
3.4.1	Neurochemistry of Mitral Cells	158
3.5	Internal Plexiform Layer and Granule Cell Layer	159
3.5.1	Neuron Types of the GCL	160
3.6	Subependymal Layer and Rostral Migratory Stream	161
3.7	Outputs and Inputs of the MOB	162
3.7.1	Outputs of MOB	162
3.7.2	Extrinsic Inputs to MOB	163
3.8	Amino Acid Receptor and Gap Junction Distribution in MOB	163
3.8.1	Iontropic Glutamate Receptors	163
3.8.2	Metabotropic Glutamate Receptors	164
3.8.3	GABA Receptors	165
3.8.4	Glycine and Glycine Receptors	166
3.8.5	Gap Junctions	166
3.9	Neuromodulatory Inputs to MOB	167
3.9.1	Cholinergic Inputs to MOB	167
3.9.2	Dopaminergic (DA) Input to MOB	168
3.9.3	Noradrenergic (NE) Input to MOB	168
3.9.4	Serotonergic (5-HT) Inputs to MOB	171
4	Olfactory Input Regulates Neurochemistry of Specific MOB Neurons	171
5	Primary Olfactory Cortex: Piriform Cortex	173

5.1	Organization and Architecture	173
5.2	PC Cell Types	173
5.2.1	Pyramidal Cells	174
5.2.2	Interneurons	174
5.3	PC Circuitry	176
5.3.1	Input from MOB	176
5.3.2	Intrinsic and Association Connections	177
5.4	Extrinsic Outputs of PC	178
5.4.1	Feedback to the MOB	178
5.4.2	Cortical Projections	178
5.4.3	Subcortical Projections	179
5.5	Amino Acid Receptor Distribution in PC	179
5.5.1	Glutamate Receptors	179
5.5.2	GABA Receptors	179
5.6	Neuromodulatory Inputs to PC	179
5.6.1	Cholinergic Inputs to PC	179
5.6.2	Dopaminergic Inputs to PC	180
5.6.3	Noradrenergic Inputs to PC	180
5.6.4	Serotonergic Inputs to PC	182

Abstract: Many aspects of animal and human behavior are guided by, or dependent on, the sense of smell-olfaction. This chapter summarizes the major facts of the anatomy, neurochemistry, molecular biology and physiology of the olfactory system. We emphasize the mammalian olfactory system, particularly rodents, because of their widespread use as a laboratory model and the rich database related to this species.

List of Abbreviations: 5-HT, serotonin; 6-OHDA, 6-hydroxydopamine; AChE, acetylcholinesterase; ACIII, adenylate cyclase type III; AMPA, alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid; AOB, accessory olfactory bulb; AON, anterior olfactory nucleus; AP4, 2-amino-4-phosphonobutyric acid; BAPTA, bis (2-aminophenoxy)ethane-N,N,N,N-tetra-acetate; BDNF, brain-derived neurotrophic factor; CaM, calmodulin; cAMP, cyclic adenosine monophosphate; CB, calbindin; CCK, cholecystokinin; ChAT, choline acetyltransferase; CNGs, cyclic nucleotide gated channels; CR, calretinin; CREB, cAMP-response element-binding protein; CRF, corticotropin-releasing factor; DA, dopamine; DB, diagonal band; DBH, dopamine- β -hydroxylase; EM, electron microscopy; ENK, enkephalin; EPL, external plexiform layer; ET, external tufted; GABA, gamma-aminobutyric acid; GAD, glutamic acid decarboxylase; GC, granule cell; GCL, granule cell layer; GL, glomerular layer; IAS, intrabulbar association system; IP₃, inositol trisphosphate; IPL, internal plexiform layer; IR, insulin receptor; JG, juxtglomerular; LC, locus coeruleus; LHRH, leutenizing hormone releasing hormone; LOT, lateral olfactory tract; M, muscarinic; MAPK/ERK, mitogen-activated protein kinase/extracellular signal-related kinase; MCL, mitral cell layer; mGluR, metabotropic glutamate receptor; MOB, main olfactory bulb; NADPH, nicotinamide adenine dinucleotide phosphate; NAG, N-acetyl-aspartyl-glutamate; NDB, nucleus of the horizontal limb of the diagonal band; NE, norepinephrine; NMDA, N-methyl D-Aspartate; NPY, neuropeptide Y; OCNC1, olfactory cyclic nucleotide gated channel subunit-1; OMP, olfactory marker protein; ON, olfactory nerve; ONL, olfactory nerve layer; ORN, olfactory receptor neurons; PC, piriform cortex; PG, periglomerular; PKC, protein kinase C; POC, primary olfactory cortex; PV, parvalbumin; RMS, rostral migratory stream; RNA, ribonucleic acid; SA, short axon; SP, substance P; SSA, superficial short axon; TH, tyrosine hydroxylase; TRH, thyrotropin-releasing hormone; Trk, tropomyosin related kinase; VG, Van Gehuchten; VIP, vasoactive intestinal polypeptide

1 Introduction

The olfactory system consists of two parallel systems, the main olfactory system and the accessory olfactory system. This chapter focuses on the main olfactory system in mammals, specifically on the main olfactory bulb (MOB) and primary olfactory cortex (POC), and it emphasizes the neuroanatomy, neurochemistry, and neurophysiology of these regions in rodents. The sense of olfaction is critically important for food consumption, emotional responses, aggression, maternal and reproductive functions, neuroendocrine regulation, and the recognition of conspecifics, predators, and prey. In many species, olfaction plays a more pivotal role in these functions than in humans, with olfactory cues exceeding visual or auditory cues in importance. The olfactory system has long been an attractive model to study cellular mechanisms underlying the encoding, transfer, processing, and decoding of sensory information. Recent interest in this area has been sparked by a series of dramatic breakthroughs over the past decade in our understanding of the organization and function of the peripheral olfactory system, cloning of the olfactory receptors, and identification of the olfactory transduction machinery (Breer, 2003). These advances, together with a wealth of accumulated knowledge about the anatomy and connectivity of the MOB (Shipley et al., 1996, 2004; Shepherd et al., 2004), have set the stage for experiments aimed at unraveling the mechanisms of early sensory processing by bulbar and cortical circuits.

2 The Olfactory Epithelium


2.1 Organization and Cell Types

Odor molecules that enter the nasal cavities and bind to first-order neurons, olfactory receptor neurons (ORNs), which are contained in a neuroepithelial sheet lining the upper and caudal aspects of the nasal

septum and the cribriform plate region of the nasal cavities. ORNs are located in a specific portion of the nasal epithelium, the olfactory neuroepithelium, which is thicker than the surrounding respiratory epithelium and contains a number of distinct cell types (● *Figure 6-1*).

■ Figure 6-1

The olfactory epithelium showing the organization of the olfactory receptor neurons (ORNs) and other cell types



2.1.1 Microvillar Cells and Sustentacular Cells

As the name suggests, microvillar cells (not shown) have superficial microvilli in contact with the mucus (Moran et al., 1982; Muller and Marc, 1984; Rowley et al., 1989; Morrison and Costanzo, 1990). They lie deep within the neuroepithelium and extend an “axon-like” projection, which can reach the MOB, as suggested by tract-tracing studies (Rowley et al., 1989). Sustentacular cells separate and partially wrap the ORNs. Like the microvillar cells, their surface contains microvilli that project into the mucus layer. They may function in mucus composition regulation (Getchell et al., 1984; Getchell and Getchell, 1992). As they express molecules of the P450 enzyme systems (Hadley and Dahl, 1982), they may play a role in detoxification.

2.1.2 Basal Cells

The basal cells are deeply located globose- and horizontal-type stem cells for the replacement of the ORNs (Cuschieri and Bannister, 1975a, b; Graziadei and Graziadei, 1979), which in rodents have a life span of approximately 40 days.

2.1.3 Olfactory Receptor Neurons

ORNs are bipolar sensory neurons with both an apical dendrite and a deep or basal axon (Cajal, 1911a, b). The dendrites extend superficially, forming an olfactory knob with multiple cilia extending into the mucus (Menco, 1984; Menco and Farbman, 1985a, b). Odor transduction takes place in the cilia. Basally, the ORNs give rise to axons—the olfactory nerve (ON) fibers. The axons are ensheathed by specialized Schwann cells, the ensheathing cells (DeLorenzo, 1957). These fibers form bundles, which then collect as groups of fascicles, pass through the cribriform plate, and synapse in the MOB.

2.1.3.1 Neurochemistry of ORNs

Olfactory Marker Protein ORNs express high levels of olfactory marker protein (OMP), which is unique in the olfactory system to ORNs (Margolis, 1972; Keller and Margolis, 1975). OMP is found in a number of mammalian species and it appears to be expressed in all mature ORNs. Studies in mice containing a null mutation for OMP suggest that this protein may play a role in ORN adaptation to odors (Ivic et al., 2000), signal amplification, and transduction (Youngentob et al., 2004).

Glutamate, Carnosine, Copper, and Zinc ORNs contain glutamate and glutamate antibodies stain ON fibers and axon terminals in the glomerular layer (GL) (Liu et al., 1989; Sassoè-Pognetto et al., 1993). Carnosine, a soluble dipeptide, is uniquely expressed in high concentrations in mammalian ORNs, and it is present in ON axon terminals in the GL (Ferriero and Margolis, 1975; Margolis, 1980; Biffo et al., 1990). Carnosine colocalizes with glutamate in the ON axon terminals (Sassoè-Pognetto et al., 1993), and it satisfies criteria for neurotransmitter candidacy, including (1) carnosine synthetic and degradative enzymes are present in ORNs, (2) the peptide is released by depolarization in a Ca^{2+} -dependent manner in ON synaptosomes, and (3) high affinity binding sites for carnosine are present in the GL (Ferriero and Margolis, 1975; Margolis, 1980; Burd et al., 1982; Rochel and Margolis, 1982; Margolis et al., 1983, 1985, 1987; Margolis and Grillo, 1984; Biffo et al., 1990). Zinc and copper are also present in high concentrations in ON axon terminals (Biffo et al., 1990). The potential neuromodulatory roles of carnosine, zinc, and copper are discussed later.

2.2 Odor Receptors, Transduction, and Physiology of ORNs

The binding of odors to receptors located on the cilia of ORNs elicits electrical signals (see Reed, 1992 for review). Olfactory receptors comprise a family of approximately 1,000 G-protein coupled receptors with 7 transmembrane domains that correspond to roughly 1,000 genes. Each mammalian ORN seems to express only a single receptor gene and approximately 10,000 ORNs express the same receptor, although there may be some exceptions (Rawson et al., 2000). The $\sim 10,000$ ORNs expressing the same receptor gene are scattered across one of the four epithelial expression zones (Buck and Axel, 1991; Ressler et al., 1993; Vassar et al., 1993; Strotmann et al., 1994; Sullivan et al., 1995). As discussed in [Section 2.3](#), collections of ORNs expressing the same receptor project focally onto very few glomeruli in the MOB. Biochemical and electrophysiological studies suggest that the cAMP signal transduction pathway predominates in mammalian ORNs (Schild and Restrepo, 1998; Firestein, 2001; Zufall and Munger, 2001). Odors activate a G-protein, G_{olf} , which activates adenylate cyclase type III (ACIII), leading to a rise in cyclic nucleotide levels (i.e., cAMP). cAMP binds to, and opens, cyclic nucleotide gated channels (CNGs) (Zufall et al., 1991). These channels are permeable to Na^+ , K^+ , and Ca^{2+} (Frings et al., 1995). The channels are preferentially permeable to Ca^{2+} , and their activation increases intracellular Ca^{2+} within the cilia (Leinders-Zufall et al., 1997, 1998). Increased Ca^{2+} leads, in turn, to activation of a Ca^{2+} -activated chloride conductance (Kleene and Gesteland, 1991; Lowe and Gold, 1993), which further depolarizes the cell (due to high intracellular Cl^- levels relative to the mucus), leading to the generation of action potentials that propagate down the axon to the MOB. Genetic null mutations for G_{olf} (Belluscio et al., 1998), CNGA2 subunits (also called OCNC1) (Brunet et al., 1996), and ACIII (Wong et al., 2000) firmly establish the essential role for these molecules in odor transduction. Mice with null mutations for any of these three transduction elements are functionally anosmic. The transduction process is negatively regulated via several mechanisms, allowing for relatively

rapid adaptation of odor responses. Increased intracellular Ca^{2+} triggers adaptation, as demonstrated by reduced adaptation when intracellular Ca^{2+} is chelated by BAPTA (Leinders-Zufall et al., 1998, 1999). The rise in intracellular Ca^{2+} activates calmodulin, which binds to CNGs, thereby decreasing their affinity for cAMP. The CNGA4 subunits are especially important for this mechanism of adaptation (Bradley et al., 2001; Munger et al., 2001). Ca^{2+} -calmodulin has also been shown to activate CaM kinase II, leading to phosphorylation of ACIII and reduced production of cAMP. CaM kinase II phosphorylates phosphodiesterase, thus increasing inactivation of cAMP. Finally, there is tonic inhibition of CNGs by extracellular Ca^{2+} , which is quite high in mucus, perhaps serving to increase the ORN signal-to-noise ratio (Zufall and Firestein, 1993).

2.3 Topography of ORN Projections to MOB


The ORNs of each olfactory epithelium give rise to several million axons that coalesce in the ON layer (ONL) of the ipsilateral MOB and synapse within the glomeruli (Meisami, 1989). Because of the widespread epithelial distribution of ORNs expressing the same receptor, each glomerulus in the MOB receives input from widely dispersed ORNs. In the same manner, adjacent ORNs may project to spatially distant glomeruli (Astic and Saucier, 1986; Saucier and Astic, 1986; Astic et al., 1987; Clancy et al., 1994; Schoenfeld et al., 1994). Early mapping studies showed that there is a crude topography such that ORNs located in the four expression zones project to homologous regions of MOB (Astic and Saucier, 1986; Saucier and Astic, 1986; Clancy et al., 1994; Schoenfeld et al., 1994). The mRNA for the receptor genes is orthogradely transported along ORN axons to their axon terminals. This property led to the discovery that despite the scattered distribution of ORNs expressing the same receptor within the epithelial zones, there is a remarkably precise topographic projection of ORNs expressing the same receptor. Specifically, the ORNs expressing the same receptor project to one or two glomeruli located on the medial and lateral side of each MOB (Figure 6-2) (Ressler et al., 1993, 1994; Vassar et al., 1994; Mombaerts et al., 1996; Wang et al., 1998; Potter et al., 2001; Treloar et al., 2002). Studies in transgenic animals showed that this projection pattern is topographically fixed across animals. That is, the same glomeruli identified in different mice receive inputs from the same restricted population of ORNs bearing the same receptor (Mombaerts et al., 1996; Wang et al., 1998; Potter et al., 2001; Treloar et al., 2002). Rough calculations confirm an approximately 1:2 ratio between the number of different types of receptors (~1,000) and the total number of glomeruli (~1,800) in mice.

The implications of these receptor localization and expression studies for the odor specificity of individual ORNs are still not fully understood. However, experiments suggest that a receptor may recognize a specific ligand or epitope expressed in common by a number of odors, and that the same receptor may respond to several similar epitopes (Krautwurst et al., 1998; Zhao et al., 1998). In turn, this suggests that an ORN expressing the same receptor responds differentially to a family of odorants containing a common epitope, as well as to other odorants containing different, but structurally similar epitopes. In support of this notion, a variety of electrophysiological studies demonstrate that individual ORNs respond with different degrees of excitation to a spectrum of odors (Malnic et al., 1999; Duchamp-Viret et al., 1999; Hamana et al., 2003). Together, these findings imply that an individual odor will activate a distinct pattern of glomerular activity that depends on the number of specific ORN-receptor gene groups activated.

The results of odor mapping studies in the bulb are consistent with the preceding model. Functional studies utilizing 2-deoxyglucose (Jourdan et al., 1980; Astic and Saucier, 1981; Benson et al., 1985; Johnson and Leon, 2000a, b; Johnson et al., 1998), *c-fos* (Onoda, 1992; Guthrie and Gall, 1995a, b), magnetic resonance imaging (Yang et al., 1998), Ca^{2+} or voltage-sensitive dye imaging (Cinelli et al., 1995; Friedrich and Korshung, 1997; Wachowiak and Cohen, 1999, 2001; Wachowiak et al., 2000), and intrinsic imaging (Rubin and Katz, 1999; Uchida et al., 2000; Belluscio and Katz, 2001; Rubin and Katz, 2001) suggest that low concentrations of single odorants activate a restricted number of glomeruli. Often, several responsive glomeruli are clustered. These studies suggest that different odors activate unique sets of glomeruli that are topographically fixed from animal to animal. Higher odorant concentrations activate larger numbers of glomeruli, probably resulting from activation of additional ORNs that express a different receptor that has a lower affinity for that odorant. These mapping studies suggest that glomeruli are functional modules that

Figure 6-2

The projections of ORNs to the glomerular layer (GL) of the MOB. Note that ORNs expressing different odorant receptor genes (shown as blue, red, or green cells) are interspersed and widely distributed, yet the axons of ORN expressing the same odorant receptor gene converge onto the same glomerulus (or pairs of medial and lateral glomeruli) in the GL (represented as blue, red, or green glomeruli). EPL, external plexiform layer; MCL, mitral cell layer; ONL, olfactory nerve layer



represent a map of the activity of ORNs. As odors typically activate groups of glomeruli, the olfactory network must extract information from such patterns to recognize and distinguish different odors. The MOB is the first component in the olfactory system that performs this neural computation.

3 Main Olfactory Bulb


The MOB in rodents is situated at the rostral pole of the cranial cavity and it is connected to the frontal cortex by a slender peduncle. The bulb can be thought of as an elongated onion composed of distinct layers or laminae that are organized as concentric circles. These layers, from superficial to deep, are the ONL, GL, external plexiform layer (EPL), mitral cell layer (MCL), internal plexiform layer (IPL), GC layer (GCL), and the ependymal layer (▶ [Figures 6-3](#) and ▶ [6-4](#)).

3.1 Olfactory Nerve Layer

The outer and most superficial MOB layer, the olfactory nerve layer (ONL), consists of ON axons and glial cells (Cajal, 1911a, b; Pinching and Powell, 1971b; Doucette, 1989). The glial or ensheathing cells envelop the ON axons and express the Schwann cell marker S100. The deepest third of the ONL also contains

Figure 6-3

Architecture of the MOB (A, B) and primary olfactory cortex (POC, C). A and B: Coronal section (Nissl stain) of the rat MOB at low (A) and high (B) magnifications. C: Coronal section through the rat brain showing several structures of the POC. Abbreviations: ac, anterior commissure; Acb, nucleus accumbens; AONvp, anterior olfactory nucleus, ventroposterior division; DP, dorsal peduncular cortex; IL, infralimbic cortex; Pir, piriform cortex; SEL, subependymal layer; Scale bar in A and C = 1 mm. Modified from Handbook of Chem. Neuroanat. Integrated Sys. CNS, Vol. 12, Part III, Chapter III, The Olfactory System, M. Shipley et al., pp. 469–573, 1996, with permission from Elsevier



astrocytes that reside between bundles of ensheathing cell-wrapped axons (Doucette, 1984; Bailey and Shipley, 1993). Astrocytes in this same region express QPRT, the degradative enzyme for quinolinic acid, an NMDA receptor agonist (Bailey and Shipley, 1993). Quinolinic acid, like glutamate, may modulate glutamatergic transmission between the ON axon terminals and postsynaptic targets. The roles of other neuroactive substances of ORN axons (glutamate, zinc, carnosine), as well as receptors expressed by the ON axon terminals (D2, GABA_B), are discussed later.

3.2 Glomerular Layer


Deep to the ONL, the GL comprises neuropil-rich ovoid structures—the glomeruli—each of which is surrounded by a shell of small neurons and glia. Within the glomeruli, ON axons synapse with mitral and tufted cells, as well as with the intrinsic neurons of the GL. Glomerular diameters are 80–160 μm in rat (Meisami and Safari, 1981; Meisami and Sendera, 1993) and 50–100 μm in mouse (Royet et al., 1988), and the number of glomeruli varies among species: 2000–3000 glomeruli/bulb for rabbits (Allison, 1949); 1800–2000 for mice (Allison, 1953; White, 1972; Royet et al., 1988); 3000 for rats (Meisami and Safari, 1981). More recent stereological studies yielded higher numbers: 6300 glomeruli in rabbit and 4200 in rat (Royet et al., 1998). Adjacent glomeruli are somewhat isolated from each other by astrocytes residing in the glomerular shell. Wedge-shaped astrocytes with somata in the glomerular shell send branched processes into the glomerular core (Bailey and Shipley, 1993; Chao et al., 1997). The astrocytes are restricted to a single glomerulus and they appear to cordon off adjacent glomeruli. This observation provides additional evidence that each glomerulus functions as a discrete unit.

3.2.1 Neuron Types of the GL

The neurons of the GL are classified as three cell types, which include: (1) periglomerular (PG) cells, (2) external tufted (ET) cells, and (3) short axon (SA) cells (Golgi, 1875; Van Gehuchten and Martin, 1891;

■ Figure 6-4

The basic circuitry of the MOB. Axons of ORNs travel in the ONL and synapse in the GL on the dendrites of mitral cells (MC), tufted cells (external tufted cell, ET; middle tufted cell, MT), and generic juxtglomerular (JG) neurons, which include periglomerular cells (PG), ET cells, and short axon cells (SA). SA cells interconnect different glomeruli. There are serial and reciprocal synapses between the apical dendrites of mitral/tufted cells and the processes of JG neurons. Superficial tufted cells (ST) are located in the superficial EPL or at the GL–EPL border. The lateral dendrites of mitral/tufted cells form serial and reciprocal synapses with the apical dendrites of granule cells (GC) in the EPL. GCs are located in the GCL and the MCL. The axons of mitral/tufted cells project locally to GCs (not shown) and also to primary olfactory cortex via the lateral olfactory tract (LOT). The bulb also contains other populations of interneurons neurons, including the van Gehuchten cells (VG) within the EPL




Blanes, 1898; Cajal, 1911a, b; Pinching and Powell, 1971a, b, c; Pinching and Powell, 1972b, c). Collectively, the intrinsic neurons of the GL are referred to as juxtglomerular (JG) cells. The term JG is also used here with regard to the cited studies in which the subtype of glomerular neuron was not specified. The morphology and features of these cells are illustrated in [Figure 6-5](#) and are only briefly reviewed here as more detailed descriptions are available (Hayar et al., 2004a, b).

3.2.1.1 ET Cells These are the largest (10–15 μm) cells in the GL, and electrophysiological recordings have confirmed that they are excitatory. They are dispersed in the juxtglomerular regions, surrounding/deep to the glomeruli. Most have one apical dendrite that arborizes extensively throughout one glomerulus (Pinching and Powell, 1971a; Hayar et al., 2004a, b). Rarely, ET cells have two or three apical dendrites that ramify in different glomeruli. Most ET cells have secondary or lateral dendrites that extend in the superficial EPL. Some ET cells have axons that appear to synapse with PG cells or SA cells, or more infrequently, project out of MOB (Pinching and Powell, 1971a). ET cells are somewhat similar to tufted cells of the EPL and mitral cells, but growing evidence suggests that all tufted cell subtypes exhibit distinct anatomical and physiological properties.

3.2.1.2 PG Cells These cells are the most numerous ones in the GL, and they are thought to be inhibitory in nature. They are small (5–8 μm), spherical or ovoid, and are distributed in the periglomerular regions

■ Figure 6-5

Neuron types in the MOB. All neurons were labeled in MOB slices by intracellular filling with biocytin during electrophysiological recordings. A–C: Reconstruction of three cells representing the three major types of JG cells in the GL: external tufted cell (A); periglomerular cell (B); short axon cell (C). D–H: Photomicrographs of biocytin-filled superficial tufted (D), EPL interneuron (E), middle tufted cell (F) in the EPL; mitral cell (G, soma in the MCL); granule cell (H, soma in the GCL). Calibration bars in A–H = 100 μ m. Figures D–F reprinted from K.A. Hamilton et al., *Neuroscience* 133: 819–829, 2005, with permission from Elsevier, Ltd



surrounding the glomeruli. Their dendrites are typically restricted to a small subregion of a glomerulus (Pinching and Powell, 1971a; Hayar et al., 2004a, b). These dendrites most commonly receive synapses from axon terminals containing spherical vesicles that have been thought to be collateral branches of ET cell axons and centrifugal fibers (Pinching and Powell, 1971c, 1972c). Other synapses along PG dendrites include asymmetrical (morphologically excitatory) synapses from the ON terminals and from mitral/tufted cell dendrites. Some of the mitral/tufted cell synapses are paired with reciprocal symmetrical (morphologically inhibitory) synapses back onto the parent mitral/tufted cell dendrites (Pinching and Powell, 1971b; Kasowski et al., 1999; Toida et al., 1998, 2000). Physiological recordings indicate that PG cells also receive monosynaptic dendrodendritic excitatory input from ET cells (Hayar et al., 2004b). PG cell axons appear to be rare (Pinching and Powell, 1971a; Hayar et al., 2004a), but they have been reported to extend over distances equivalent to four to five glomeruli. They appear to form symmetrical

(morphologically inhibitory) synapses onto mitral/tufted cell dendrites and ET cells and other JG cells (Pinching and Powell, 1971c).

3.2.1.3 SA Cells These cells are roughly of the same size (8–12 μm) as the ET cells. They are distinguished by multiple dendrites that seem to harvest information from multiple glomeruli (Hayar et al., 2004b). The dendrites may receive synaptic inputs from ET cell dendrites (Hayar et al., 2004b), tufted cell collaterals, or from other SA cells. SA cells have axons that can extend up to 1–2 mm within the GL (Aungst et al., 2003). The axons appear to synapse onto the dendrites of PG cells (Pinching and Powell, 1971a). They do not receive direct ON input (Pinching and Powell, 1971c; Hayar et al., 2004b).

3.2.2 Neurochemistry of JG Neurons

As illustrated in [Tables 6-1](#) and [6-2](#), GL neurons are rich in neuroactive substances and mapping studies have revealed that PG cells in particular are neurochemically heterogeneous (Halász, 1990).

Table 6-1

Neurotransmitters in MOB neurons

Transmitter	Cell type (layer)	References
Aspartate	Mitral	Fuller and Price (1988), Halász (1987), Watanabe and Kawana (1984)
CCK	JG, ET, Tufted, Cajal	Bonnemann et al. (1989), Matsutani et al. (1988), Seroogy et al. (1985)
CRF	Mitral, Tufted	Bassett et al. (1992), Imaki et al. (1989)
DA	JG	Baker et al. (1983,1984,1988), Baker (1986), Gall et al. (1987), Halász et al. (1981), McLean and Shipley (1988)
ENK	PG, Granule, EPL Int,	Bogan et al. (1982), Davis et al. (1982), Kosaka et al. (1995)
GABA	PG, Granule, EPL Int.	Gall et al. (1987), Kosaka et al. (1987d), Mugnaini et al. (1984a, b)
GABA + DA	PG	Kosaka et al. (1985,1988)
GABA + Parvalbumin	Sup. Tufted (EPL)	Kosaka et al. (1987d)
GABA + ENK	Granule	Kosaka et al. (1987d)
GABA and SP and DA	PG	Davis et al. (1982)
Glutamate	JG, Mitral, Tufted	Liu et al. (1989)
NAG	Mitral	Blakely et al. (1987), Ffrench-Mullen et al. (1985)
NADPH-diaphorase	SA (GL), EPL Int.	Alonso et al. (1995), Davis (1991), Scott et al. (1987), Villalba et al. (1989)
Neurotensin	EPL Int.	Matsutani et al. (1988)
NPY	PG, GCL, EPL Int., SA	Briñón et al. (1992), Gall et al. (1986), Nakajima et al. (1996), Ohm et al. (1988), Sanides-Kohlrausch and Wahle (1990a), Scott et al. (1987), Seroogy et al. (1989)
Somatostatin	JG, SA (GL), Deep SA, EPL Int.	Davis et al. (1982), Matsutani et al. (1988), Scott et al. (1987), Seroogy et al. (1989)
SP	JG, Sup. Tufted (EPL), EPL Int.	Baker (1986), Davis et al. (1982), Kream et al. (1984), Wahle et al. (1990)
TRH	PG	Merchenthaler et al. (1988), Tsuruo et al. (1988)
VIP	PG, Sup. Tufted (EPL), Van Gehuchten (EPL)	Crespo et al. (2002), Gall et al. (1986), López-Mascaraque et al. (1989), Sanides-Kohlrausch and Wahle (1990b)

EPL Int., EPL interneuron; Sup. Tufted, superficial tufted cell. Adapted from Handbook of Chem. Neuroanat. Integrated Sys. CNS, Vol. 12, Part III, Chapter III, The Olfactory System, M. Shipley et al., pp. 469–573, 1996, Elsevier, Ltd

■ Table 6-2

Relative frequency of neurons expressing calcium binding proteins in a 25- μ m-thick rat MOB section

Cell Type (Layer/s)	Calcium Binding Protein			
	Neurocalcin	Cabindin D-28k	Calretinin	Parvalbumin
Periglomerular (GL)	+++++	+++++	+++++	++
Sup. Short Axon (GL, EPL)	–	++	+++	++++
Sup. Tufted (EPL)	+++++	–	–	–
Van Gehuchten (EPL)	+	+	++	+++
Mitral (MCL)	–	–	+++	–
Deep Short Axon (MCL/IPL, IPL, GCL)	+	+	+	++
Granule (MCL, IPL, GCL)	++++	–	+++++	+

+++++, more than 500 cells per section; +++++, 250–500 cells per section; +++, 100–250 cells per section; ++, 25–100 cells per section; +, less than 25 cells per section; –, immunonegative. MCL/IPL, IPL border with MCL. Adapted from Briñón et al., *J. Comp. Neurol.* 407: 404–414 (1999), with permission from John Wiley & Sons, Inc

Electrophysiologically, ET cells appear to be glutamatergic (Hayar et al., 2004b) and they may correspond to JG cells stained by glutamate antibodies (Liu et al., 1989). The GL also contains the largest population of dopamine (DA) containing cells in the brain. The majority of DA containing cells are PG cells, but some ET cells are also dopaminergic (Halász et al., 1977; Davis and Macrides, 1983; Halász et al., 1985; Gall et al., 1986; McLean and Shipley, 1988). Another major neurotransmitter of JG neurons is GABA, which appears to be predominantly contained in PG cells (Ribak et al., 1977; Mugnaini et al., 1984a; Kosaka et al., 1985, 1987a, b, c, d, 1988). Some JG cells that send axons to the deeper GCL contain nicotinamide adenine dinucleotide phosphate (NADPH)-diaphorase, neuropeptide-Y (NPY), and somatostatin, which Scott et al. (1987) suggested may provide a direct route for PG cells to influence granule cells (GCs). Davis (1991) concluded that NADPH in the GL is primarily, if not exclusively, contained in PG cells. The NADPH PG cells are distinct from cells that express calbindin immunoreactivity (Alonso et al., 1993). Other neurotransmitter and neurochemical markers of JG cells include: (1) enkephalin (ENK) (Bogan et al., 1982; Davis et al., 1982; Kosaka et al., 1987d), (2) thyrotropin-releasing hormone (TRH), (Merchenthaler et al., 1988; Tsuruo et al., 1988), (3) acetylcholinesterase (AChE), a marker for cholinceptive neurons (Nickell and Shipley, 1988b), (4) cholecystokinin (CCK) (Seroogy et al., 1985; Matsutani et al., 1988), (5) aspartic acid (Watanabe and Kawana, 1984; Halász, 1987; Fuller and Price, 1988), (6) vasoactive intestinal polypeptide (VIP) (Gall et al., 1986; Sanides-Kohlrausch and Wahle, 1990a), (7) protein kinase C (Saito et al., 1988) and two Ca^{2+} binding proteins, (8) calretinin (CR) (Jacobowitz and Winsky, 1991; Rogers, 1992), and (9) calbindin-d28k (CB) (Baimbridge and Miller, 1982; Celio, 1990; Briñón et al., 1992).

Three-dimensional analyses using serial-sectioning/mirror-imaging and immunocytochemical double-labeling techniques revealed that about half of the PG cell population can be classified into at least three cell types, including: (1) GABA-positive, (2) CR-positive, and (3) CB-positive (Kosaka et al., 1995, 1998). Moreover, the DA, TRH, and ENK neurons were identified as further subpopulations. About 80% of the DA cells contained GABA in the rat MOB, and in turn, corresponded to about 50% of GABAergic cells (Kosaka et al., 1985, 1987a, c, 1988, 1995; Gall et al., 1987). Therefore, the DA PG neurons are regarded as a subpopulation of GABAergic neurons. Clearly, the neurochemical diversity of JG neurons is remarkable and at present, the role of these neuronal subtypes and their neurochemical constituents in olfactory processing is unclear.

3.2.3 Synaptic and Bicompartamental Organization of the GL

3.2.3.1 Excitatory Systems in the GL In addition to ON axons, each glomerulus contains the apical-dendritic tufts of about 20 mitral cells, 200 tufted cells and 1500–2000 JG cells (Shipley et al., 1996).

Sensory transmission from ON axon terminals is mediated by glutamate acting at AMPA and NMDA ionotropic glutamate receptor subtypes (Bardoni et al., 1996; Ennis et al., 1996, 2001; Aroniadou-Anderjaska et al., 1997; Chen and Shepherd, 1997; Keller et al., 1998). The distribution of these receptors in MOB is discussed later (▶ [Section 3.8](#) and ▶ [Table 6-3](#)). Electrophysiological studies, described later,

■ **Table 6-3**
Neurotransmitter receptors in the MOB

Receptor	GL	EPL	MCL	IPL	GCL	References
Cholinergic						Fonseca et al. (1991a, b), Le Jeune et al. (1996), Rotter et al. (1979), Sahin et al. (1992), Spencer et al. (1986)
M1	-	++	-	++	++	
M2	++	++	++	++	++	
M3	-	++	-	++	-	
M4	-	++	-	++	-	
Nicotinic	++	++	++	-	++	
Noradrenergic						Booze et al. (1989), Day et al. (1997), Domyancic and Morilak (1997), McCune et al. (1993), Nicholas et al. (1993), Palacios and Kuhar (1980), Pieribone et al. (1994), Rosin et al. (1996), Sargent-Jones et al. (1985), Talley et al. (1996), Wanaka et al. (1989), Winzer-Serhan et al. (1997a), Woo and Leon (1995), Young and Kuhar (1980a), Yuan et al. (2003)
$\alpha 1$	++	++	++	++	++	
$\alpha 2$	++	++	++	++	++	
$\beta 1$	++	++	++	-	++	
$\beta 2$	++	-	-	++	++	
Dopaminergic						Coronas et al. (1997), Koster et al. (1999), Mansour et al. (1990a, b), Nickell et al. (1991)
D1	++	-	-	-	-	
D2	++	-	-	-	++	
Serotonergic						Clemett et al. (2000), Cornea-Hebert et al. (1999), Hamada et al. (1998), McLean et al. (1995), Pompeiano et al. (1992, 1994), Tecott et al. (1993), Whitaker-Azmitia et al. (1993), Wright et al. (1995), Yuan et al. (2003)
5-HT _{1A}	++	++	++	++	++	
5-HT _{1C}	-	-	-	-	++	
5-HT _{2A/C}	++	++	++	NR	++	
5-HT ₃	++	-	-	-	-	
Glutamatergic						Duvoisin et al. (1995), Gall et al. (1990), Giustetto et al. (1997), Hamilton and Coppola (2003), Kinoshita et al. (1996, 1998), Kinzie et al. (1995, 1997), Martin et al. (1992, 1993), Masu et al. (1991), Miller et al. (1990), Molnar et al. (1993), Monaghan and Cotman (1982), Montague and Greer (1999), Monyer et al. (1994), Ohishi et al. (1993a, b, 1995, 1998), Petralia and Wenthold (1992), Petralia et al. (1994a, b), Romano et al. (1995), Sahara et al. (2001), Sassoè-Pognetto and Ottersen (2000), Saugstad et al. (1997), Shigemoto et al. (1992, 1993), Sun et al. (2000), Tanabe et al. (1992), Van Den Pol (1995), Wada et al. (1998), Watanabe et al. (1993), Wisden and Seeburg (1993)

■ **Table 6-3 (continued)**

Receptor	GL	EPL	MCL	IPL	GCL	References
Kainate	–	++	++	++	–	
NMDA	++	++	++	++	++	
AMPA	++	++	++	++	++	
mGlu	++	++	++	++	++	
GABAergic						Bonino et al. (1999), Bowery et al. (1987), Chu et al. (1990), Fritschy et al. (1992), Laurie et al. (1992), Margeta-Mitrovic et al. (1999), Persohn et al. (1992), Richards et al. (1987)
GABA _A	++	++	++	–	++	
GABA _B	++	–	++	NR	++	
CRF	NR	++	++	NR	++	Chen et al. (2005), Van Pett et al. (2000)

++, receptors present; –, receptors absent; NR, not reported. Adapted from Handbook of Chem. Neuroanat. Integrated Sys. CNS, Vol. 12, Part III, Chapter III, The Olfactory System, M. Shipley, pp. 469–573, 1996, Elsevier, Ltd

have shown that mitral/tufted cells and ET cells are excited by these synapses. ON axon terminals synapse with some subclasses of PG cells, but not with others, and that they do not synapse with SA cells (Hayar et al., 2004a, b; for review see Kosaka et al., 1998). As will also be discussed later, mitral/tufted cells are glutamatergic, and dendritic release of glutamate from these cells has been reported to produce synaptically and nonsynaptically mediated excitation of neighboring cells (i.e., glutamate spillover).


Recent *in vitro* electrophysiological findings have also provided a new insight into the roles of the different types of JG neurons in olfactory processing (Hayar et al., 2004a, b). Notably, these studies have shown that the spontaneous and sensory-evoked activity patterns of ET cells differ markedly from those of SA and PG cells. ET cells spontaneously generate rhythmic spike bursts and receive monosynaptic ON input, as has also been reported for JG cells *in vivo* (Getchell and Shepherd, 1975; Wellis and Scott, 1990). SA and PG cells, by contrast, have relatively low levels of spontaneous spike activity, do not possess the intrinsic capacity to generate spike bursts, and mostly respond di- or polysynaptically to ON stimuli. Some dopaminergic PG cells exhibit intrinsic pacemaking activity however (Puopolo et al., 2005). Thus, most SA and PG cells do not appear to receive direct ON input, perhaps because their dendrites ramify in glomerular compartments devoid of ON terminals. Anatomical studies have revealed that the glomeruli have a bicompartmental organization. Each glomerulus has several interdigitating compartments, one of which is rich in ON terminals and the second devoid of ON input (Kosaka et al., 1997; Kasowski et al., 1999). Calbindin-positive JG neurons extend their dendrites only into glomerular compartments devoid of ON terminals, suggesting that they do not receive direct sensory innervation (Toida et al., 1998, 2000). It is reasonable to speculate, therefore, that the SA and PG cells that do not receive direct input (Hayar et al., 2004b) may correspond to these calbindin-positive glomerular interneurons. If most PG cells and all SA cells are devoid of ON input, what circuit mediates their polysynaptic responses to ON stimulation? Recent studies have shown that ET cells, which respond monosynaptically to ON stimulation, in turn relay this input to PG and SA cells (Hayar et al., 2004b). These studies confirmed that ET cells provide direct excitatory glutamatergic input to PG and SA cells. Notably, this microcircuit is intra- versus interglomerular in nature. Mitral/tufted cell dendrites are also likely to provide intraglomerular excitatory input to the PG and SA cells based on anatomical grounds, but this has not been confirmed electrophysiologically (🔗 [Figures 6-6](#)).

3.2.3.2 Inhibitory Systems in the GL

Dendrodendritic Inhibition As noted earlier, PG cells contain GABA and they form inhibitory synapses onto mitral/tufted cell dendrites in the glomeruli. In the periglomerular spaces, their axons also form symmetrical synapses onto the mitral/tufted cell dendrites and PG and SA cells (Pinching and Powell, 1971c). Physiological studies support the notion that PG cells receive excitatory input from mitral/tufted cells and ET cells, and in return, make feedforward and feedback inhibitory GABAergic synapses onto these

■ Figure 6-6

Intra- and interglomerular circuitry. Glutamatergic ET cells rhythmically burst at sniffing frequencies and receive monosynaptic ON input. ET cells of the same glomerulus burst synchronously and trigger monosynaptic bursts of EPSPs in other JG interneurons (PG and SA cells), most of which do not receive direct ON input. ET cells are synchronized by ON input, glutamate spillover, and interconnections via gap junctions (ovals). ET cells are also synchronized by IPSP bursts from the same and different sets of PG cells. PG cell dendrites ramify in a restricted portion of a single glomerulus and provide local intraglomerular inhibition of ET cells via dendrodendritic interactions, and also exert presynaptic inhibition of ON terminals. By contrast, SA cells have dendrites and axons extending throughout several glomeruli, and thus might subserve interglomerular interactions (i.e., lateral inhibition). The synchronously bursting ET cell glomerular “ensemble” may constitute an oscillating rhythm generator that monosynaptically synchronizes the activity of PG and SA cells within the same glomerulus, and perhaps also coordinates the activity of mitral/tufted cells via glutamate spillover



cells (Shepherd and Greer, 1998; Hayar et al., 2004b). Such inhibition is thought to be primarily mediated by GABA_A receptors (see ▶ Section 3.8.3). Because PG cell dendrites ramify within a restricted portion of a glomerulus, their inhibition is presumably localized to microdomains of the extensive mitral/tufted cell dendritic arbors or to nearby JG cells (Kasowski et al., 1999). By contrast to PG cells, SA cells extend dendrites and axons across multiple glomeruli, suggesting that these cells mediate interglomerular functions, such as lateral inhibition of output neurons in neighboring glomeruli (Aungst et al., 2003). The latter study suggests the intriguing possibility that SA cells are excitatory. The possible interactions among ET, PG, and SA cells are illustrated in a simplified network model in ▶ Figure 6-6. Detailed EM-immunocytochemical work from the Kosaka and Toida laboratories has deduced the synaptic interactions of subclasses of GABA-, DA-, and CB-positive PG neurons with mitral/tufted cell dendrites. A review of this work is beyond the scope of this chapter and interested readers are referred to the original publications (Toida et al., 1998, 2000; Kosaka et al., 2001). PG cells and ET cells also express GABA_A receptors (see ▶ Section 3.8.3), and there is electrophysiological evidence that some PG cells can release GABA onto themselves, and perhaps neighboring PG cells, in certain circumstances. This occurs through Ca²⁺-dependent release of GABA (Smith and Jahr, 2002). GABA has been reported to depolarize PG cells at their resting potential, probably due to the elevated intracellular chloride concentrations (Siklos et al., 1995; Smith and Jahr, 2002). It was suggested that GABA inhibits PG cells by activating a chloride conductance that reduces the neuronal input resistance and shunts excitatory inputs. GABA released from PG cells was recently shown to inhibit other PG cells in the same glomerulus via GABA_A receptors (Murphy et al., 2005).

Presynaptic Inhibition of ON Axon Terminals

Dopamine and D2 receptors The GL contains several hundred thousand DA neurons, but the MOB receives no known extrinsic DA input. In mammals, D1 receptor mRNA is expressed in the GL and GCL (Coronas et al., 1997). Immunocytochemical localization of D1 receptors is faint and primarily in the GCL (Levey et al., 1993). By contrast, D1-like ligand binding is present at very low levels in all layers of the MOB

with the exception of the ONL (Mansour et al., 1990a; Nickell et al., 1991; Coronas et al., 1997). The functional significance of D1 receptors in MOB remains unclear. By contrast, only the ONL and GL have high densities of D2 receptors in rats and mice (Mansour et al., 1990a; Nickell et al., 1991; Levey et al., 1993; Coronas et al., 1997; Koster et al., 1999) and to a lesser extent in guinea pigs, but not in cats or monkeys (Camps et al., 1990). In the GL, the JG neurons express D2 receptors (Mansour et al., 1990a). Some immunocytochemical labeling for D2 receptors, as well as in situ hybridization, has been reported in the GCL and EPL, however D2 binding sites are consistently restricted to the ONL and GL (Mansour et al., 1990a; Levey et al., 1993; Coronas et al., 1997). Other anatomical evidence indicates that most, if not all, of the D2 receptors in the GL occur on ON axon terminals. ORNs express D2 receptors and bulbectomy, a manipulation that causes retrograde degeneration of ORNs, eliminates D2 receptor mRNA in the olfactory epithelium (Koster et al., 1999). Taken together, these findings indicate that DA released from JG neurons may presynaptically modulate ON terminals via activation of D2 receptors.

In agreement with this, DA and D2 receptor agonists reduced spontaneous and ON-evoked activity in mitral cells, as well as odor-evoked activity in the GL and odor detection performance, in a variety of species (Nowycky et al., 1983; Doty and Risser, 1989; Sallaz and Jourdan, 1992; Wachowiak and Cohen, 1999; Hsia et al., 1999; Berkowicz and Trombley, 2000; Ennis et al., 2001). These effects are mediated by presynaptic suppression of glutamate release from ON terminals via inhibition of Ca^{2+} influx (Wachowiak and Cohen, 1999). In a similar manner, DA and D2 receptor agonists suppressed spontaneous and ON-evoked activity in JG cells, but had no effect on mitral-to-JG cell transmission (Ennis et al., 2001). The inhibitory effects of DA were abolished in D2 receptor knockout mice (Ennis et al., 2001). Intriguingly, this presynaptic regulation occurs in the absence of conventional anatomical synapses from JG cells to ON terminals. The close proximity of the dendrites of DA-positive and GABA-positive PG cells to ON terminals may facilitate this “nonsynaptic” presynaptic regulation. Further, synaptic vesicles are present in some dendrites opposed to ON axon terminals (Bonino et al., 1999). What role might such presynaptic inhibition serve? One possibility is that presynaptic inhibition of ON terminals by DA provides a mechanism for increasing the range of concentrations that can be processed by MOB neurons: as activity increases in ON terminals, dopaminergic JG cells are more strongly excited. This in turn provides negative feedback to ON terminals reducing the release of glutamate. Such a scheme would effectively increase the dynamic range of information transfer from ORNs to MOB neurons. Interestingly, systemic administration of D2 receptor agonists has been reported to prevent odorant specific 2-deoxyglucose patterns in MOB and to reduce odorant detectability (Doty and Risser, 1989; Sallaz and Jourdan, 1992). Related to this question of how DA participates in odor processing is the degree to which these receptors are tonically active in vivo? If, for example, ON terminals are tonically inhibited by DA via D2 receptors, this might serve to filter out weak signals (“noise”). This might sharpen the spatial pattern of active glomeruli, and facilitate the detection of predominant odors. There is experimental support for this possibility. Blockade of D2-like receptors by systemic administration of spiperone increased the number of mitral cells that responded to single or multiple odorants (Wilson and Sullivan, 1995). One interpretation of this study is that reduced D2 presynaptic inhibition of ON terminals increases the odor responsiveness of mitral cells, but does so at the cost of reduced odorant discrimination. Other effects of both D1 and D2 receptor activation have been reported (Duchamp-Viret et al., 1997; Brünig et al., 1999; Davison et al., 2004).

GABA, GABA_B Receptors, and Taurine GABA_B receptors play a presynaptic inhibitory role, apparently very similar to that described for D2 receptors. As noted earlier, GABAergic PG cells represent a large population of GL interneurons. In the rat MOB, the glomeruli have the highest concentration of GABA_B receptors as determined by radioligand binding (Bowery et al., 1987; Chu et al., 1990) and immunohistochemical localization of GABA_B receptor subunits (Margeta-Mitrovic et al., 1999). EM-immunohistochemistry revealed that the dense labeling in the GL is due to the presence of GABA_B receptors on ON terminals and the somata of PG cells (Table 6-4) (Bonino et al., 1999). A variety of imaging and electrophysiological studies have provided solid evidence that GABA released from PG neurons presynaptically inhibits glutamate release from ON terminals via activation of these GABA_B receptors (Keller et al., 1998; Wachowiak and Cohen, 1999; Aroniadou-Anderjaska et al., 2000; Palouzier-Paulignan et al., 2002; Murphy et al., 2005). The MOB also contains the highest levels of the putative inhibitory transmitter

■ Table 6-4

Gap junctions and connexins in the MOB

Layer/Cell Type	Gap Junctions ^a	Connexin mRNA ^b	Connexin Protein ^c	References
MOB			C×36	Rash et al. (2000)
ONL			C×36, 43	Paternostro et al. (1995), Reyher et al. (1991), Teubner et al. (2000), Zhang and Restrepo (2003)
GL			C×36, 43	Belluardo et al. (2000), Christie et al. (2005), Paternostro et al. (1995), Theis et al. (2003), Teubner et al. (2000), Zhang and Restrepo (2003)
JG cells		C×36		Christie et al. (2005)
PG cells		C×36		Kosaka and Kosaka (2003, 2004), Landis et al. (1974)
M/T-PG dendrites	TEM			Kosaka and Kosaka (2003, 2005)
M/T-M/T dendrites	TEM			Christie et al. (2005), Kosaka and Kosaka (2004, 2005), Rash et al. (2005)
M/T-unidentified dendrites	TEM			Kosaka and Kosaka (2003, 2005), Kosaka et al. (2005)
EPL			C×43	Reyher et al. (1991)
Mitral cell proximal dendrites			C×45	Teubner et al. (2000)
Tufted cells (and/or interneurons?)		C×36, 43		Condorelli et al. (1998, 2000), Miragall et al. (1996)
Unidentified cells			C×43	Theis et al. (2003)
Tufted cells-Interneurons	TEM			Kosaka and Kosaka (2003)
M/T-Interneuron dendrites	TEM			Kosaka and Kosaka (2003)
M/T-Unidentified dendrites	TEM			Kosaka and Kosaka (2003)
M/T-GC dendrites	FF			Landis et al. (1974)
EPL/MCL		C×36	C×36, 43	Belluardo et al. (2000)
MCL		C×36	C×36, 43	Belluardo et al. (2000), Christie et al. (2005), Condorelli et al. (2000), Reyher et al. (1991)
Mitral cells	FF	C×36, 43, 45	C×45	Belluardo et al. (2000), Condorelli et al. (1998), Miragall et al. (1996), Paternostro et al. (1995), Theis et al. (2003), Teubner et al. (2000), Zhang and Restrepo (2002)
IPL			C×43	Reyher et al. (1991), Kosaka et al. (2005), Rash et al. (2005)
GCL				
GCs	TEM, FF	C×36	C×43	Christie et al., (2005), Condorelli et al. (1998), Reyher et al. (1991), Kosaka et al. (2005)
Unidentified cells			C×43	Theis (2003)

^aStandard transmission electron microscopy (TEM) showing junctions between two cell types or freeze-fracture (FF) replica analysis or intramembranous particles

^bIn situ hybridization or nuclease protection assay

^cWestern blots (entire MOB, presumably including AOB), immunocytochemistry (layers) or immunohistochemistry (cells)
Abbreviations: M/T, mitral/tufted

taurine, exceeding concentrations of GABA and glutamate (Collins, 1974; Margolis, 1974; Banay-Schwartz et al., 1989a, b; Ross et al., 1995; Kamisaki et al., 1996). Taurine is found in ON axons, in various neurons, and in astrocytes (Kratskin et al., 2000; Kratskin and Belluzzi, 2002; Pow et al., 2002). In the ON terminals and some postsynaptic dendrites, taurine is colocalized with glutamate (Didier et al., 1994). Observation of spontaneous taurine release from MOB synaptosomes suggests that taurine may be abundantly released (Kamisaki et al., 1996). In electrophysiological recordings, taurine directly activated presynaptic GABA_B receptors and inhibited ON terminals, and also induced Cl⁻ currents in mitral/tufted cells. Surprisingly, taurine had no direct effect on PG cells (Belluzzi et al., 2003).

3.2.3.3 Neuromodulation in the GL Neuromodulatory systems in the GL include carnosine and certain heavy metals (copper and zinc), metabotropic glutamate receptors (mGluRs), and neuromodulatory inputs from centrifugal afferents. mGluRs and centrifugal inputs are discussed later (▶ [Sections 3.8.2](#) and ▶ [3.9](#)). Carnosine, a dipeptide synthesized by ORNs, is localized in ON terminals in the GL and fulfils many criteria for neurotransmitter candidacy. However, no direct postsynaptic actions of carnosine have been revealed to date (MacLeod and Straughan, 1979; Nicoll and Alger, 1980; Frosch and Dichter, 1982; Trombley et al., 1998). Carnosine did not affect currents evoked by glutamate, GABA, or glycine in cultured MOB neurons (Trombley et al., 1998). Because carnosine is a chelator of both zinc and copper, it has been suggested that it might modulate transmission at ON synapses by regulating zinc and copper. Depending on the concentration, both zinc and copper can augment or block responses mediated by NMDA and GABA receptors. Both zinc and copper inhibit NMDA and GABA receptor-mediated currents and synaptic transmission in MOB neurons (Trombley and Shepherd, 1996; Trombley et al., 1998). Carnosine prevented the action of copper and reduced the effect of zinc. These results suggest that carnosine may indirectly influence glutamate actions on MOB neurons by modulating the effects of synaptically released zinc and copper. What roles might carnosine, zinc and copper play in olfactory processing? At present, there are no clear answers to this question. Is carnosine, like other peptide transmitters, preferentially released by high frequencies of ON activity? If so, carnosine may be preferentially released during intense odor stimulation. Zinc is known to be preferentially released during high frequency neural activity. There are neurotoxic effects of zinc and copper, and Trombley (Horning et al., 2000) has speculated that carnosine, by preventing the actions of these metals, may serve an important neuroprotective function, perhaps to protect MOB neurons during intense or high frequency activity.

3.3 External Plexiform Layer

The EPL lies beneath or deep to the glomeruli, and it primarily consists of dense neuropil formed by the dendrites of mitral cells and GCs that ascend from the MCL and GCL, respectively. Relative to other MOB layers, the EPL has a low cell density. In Nissl-stained sections, however, it can nevertheless be seen to contain significant numbers of neurons (▶ [Figure 6-3](#)). These include several subtypes of tufted cells and intrinsic interneurons, which are described later. Because tufted cells are in many aspects similar to mitral cells, and as mitral and tufted cell dendrites cannot be distinguished ultrastructurally, the term mitral/tufted cell is often used when generalizing to these two cell populations. The dominant feature of the EPL is nevertheless the extensive dendrodendritic synapses between mitral/tufted cells and GCs.

3.3.1 Neuron Types of the EPL

3.3.1.1 Tufted Cells Tufted cells are the most numerous cells of the EPL. Like ET cells and mitral cells, tufted cells of the mammalian EPL have one (or rarely several) apical (primary) dendrite(s) ending in a glomerular tuft, hence the name (Cajal, 1890) (▶ [Figure 6-5](#)). The size of the tufted cell bodies gradually increases from the superficial to the deep EPL (Cajal, 1890; Pinching and Powell, 1971a; Macrides and Schneider, 1982; Orona et al., 1984). Golgi staining and dye injection studies (▶ [Figure 6-5](#)) have shown that superficial tufted cells tend to have asymmetrical branching patterns, extending sparsely branched

lateral dendrites obliquely to the projection path of the primary dendrites (Macrides and Schneider, 1982; Orona et al., 1984). Middle tufted cells, deep tufted cells, and mitral cells exhibit more symmetrical branching patterns (i.e., shaped like a bishop's miter), with one apical dendrite and tangentially arrayed lateral dendrites (▶ [Figure 6-5](#)). Tufted cells that closely resemble mitral cells in size have also been referred to as displaced mitral cells (Blanes, 1898), but this term is now generally reserved for the deepest tufted cells located near the MCL (Mori et al., 1983; Kiyoshi et al., 1984; Mori, 1987a, b). Tufted cells utilize glutamate as their principle transmitter (Liu et al., 1989; Christie et al., 2001), but they also stain for a number of other substances (▶ [Table 6-1](#)). Many superficial tufted cells contain CCK (Seroogy et al., 1985), and they form a topographically organized, reciprocal network (Schoenfeld et al., 1985), which is described in ▶ [Section 3.3.2.4](#). In the Chinese hamster, many superficial and middle tufted cells are immunoreactive for substance P. The rat has some of these cells, but in other rodents, few if any occur (Baker et al., 1986; Matsutani et al., 1988). The deep EPL of the Syrian hamster has a few NADPH diaphorase-positive tufted cells (Davis, 1991), and it also has a relatively large population of dopaminergic tufted cells (Halász et al., 1981; Baker, 1986; Gall et al., 1987). The latter cells are relatively rare in other rodent species, but are observed in primates (Smith et al., 1991) and amphibians (Boyd and Delaney, 2002). TH-immunoreactive tufted cells do not coexpress GABA, unlike TH-positive PG cells (Gall et al., 1987). As discussed in ▶ [Section 3.4](#), many mitral/tufted cells are also immunoreactive for corticotropin-releasing factor (CRF) (Imaki et al., 1989; Bassett et al., 1992), and the rodent EPL exhibits a high density of CRF receptors (De Souza et al., 1985).

Axonal projections of middle and deep tufted cells are similar, but not identical, to those of mitral cells (Schoenfeld and Macrides, 1984; Schoenfeld et al., 1985). The local axon collaterals of tufted cells course mainly in the IPL. The projections of tufted cells beyond the MOB terminate densely in the AON and, to a lesser extent, in other rostral olfactory cortical structures (Schoenfeld et al., 1985; Scott, 1986). Few tufted cell axons project into more caudal POC regions.

3.3.1.2 EPL Interneurons The EPL contains anatomically and neurochemically heterogeneous subtypes of intrinsic interneurons. From a functional standpoint, these cells are not well understood. More information on their synaptic interactions is clearly needed to understand the role of these cells in olfactory processing. Their interactions with tufted cells may be particularly important, because tufted cells are thought to comprise a parallel “channel” for transmitting most of the activity from the MOB, separately from the mitral cells (Shepherd et al., 2004). With Golgi staining methods, the EPL was originally described in the cat as containing multipolar neurons having multiple dendrites originating from two poles, but no axons (Van Gehuchten and Martin, 1891). Other interneuron subtypes were subsequently described in other species (for a review, see Halász, 1990). These subtypes include bipolar interneurons with long processes arrayed tangentially to the MCL (horizontal cells) (Valverde, 1965) and multipolar neurons with varicose dendrites, referred to as Van Gehuchten (VG) cells, which occur together with superficial short axon (SSA) cells resembling those of the GL (▶ [Figure 6-5](#)) (Schneider and Macrides, 1978). VG cells are characterized by several relatively thick primary dendrites that remain in the EPL and axons terminating around mitral and tufted cells. VG cells with smooth dendrites, horizontal cells thought to be equivalent to SSA cells, and small satellite cells with varicose dendrites that envelop small segments of mitral cell dendrites, have been described (López-Mascaraque et al., 1990). Other VG and SA cells typically have varicose dendrites that branch within either the superficial or deep half of the EPL. With parvalbumin (PV) immunostaining however, interneurons with dendrites that are either varicose or relatively smooth have been observed, and some of these interneurons exhibit intermediate branching patterns (Kosaka et al., 1994a, b). Many EPL interneurons stain for GABA (Mugnaini et al., 1984a; Gall et al., 1987; Kosaka et al., 1987d), including some of those identified as VG and SSA cells (Ohm et al., 1990), and are therefore presumed to be inhibitory. Some of the GABA interneurons are also immunoreactive for PV (Kosaka et al., 1987d). As summarized in ▶ [Tables 6-1](#) and ▶ [6-2](#), EPL interneuron subtypes are neurochemically diverse and stain for (1) NADPH diaphorase (Scott et al., 1987; Villalba et al., 1989; Alonso et al., 1995), (2) AChE (Le Jeune and Jourdan, 1994), (3) SP (Baker, 1986; Wahle et al., 1990), (4) ENK (Bogan et al., 1982; Davis et al., 1982), (5) NPY (Gall et al., 1986; Scott et al., 1987; Seroogy et al., 1989; Sanides-Kohlrausch and Wahle, 1990a), (6 and 7) neurotensin and somatostatin (Matsutani et al., 1988), (8) VIP (Gall et al., 1986; López-Mascaraque et al., 1989;

Sanides-Kohlrausch and Wahle, 1990b; Nakajima et al., 1996), (9) PV (Cielo, 1990; Ohm et al., 1990; Kosaka et al., 1994a, b; Alonso et al., 1995, 1998; Kakuta et al., 1998; Briñón et al., 1999, 2001; Jia and Halpern, 2004), (10) CB, (11) neurocalcin, and (12) CR (Ohm et al., 1991; Briñón et al., 1992, 1999, 2001; Alonso et al., 1995, 1998; Jia and Halpern, 2004).

3.3.2 Circuitry, Synaptology, and Sublaminar Organization of the EPL

3.3.2.1 Overview of EPL Circuitry and Reciprocal Mitral/Tufted Cell—Granule Cell Dendrodendritic Interactions The neuropil of the EPL contains the vertically oriented apical dendrites of mitral/tufted cells ascending to receive monosynaptic ON input in the GL, and laterally oriented secondary dendrites of the mitral/tufted cells. The majority of the synapses of the EPL are between the lateral dendrites of mitral/tufted cells and the dendrites of GCs. Most of the synapses are: (1) asymmetrical synapses from the mitral/tufted cell lateral dendrites onto GC dendrites, and (2) symmetrical synapses from the spines (“gemmules”) of GC dendrites onto the mitral/tufted cell lateral dendrites (Price and Powell, 1970b–d). These synapses are mostly reciprocal (Hirata, 1964; Rall et al., 1966; Price and Powell, 1970d; Woolf et al., 1991), and therefore occur in roughly equal proportion (Jackowski et al., 1978). Similar synapses are formed by mitral/tufted cell bodies and GC dendrites. The mitral/tufted cell lateral dendrites are tangentially arrayed relative to the MOB layers and can extend up to ~1–2 mm from the soma (Mori et al., 1983). One mitral/tufted cell can therefore receive feedback inhibition and lateral inhibition mediated by GCs that are excited by other mitral/tufted cells. As detailed knowledge of the functions of these synapses has been derived from studies of mitral cells, they are described in greater detail in [Section 3.4.1.1](#). The mitral/tufted cell dendrites also form synapses with intrinsic EPL interneurons. Through the mitral/tufted cells, these EPL interneurons may therefore indirectly receive input related to the overlying glomeruli (Hamilton et al., 2005). The third element of the EPL is an intrabulbar association system (IAS), and the fourth element is the centrifugal afferent input to the EPL from the POC. The interneuron synapses, IAS, and centrifugal inputs are also described separately later.

3.3.2.2 Synaptology of EPL Tufted Cells Tufted cells are, in many respects, similar to mitral cells in terms of synaptology. The dendritic tufts receive monosynaptic input from ON axon terminals and the lateral dendrites synapse with JG cells in the GL and with GC dendrites in the EPL (Shepherd, 1972; Shepherd et al., 2004). Like mitral cells ([see Section 3.4](#)), tufted-to-GC synaptic interactions involve a strong NMDA receptor component (Christie et al., 2001). The lateral extent of tufted cell inhibition is limited to several glomerular widths, however, ~400 μm as opposed to 750 μm for mitral cells in rats (Christie et al., 2001). The weaker lateral inhibition of tufted cells could merely be due to the fact that their lateral dendrites are relatively short (Mori et al., 1983; Orona et al., 1984). However, recent evidence suggests that inhibition from the intrinsic EPL interneurons, not from GCs, tunes the responses of middle tufted cells to odor stimulation (Nagayama et al., 2004).

3.3.2.3 Synaptology of EPL Interneurons The EPL interneurons were originally thought to interact chiefly with GC dendrites (Schneider and Macrides, 1978). However, more recent studies have suggested that they instead interact with mitral/tufted cell dendrites (López-Mascaraque et al., 1990; Nagayama et al., 2004), a view that is supported by ultrastructural studies. Ultrastructural studies have shown that the PV-IR interneurons of the rat EPL receive asymmetrical synapses from mitral/tufted cell bodies and dendrites, and form symmetrical synapses onto the mitral/tufted cells, many of which are reciprocal (Toida et al., 1994, 1996; Crespo et al., 2001). Surprisingly, although the PV-IR interneurons were observed to form frequent contacts with each other, no synapses were observed at the contact points. However, symmetrical synapses from unidentified PV-immunonegative axons were observed on the PV-immunoreactive cell bodies, where they would be expected to shunt the more distal mitral/tufted cell inputs. The PV-immunoreactive interneurons also receive asymmetrical synapses from unidentified axons (Toida et al., 1996), which could be centrifugal afferents. Synapses from centrifugal fibers have been observed on SA cells (Pinching and Powell, 1972a–c). Combined morphological and electrophysiological analyses showed that multipolar

EPL interneurons with highly varicose dendrites are excited polysynaptically by ON stimulation, most likely via input from mitral/tufted cells (Hamilton et al., 2005). The multipolar interneurons exhibit robust spontaneous excitatory synaptic activity mediated by AMPA/kainate receptors, consistent with immunocytochemical staining studies showing that mouse EPL neurons are strongly reactive to the GluR1 AMPA receptor subunit (Petralia and Wenthold, 1992; Giustetto et al., 1997; Montague and Greer, 1999; Hamilton and Coppola, 2003). Similar electrophysiological recordings were obtained from varicose multipolar interneurons in slices from normal mice and from fluorescent cells exhibiting the same morphology in slices from transgenic mice labeled to reveal GABAergic interneurons. The dendrites of the interneurons that were located in the superficial EPL bridged the space below several adjacent glomeruli (Figure 6-5). Interneurons located in the deeper zones bridged an equivalent EPL area, which was approximately equal to the glomerular width (Hamilton et al., 2005). This suggests that multipolar EPL interneurons are excited by mitral/tufted cells and may in turn inhibit mitral/tufted cells within EPL domains that are topographically related to pairs of overlying glomeruli.

3.3.2.4 Sublaminar Organization of the EPL As noted earlier, superficial, middle, and deep tufted cells are distinguished by cell body location and size (Cajal, 1890; Pinching and Powell, 1971a; Macrides and Schneider, 1982; Orona et al., 1983). The lateral dendrites of the superficial, middle, and deep tufted cells course within superficial, intermediate, and deep EPL zones, respectively (Scott and Harrison, 1987). Type I and II mitral cell subtypes have also been identified according to the distribution of their lateral dendrites within the deep and intermediate zones, respectively (Kiyoshi et al., 1982; Orona et al., 1984). Electrophysiological recordings have shown that the sensitivity of these tufted cell and mitral cell subtypes to ON stimulation is correlated with the depth of the lateral dendrites within the EPL. The more superficial tufted cells are more easily excited by ON stimulation than the deeper tufted cells, and the type II mitral cells are more easily excited than type I mitral cells (Schneider and Scott, 1983; Wellis et al., 1989; Ezeh et al., 1993). In response to odor stimulation, superficially located tufted cells also exhibit more prolonged excitation than mitral cells (Luo and Katz, 2001). Thus, anatomically distinct mitral/tufted cell subtypes appear to be functionally distinct.

A differential sublaminar distribution is also observed for the dendrites of superficial and deep GCs. The apical dendrites of superficial GCs have very dense spines, and they terminate within both the superficial and deep portions of the EPL. By contrast, apical dendrites of deeper GCs terminate exclusively within the deep EPL (Orona et al., 1983; Greer, 1987). The dendrites of a third GC subtype produce spines predominantly within the superficial EPL (Mori, 1987a, b). Perhaps as a result of these morphological differences, the superficial EPL stains more strongly for glutamic acid decarboxylase (GAD) (Mugnaini et al., 1984b), for the $\alpha 3$ GABA_A receptor subunit (Panzanelli et al., 2005), and for the AMPA receptor subunit, GluR4 (Montague and Greer, 1999). Both the superficial and deep EPL stain faintly for cytochrome oxidase activity, whereas the intermediate EPL stains darkly (Mouradian and Scott, 1988). Together, these observations suggest that morphologically and neurochemically distinct subsets of mitral/tufted cells might synapse with different subsets of GCs and intrinsic interneurons within the superficial and deep EPL sublaminae (Macrides et al., 1985; Mori, 1987a, b).

3.3.2.5 Intrabulbar Association System Many superficial tufted cells do not appear to project beyond the bulb. The axons of these cells contain CCK (Seroogy et al., 1985), and they form a topographically organized, reciprocal network, the IAS, connecting lateral and medial regions of each MOB (Schoenfeld et al., 1985). The axons of the IAS travel through the EPL and MCL to the IPL, where they coalesce into tracts that travel to the opposite side of the bulb and terminate on the apical dendrites of GCs coursing through the IPL en route to the EPL (Liu and Shipley, 1994). The IAS projection exhibits a high degree of point-to-point, topographical organization. As noted in Section 2.3 earlier, the ORNs which express the same receptor project to one glomerulus in the lateral bulb and to a second glomerulus in the medial bulb (Ressler et al., 1993; Vassar et al., 1994; Mombaerts et al., 1996; Wang et al., 1998). The IAS projections from a single glomerulus preferentially target the IPL deep to the second glomerulus (Belluscio et al., 2002; Lodovichi et al., 2003). Because CCK typically causes membrane depolarization, activation of the CCK tufted cells presumably depolarizes GCs located on the opposite side of the bulb. Thus, the IAS appears to

modulate the activities of subsets of MOB neurons receiving input from ORNs expressing the same receptor, on opposite sides of the MOB.

3.4 Mitral Cell Layer, Mitral/Tufted Cell Synapses, and Neurochemistry

Deep in the EPL is the MCL, a thin layer that contains the somata of mitral cells (25–35 μm diameter, rat) as well as numerous GCs (Cajal, 1911a, b). There are $\sim 40,000$ mitral cells (Meisami, 1989) and $\sim 100,000$ GCs in the MCL (Frazier and Brunjes, 1988). Thus, mitral cells make up less than 50% of the cells in this layer. Together with tufted cells, mitral cells are the major class of output cells of the bulb. They extend a single apical dendrite through the EPL to the GL, where it arborizes extensively throughout much of a single glomerulus (Figure 6-5). There are about 25 mitral cells (and 50 tufted cells) associated with a single glomerulus (Cajal, 1911a, b; Allison, 1953). The apical dendrites are synaptically contacted by ON terminals (Price and Powell, 1970a; Shepherd, 1972). The secondary dendrites of mitral cells may extend up to 2 mm in the EPL and are oriented tangentially, i.e., parallel to the surface of the bulb. Mitral cells have been subdivided into two classes based on the organization of the lateral dendrites as reviewed in detail elsewhere (Orona et al., 1984; Shipley et al., 1996). These lateral dendrites participate in dendrodendritic synapses with dendrites of GCs, as reviewed earlier and later. In addition, they may receive centrifugal axon inputs and inputs from VG EPL interneurons (Jackowski et al., 1978; Toida et al., 1996). Mitral cells give off axon collateral which terminate, within the bulb, in the IPL and GCL (Mori et al., 1983; Price and Powell, 1970c), or exit the MOB and innervate a number of olfactory-related brain regions collectively known as the POC. Projections to POC are discussed later.

3.4.1 Neurochemistry of Mitral Cells

Mitral (and tufted) cells are glutamatergic (Liu et al., 1989; Christie et al., 2001). A few small mitral cells appear to contain aspartate and project to the largest component of POC, piriform cortex (PC) (Fuller and Price, 1988). Another proposed transmitter for mitral cells is N-acetyl-aspartyl-glutamate (NAG) (Table 6-1) based on anatomical grounds (Ffrench-Mullen et al., 1985; Blakely et al., 1987), but there has been little neurophysiological support for this notion (Whittemore and Koerner, 1989; see Section 5.3.1.1). The neuropeptide CRF has been demonstrated in mitral and some tufted cells using both immunocytochemistry and in situ hybridization in the rat (Imaki et al., 1989; Bassett et al., 1992). CRF fibers were also observed in layer Ia of the PC, consistent with the dense axonal projections of mitral cells to layer Ia of the PC. Mitral cells express type 1 CRF receptor mRNA (Van Pett et al., 2000), and mouse mitral cells also label for a soluble splice variant of the type 2 CRF receptor (Chen et al., 2005). SP mRNA transcripts have been detected in up to one-half of the rat mitral cells (Warden and Young, 1988), but to date no studies using immunocytochemistry have detected SP in mitral cells of any species (Inagaki et al., 1982; Shults et al., 1984; Baker, 1986). Finally, CR, a Ca^{2+} -binding protein, has been shown by immunohistochemistry to be localized in mitral cells (Table 6-2) (Jacobowitz and Winsky, 1991). Transmitter candidates for mitral and tufted cells are discussed further in Section 5.3.1.1.

3.4.1.1 Glutamate

ON Input As noted earlier, mitral/tufted cell apical dendrites respond to glutamate released from the ON via AMPA and NMDA receptors (Ennis et al., 1996, 2001; Aroniadou-Anderjaska et al., 1997; Chen and Shepherd 1997). The NMDA receptor-mediated response component is of unusually long duration, leading to a late spiking component in response to ON input (Ennis et al., 1996, 2001; Aroniadou-Anderjaska et al., 1997). An mGluR-mediated component of ON-evoked currents in mitral cells also occurs but is typically small under normal conditions (De Saint Jan and Westbrook, 2005; Ennis et al., 2006), yet, as will be discussed later, blockade of these receptors significantly alters ON-evoked spiking.

Self And Neighbor Excitation Electrophysiological studies indicate that mitral and tufted cells have functional ionotropic glutamate autoreceptors that mediate recurrent excitatory interactions among their

apical and lateral dendrites (Isaacson, 1999; Aroniadou-Anderjaska et al., 1999b; Carlson et al., 2000; Schoppa and Westbrook, 2001; Didier et al., 2001; Salin et al., 2001; Urban and Sakmann, 2002). Mitral cell bodies and lateral dendrites have NMDA (Isaacson, 1999; Isaacson and Murphy, 2001), AMPA, and kainate autoreceptors (Lowe, 2003). Ultrastructural studies indicate that NMDA receptors of the lateral dendrites are extrasynaptic (Sassoè-Pognetto et al., 2003). Mitral cell primary dendrites have AMPA autoreceptors (Salin et al., 2001). NMDA autoreceptors on mitral cells increase the firing frequency during prolonged discharges (Friedman and Stowbridge, 2000). Thus, glutamate released from mitral/tufted cell dendrites appears to spillover and excite glutamate receptors on the same cell and on neighboring mitral/tufted cell dendrites. Such excitatory interactions among mitral tufted cell apical dendrites in the glomeruli are thought to synchronize the discharge of mitral cells associated with the same glomerulus (Carlson et al., 2000; Schoppa and Westbrook, 2001).

mGluRs In dissociated rat and frog, MOB culture preparations, Group I mGluRs increased Ca^{2+} release from internal stores in mitral/tufted cells as well as in MOB interneurons (Geiling and Schild, 1996; Carlson et al., 1997), or it depolarized and increased the frequency of miniature excitatory postsynaptic currents (Schoppa and Westbrook, 1997). Other studies indicate that activation of Group III mGluRs with AP4 inhibits Ca^{2+} currents in mitral cells and presynaptically decreases mitral cell-to-GC synaptic transmission (Trombley and Westbrook, 1992). More recent studies in rat and mouse MOB slices demonstrate that activation of mGluR1 directly depolarizes and increases the firing of MCs, and that these effects persist in the presence of blockers of fast synaptic transmission (Heinbockel et al., 2004). The same study showed that mGluR1 induces a voltage-dependent inward current consisting of multiple components sensitive to K^+ and Ca^{2+} channel blockade and intracellular Ca^{2+} chelation. mGluR1 antagonists also altered mitral cell membrane potential bistability, increasing the duration of the up- and downstates, and substantially attenuated ON-evoked spikes. These findings suggest that endogenous glutamate tonically modulates MC excitability and responsiveness to ON input via activation of mGluR1.

Mitral/Tufted-to-JG and Mitral-to-GC Excitation Glutamate released from mitral/tufted cells, acting at ionotropic receptor subtypes, mediates dendrodendritic transmission at synapses with JG cells and with GCs (Bardoni et al., 1996; Isaacson and Stowbridge, 1998; Schoppa et al., 1998; Aroniadou-Anderjaska et al., 1999a; Chen et al., 2000; Christie et al., 2001). The mitral/tufted-to-GC cell excitation triggers GABA release from the GC dendrites (Isaacson and Stowbridge, 1998; Schoppa et al., 1998; Chen et al., 2000), which inhibits mitral cells via GABA_A receptors (Shepherd, 1972; Jahr and Nicoll, 1982; Schoppa et al., 1998; Chen et al., 2000; see [Section 3.8.3](#)). Ca^{2+} influx via the NMDA receptor channel is sufficient to trigger GABA release from GCs, and thus, to drive dendrodendritic inhibition of mitral cells (Chen et al., 2000; Halabisky et al., 2000); however, NMDA-evoked depolarization can also trigger GABA release from these cells via activation of voltage-dependent Ca^{2+} channels.

Modulation of Mitral/Tufted-to-Granule Cell Dendrodendritic Interactions mGluRs may also occur at the mitral-to-granule synapses, although their role is unclear. As noted earlier, activation of Group III receptors was reported presynaptically to decrease mitral-to-GC synaptic transmission (Trombley and Westbrook, 1992), and activation of GABA_B receptors has been observed to reduce dendrodendritic inhibition via suppression of GABA release from GCs due to reduction of high voltage-activated Ca^{2+} currents (Isaacson and Vitten, 2003). Although DA has no direct effect on resting membrane properties of mitral/tufted cells (Ennis et al., 2001), activation of D2 receptors was found to reduce glutamate release onto GCs via inhibition of N and/or P/Q high voltage-activated Ca^{2+} channels (Davila et al., 2003; Davison et al., 2004).

3.5 Internal Plexiform Layer and Granule Cell Layer

The IPL is the relatively thin layer that lies deep to the MCL. It has a low density of cells, composed mostly of dendrites derived from GCs and axons derived from mitral/tufted cells and centrifugal sources (discussed later). As noted earlier, the IPL contains a network of CCK-positive axons and terminals, which derive

from CCK-containing tufted cells as part of the IAS (Liu and Shipley, 1994). The major intrinsic cell type in the IPL appears to be the horizontal cell. Horizontal cell dendrites are oriented parallel to the MCL and horizontal cell axons extend into the EPL (Price and Powell, 1970c). Neurocalcin (Bastianelli et al., 1993; Porteros et al., 1996), CB (Briñón et al., 1992), NADPH diaphorase (Alonso et al., 1995), and ENK (Bogan et al., 1982) have been found in these cells. The IPL also contains a few multipolar neurons, which are larger than GCs and express AChE (Carson and Burd, 1980; Nickell and Shipley, 1988b; Le Jeune and Jourdan, 1994).

The GCL is the deepest neuronal layer in the bulb, and it contains the largest number of cells. Most of the neurons of the GCL are the GCs, but there are also small numbers of Golgi cells, Cajal cells, and Blanes cells. As discussed earlier, the GCs are inhibitory GABAergic cells that form dendrodendritic synapses with mitral/tufted cells in the EPL.

3.5.1 Neuron Types of the GCL

3.5.1.1 GCs GCs are axon-less cells with small cell bodies that are mostly tightly packed into row-like aggregates of 3–9 somata (Figure 6-5) (Reyher et al., 1991). GC bodies are also found mixed with mitral cell bodies within the MCL, however. Golgi studies indicate that most GCs have a thicker, longer apical dendrite that ramifies within the EPL and shorter basal dendrites that ramify within the GCL. As described earlier, several types of GCs occur, which have different dendritic ramifications within the superficial and deep portions of the EPL. Not all GCs follow this pattern, however. The exceptions include GCs with basal dendrites that project deeper, toward the center of the MOB, and GCs with apical dendrites that do not enter the EPL (Schneider and Macrides, 1978). The superficial and deep GCs can also be distinguished using a variety of staining methods. For example, superficial GCs have more dense spines on their apical dendrites and they stain for the transcription factor Er81 (Stenman et al., 2003), whereas deep GCs stain for Ca^{2+} /calmodulin-dependent kinase IV (Baker et al., 2001). The GC spines are also immunoreactive for Ca^{2+} -camodulin-dependent kinase II, expression of which is developmentally regulated. Because MOB interneurons develop postnatally and are added throughout life, the differential staining of cell bodies in the superficial versus deep GCL could be related to the migration and maturation of interneuron progenitors (see later). One function of the cholinergic input to the MOB appears to be related to the survival of these cells (Cooper-Kuhn et al., 2004), which is influenced by nicotinic receptors containing the β_2 subunit (Mechawar et al., 2004). Subpopulations of GCs are either intensely or lightly immunoreactive for CR (Jacobowitz and Winsky, 1991). GCs and some SA cells are also immunoreactive for methionine-enkephalin (Bogan et al., 1982; Davis et al., 1982), whereas deep SA cells are immunoreactive for somatostatin (Davis et al., 1982). Other possible transmitters of GCs are summarized in Table 6-1.

As discussed earlier, GCs are excited by mitral/tufted cells at asymmetrical synapses in the EPL. Most GCs contain GABA (Ribak et al., 1977), which, via GABA_A receptors, inhibits mitral/tufted cells (Shepherd, 1972; Jahr and Nicoll, 1982; Schoppa et al., 1998; Chen et al., 2000) at symmetrical synapses within the EPL. Within both the EPL and GCL, the GCs also receive asymmetrical synapses from a variety of centrifugal afferents, however, which modulate these synaptic interactions. Some of the centrifugal afferents have restricted terminations on the GC bodies and dendritic arbors, suggesting that different sources of centrifugal input could differentially modulate GC synaptic interactions with mitral/tufted cells (Price and Powell, 1970c). In large part, these centrifugal fibers arise from neurons in POC (e.g., PC and AON), which form the bulk of synaptic contacts onto GC somata and proximal dendrites within the GCL (Price and Powell, 1970c). Deep to the EPL, GCs also receive synapses from the collateral branches of mitral and tufted cell axons. During mitral cell excitation, coincident, γ -frequency stimulation of GC proximal dendrites relieves the Mg^{2+} blockade of their NMDA receptors and facilitates recurrent mitral cell inhibition (Halabisky and Stowbridge, 2003), presumably via activation of these axon collaterals. In addition, GCs receive minor inputs from Golgi, Cajal, and Blanes cells (Price and Powell, 1970c) and GCs also receive symmetrical synapses thought to be from the axons of deep SA cells, described later. The SA cells could be a source of GC inhibition as noted earlier. GABAergic centrifugal afferents from the nucleus of the horizontal limb of the diagonal band are another likely source of this inhibition (Kunze et al., 1992).

GCs express high levels of the transient outward potassium current activated by depolarization—the A-type current (Schoppa and Westbrook, 1999). This current plays an important role in dendrodendritic transmission with mitral cells in the sense that it dampens the AMPA-mediated synaptic depolarization of GCs and thereby allows NMDA receptor-mediated depolarization to play a relatively greater role. A-currents may also negatively modulate backpropagation of spikes in GC lateral dendrites (Christie and Westbrook, 2003). As noted earlier, activation of GABA_B receptors on GCs has been reported to reduce GABA release from these cells via inhibition of high voltage-activated Ca²⁺ channels (Isaacson and Vitten, 2003). GCs also receive excitatory glutamatergic projections arising from POC (see [Section 3.7.2.1](#)). GCs express type 1 and 2 CRF receptor mRNA (Van Pett et al., 2000; Chen et al., 2005).

3.5.1.2 Other Deep Interneurons The deep MOB layers contain several subtypes of SA cells (Price and Powell, 1970b; Schneider and Macrides, 1978; López-Mascaraque et al., 1990). The somata of these cells mostly occur within the IPL and GCL, and they project multiple dendrites and axons that ramify within the EPL, MCL, and GCL. The deep SA cells include horizontal cells, similar to those seen in the EPL, and Cajal cells (Cajal, 1890; Van Gehuchten and Martin, 1891; Blanes, 1898). The Cajal cells, which are also known as vertical cells of Cajal, have fusiform cell bodies that project dendrites perpendicularly rather than tangentially to the MCL. The apical dendrites extend through the MCL into the EPL, the basal dendrites extend into the GCL, and the axons project out of the GCL into the EPL (Cajal, 1911a, b). By contrast, the Blanes cells have numerous dendrites emerging from all sides of the soma. The axons from these cells can extend considerable distances, but they remain within the GCL (Cajal, 1911a, b). Larger Golgi cells (Golgi, 1875) and Blanes cells (Blanes, 1898) occur in the deeper layers. Golgi cell morphology varies considerably, but the dendrites are generally radial and the axons remain within the GCL (Cajal, 1911a, b). The dendrites of all four of these deep interneuron subtypes are varicose, but the Blanes cell dendrites are also densely spiny (Schneider and Macrides, 1978). For a detailed discussion of early literature concerning these cells, see Halász (1990).

Very little is known about the functions of the deep interneurons, but they are presumed to be inhibitory. Their axon terminals have been thought to be a source of the numerous symmetrical synapses observed on the cell bodies, basal dendrites, and spines of GCs (Price and Powell, 1970d), and some types have been shown to be immunoreactive for GABA (Gracia-Llanes et al., 2003). Recent ultrastructural evidence indicates that the VIP-containing deep short-axon cells form symmetrical synapses onto other deep interneurons containing VIP, calbindin, or NPY, but not onto GCs (Gracia-Llanes et al., 2003). Deep interneurons have also been shown to receive asymmetrical synapses that are presumed to be from mitral/tufted cell axon collaterals (Price and Powell, 1970b). It therefore appears that at least some of these interneurons inhibit other deep interneurons in response to their excitation by mitral/tufted cells. As with the more superficial interneurons, however, the deep interneurons are immunoreactive for a rich variety of substances, most notably neuropeptides ([Table 6-1](#)) and Ca²⁺ binding proteins ([Table 6-2](#)), suggesting they could serve a variety of functions. Some GCL interneurons contain NPY (Gall et al., 1986; Ohm et al., 1988). These are probably Golgi or Cajal cells, giving rise to axons that ramify in the more superficial layers such as the EPL and GL. Cajal cells stain for NADPH diaphorase, and they are immunoreactive for neurocalcin, CB, and potentially, NPY (Gall et al., 1986; Briñón et al., 1992; Bastianelli et al., 1993; Alonso et al., 1995; Porteros et al., 1996). Blanes cells also stain for NADPH diaphorase, and immunoreactivity for NPY has been observed in a population of “Blanes-like” cells near the ventricular layer (Scott et al., 1987), although the exact identity of these NADPH diaphorase/NPY cells has yet to be confirmed. Recent work shows that Blanes cells mediate persistent monosynaptic GABAergic inhibition of GCS (Pressler and Strowbridge, 2006).

3.6 Subependymal Layer and Rostral Migratory Stream

The deepest layer in the MOB is the subependymal layer (also called subventricular zone), which is a cell-poor region lining the ventricle (if present) in adults. Most MOB interneurons originate postnatally from progenitor cells within this layer (Hinds, 1968; Altman, 1969; Bayer, 1983). In adults, interneurons (primarily GCs and PG cells) are also continually generated from these progenitor cells and their offspring

generated en route migrate to the MOB within the rostral migratory stream (RMS) (Luskin, 1993; Lois and Alvarez-Buylla, 1994; Smith and Luskin, 1998; Wichterle et al., 2001). The RMS progenitors express the ETS transcription factor *Er81*, and they originate from the lateral ganglionic eminence (Wichterle et al., 2001; Stenman et al., 2003). They continue to migrate and divide in the absence of the bulb (Kirschenbaum et al., 1999), but they only reach their final destinations and mature if the GL and MCL+plexiform layers are present (Liu and Rao, 2003). Recent evidence shows that the extracellular matrix glycoprotein tenascin-R is important to this process (Saghatelian et al., 2004). Many of the offspring nevertheless die (Kaplan et al., 1985; Brunjes and Armstrong, 1996; Kato et al., 2000), with olfactory stimulation apparently being critical to the survival of new interneurons during development (Frazier-Cierpial and Brunjes, 1989; Corotto et al., 1994; Najbauer and Leon, 1995; Fiske and Brunjes, 2001) and of mature, presumably synaptically connected, interneurons in adults (Petreanu and Alvarez-Buylla, 2002; Hack et al., 2005). Two subsets of new interneurons have been identified, which express either TH, which is required for DA synthesis, or Ca^{2+} -calmodulin-dependent protein kinase IV during migration (Baker et al., 2001). The transcription factors *Pax6* and *Olig2* play important roles in determining the dopaminergic phenotype of a periglomerular subset of these interneurons (Hack et al., 2005).

Both the progenitors and new interneurons express functional GABA_A receptors. Electrophysiological studies indicate that the new interneurons subsequently express functional AMPA receptors, then NMDA receptors, before they exhibit spiking activity, responses to ON stimulation, and spontaneous glutamatergic EPSCs (Belluzzi et al., 2003; Carleton et al., 2003). These electrophysiological studies provide evidence that the new interneurons become functionally integrated into the MOB circuitry. Neural cell adhesion protein-deficient mice, which exhibit defective migration of new interneurons into the GCL resulting in reduced MOB size, have been shown to exhibit impaired odor discrimination (Gheusi et al., 2000). Although threshold detection and short-term odor memory were normal in these mice, in normal mice both olfactory memory and survival of the new interneurons were improved following rearing in an odor-enriched environment (Rocheffort et al., 2002). Thus, the neurons of the subependymal layer and RMS appear to be important both to MOB development and to certain olfactory functions during adulthood.


3.7 Outputs and Inputs of the MOB

3.7.1 Outputs of MOB

The output neurons of the MOB are mitral and tufted cells. These cells project to a number of structures collectively designated as the POC (de Olmos et al., 1978). The features of POC, including input from MOB, are described later (▶ *Figure 6-7*).

■ Figure 6-7

Major connections of the main olfactory system. Axons of MOB mitral/tufted cells (circles in the EPL and MCL, respectively) project as the LOT to synapse in a number of structures collectively referred to as primary olfactory cortex (POC). Centrifugal inputs to MOB include feedback projections from POC as well as inputs from subcortical forebrain and brainstem neuromodulatory cell groups. Abbreviations: AON, anterior olfactory nucleus; DP, dorsal peduncular cortex; Ent, entorhinal cortex; IG-AHC, indusium griseum-anterior hippocampal continuation; LC, locus coeruleus; NDB, nucleus of the diagonal band; PeCo, periamygdaloid cortex; PC, piriform cortex; RN, raphe nuclei (dorsal and median raphe); TT, taenia tecta; Tu, olfactory tubercle



3.7.2 Extrinsic Inputs to MOB

Extrinsic afferent input to MOB, also referred to as centrifugal fibers, can be subdivided into two classes: (1) inputs arising from olfactory-related structures, in particular those arising from POC, and (2) inputs arising from non-olfactory, so-called neuromodulatory transmitter systems including ACh, NE, and 5-HT. The modulatory transmitter systems are considerable and are described in [Section 3.9](#).

3.7.2.1 Neurochemistry of POC inputs to MOB Before departing to the MOB, it is germane to consider the neurochemistry of the input to the MOB from POC. As described later, feedback projections to the MOB from POC arise predominantly from glutamatergic, pyramidal neurons in layer II and III of PC, as well as other POC structures. These projections massively target GCs, where they form asymmetrical synapses on the cell bodies, basal dendrites, and spines of GCs (Price and Powell, 1970a). Activation of these feedback projections produces a negative field-potential recorded in the GCL (Walsh, 1959; Nakashima et al., 1978), as expected, if excitatory currents are flowing into GCs. Similar stimulation elicits IPSPs in mitral cells (Yamamoto et al., 1963; Nicoll, 1971; Mori and Takagi, 1978), due to GC excitation and subsequent GABA release (Halász and Shepherd, 1983). The transmitter of these feedback projections to GCs is glutamate. Activation of POC input to MOB excites GCs as measured by voltage-sensitive dye and field-potential recordings *in vitro*. This excitation is mediated by glutamate acting at both AMPA and NMDA receptors (Laaris and Ennis, 2002). A major function of these projections is to provide an inhibitory regulation of the firing rate and excitability state of mitral/tufted cells. Activation of these inputs modifies (inhibits) odor responses in MOB (Kerr and Hagbarth, 1955; von Baumgarten et al., 1962).

3.8 Amino Acid Receptor and Gap Junction Distribution in MOB

The following sections review the distributions of glutamate and GABA receptors and gap junction proteins in the MOB. In some cases, additional details about the neurophysiology and functional significance of the receptors are provided in the layer-specific sections earlier.

3.8.1 Ionotropic Glutamate Receptors

The three classes of ionotropic glutamate receptors- NMDA, AMPA, and kainate- are distributed extensively throughout the MOB. NMDA receptors are found in every layer of MOB except the IPL (Watanabe et al., 1993; Monyer et al., 1994; Petralia et al., 1994b). AMPA receptors are found in all layers except the ONL and subependymal layer (Molnar et al., 1993; Martin et al., 1993; Petralia and Wenthold, 1992; van den Pol, 1995). Kainate receptors are found in the MCL, EPL, and IPL (Monaghan and Cotman, 1982; Gall et al., 1990; Miller et al., 1990; Wisden and Seeburg, 1993).

3.8.1.1 Distribution in the GL Immunoreactivity for the NR1 NMDA receptor subunit is present at synapses between ON terminals and profiles that mostly appear to be mitral and tufted cell dendrites (Giustetto et al., 1997). JG neurons, including PG cells, also express NMDA NR1 subunit, which is also present at the dendrodendritic synapses with mitral/tufted cells in the glomeruli, as well as at synapses from centrifugal fibers and/or tufted cell collaterals in the periglomerular regions (Giustetto et al., 1997; Montague and Greer, 1999). Immunocytochemical studies show that AMPA receptors are also localized to processes and cell bodies of mitral, tufted, and PG cells (Petralia and Wenthold, 1992; Molnar et al., 1993). JG neurons, including PG cells, express GluR1, GluR2/3, and GluR4 AMPA receptor subunits (Giustetto et al., 1997; Montague and Greer, 1999); GluR4 staining is also associated with glia in the GL. Like NMDA receptors, AMPA receptors are present postsynaptically at synapses formed by ON terminals and at mitral/tufted-to-PG cell dendrodendritic synapses. Thus, NMDA and AMPA subunits are localized to ON-to-mitral/tufted cell synapses and to dendrodendritic synapses in the glomeruli (Giustetto et al., 1997).

3.8.1.2 Distribution in the Deeper Layers In the deeper layers, both mitral and tufted cells express NR1, GluR1, GluR2/3, and GluR4 AMPA receptor subunits, and kainate GluR5/6/7 receptor subunits, as indicated by immunohistochemical staining (Giustetto et al., 1997; Montague and Greer, 1999). Additionally, the interneurons of the EPL express GluR1, as do GCs (Giustetto et al., 1997; Montague and Greer, 1999; Hamilton and Coppola, 2003). GCs also express GluR2/3 subunits, but they are weakly immunoreactive for GluR1 and 4 receptor subunits (Montague and Greer, 1999). At the mitral/tufted-to-GC synapses, GluR2/3 and/or GluR1 AMPA receptor subunits are colocalized with NR1 NMDA receptor subunits. The GluR2 and 4 subunits occur in equal proportion as both flip and flop splice variants, whereas GluR1 and 3 subunits occur mostly as flip variants (Horning et al., 2004). In primary cell cultures, AMPA receptors of presumptive interneurons (most likely GCs), desensitize more rapidly and completely than AMPA receptors of presumptive mitral/tufted cells. This could be due to the occurrence of a higher proportion of flip subunits (Blakemore and Trombley, 2003). In dissociated preparations, AMPA receptors of presumptive GCs have also been reported to exhibit limited Ca^{2+} permeability, suggesting that GC AMPA receptors must include GluR2 subunits, which regulate Ca^{2+} permeability (Jardemark et al., 1997). The AMPA receptors of mitral cells and JG cells appear to exhibit Ca^{2+} permeability, however (Blakemore and Trombley, 2005; Ma and Lowe, 2005).

3.8.2 Metabotropic Glutamate Receptors

Neuroanatomical studies demonstrate that the MOB expresses high levels of mGluRs, suggesting that these receptors play important roles in olfactory processing. The eight mGluRs identified to date are subdivided into three groups based on sequence homology, signal transduction mechanisms, and pharmacology (Conn and Pin, 1997): Group I mGluRs (mGluR1, mGluR5), Group II mGluRs (mGluR2, mGluR3), and Group III mGluRs (mGluR4, mGluR6-8) (🔗 [Figure 6-8](#)).


3.8.2.1 Distribution in the GL JG cells express several types of mGluRs. Deciphering mGluR expression in the GL from the literature is difficult because mGluR expression has not been systematically examined in the different JG cell types. ET cells appear to have the same mGluR makeup as the deeper mitral/tufted cells; i.e., they express high levels of mGluR1 and lower levels of mGluR7 and mGluR8 (Ohishi et al., 1995; Kinzie et al., 1995; Saugstad et al., 1997). Other, as of yet unidentified JG cells have been reported to express low-to-moderate levels of Group I (mGluR5), Group II (mGluR2), and Group III (mGluR4/7/8) receptors (Ohishi et al., 1993a, 1998; Romano et al., 1995; Saugstad et al., 1997).

3.8.2.2 Distribution in Mitral/Tufted Cells Mitral/tufted cells express high levels of mGluR1 (Masu et al., 1991; Martin et al., 1992; Shigemoto et al., 1992; van den Pol, 1995; Sahara et al., 2001). Electron microscopy studies demonstrated that mGluR1 is present on the somata and apical and lateral dendrites of mitral/tufted cells (van den Pol, 1995). This expression pattern suggests that mGluR1 could mediate mitral/tufted cell responses to glutamatergic inputs from ON terminals, and/or could function as an auto- or heteroreceptor for glutamate released from apical or lateral dendrites of mitral/tufted cells. Mitral/tufted cells also express mRNA for mGluR7 and mGluR8 (Kinzie et al., 1995; Saugstad et al., 1997), although immunocytochemical studies indicate that these receptors are present on mitral/tufted cell axon terminals in the GCL and PC (Kinoshita et al., 1998; Wada et al., 1998). Mitral/tufted cells do not express Group II receptors (mGluR2/3), which contrast markedly with the strong expression of mGluR2 in mitral cells in the Accessory olfactory Bulb (AOB) (Ohishi et al., 1993b, 1998).

3.8.2.3 Distribution in the GCL GCs express the highest levels of mGluR5 (Group I) in the brain (Romano et al., 1995). EM studies have shown that mGluR5 is localized to portions of GC dendrites in the EPL apposed to presynaptic glutamatergic synapses from mitral/tufted cell lateral dendrites (van den Pol, 1995). This suggests that mGluR5 may mediate, at least in part, responses of GCs to glutamatergic inputs from mitral/tufted cells. GCs also express low-to-moderate levels of mGluR2 (Group II) (Ohishi et al., 1993a, 1998), and low levels of mGluR4 and mGluR7 (Group III) (Kinzie et al., 1995; Ohishi et al., 1995; Saugstad

■ Figure 6-8

The distribution of Group I, Group II, and Group III mGluRs in the MOB. See [Section 3.8.2](#) for details



et al., 1997; Wada et al., 1998). The precise cellular localization of these Group II/III mGluRs on GCs is not known. Recent experiments have shown that activation of mGluR5 directly and potently activates GCs. This action is mediated by an apparent inward current that involves, at least in part, a closure of K^+ channels (Heinbockel and Ennis, 2003).

3.8.3 GABA Receptors

GABA_A receptors are present in every layer of MOB except for the IPL (Palacios et al., 1981b; Bowery et al., 1987; Richards et al., 1987; Chu et al., 1990; Zhang et al., 1991; Persohn et al., 1992; Fritschy et al., 1992; Laurie et al., 1992). PG cells moderately express mRNAs for the $\beta 2$ and $\beta 3$ GABA_A receptor subunits (Laurie et al., 1992), and several ET cell subpopulations occur, which are differentially immunoreactive for the $\alpha 1$, $\alpha 3$, or both $\alpha 1$ and $\alpha 3$ receptor subunits (Panzanelli et al., 2005). Putative SA cells of the GCL express $\alpha 1$ and $\beta 2$ receptor subunits (Laurie et al., 1992). At the granule-to-mitral/tufted cell synapses of the EPL, $\alpha 1$, $\beta 2$, $\beta 3$, and $\gamma 2$ GABA_A receptor subunits occur (Fritschy et al., 1992; Giustetto et al., 1998; Sassoè-Pognetto et al., 2000). Mitral/tufted cells also express $\beta 1$ GABA_A receptor subunits (Laurie et al., 1992). In addition,

some mitral cells, middle tufted cells, and deep tufted cells express the $\alpha 3$ GABA_A receptor subunit, and many of the mitral cells coexpress CR (Panzanelli et al., 2005). The $\alpha 1$ GABA_A receptor subunits are perisynaptic, which suggests that GABA overflow from nearby synapses might regulate glutamate release from mitral/tufted cell dendrites (Panzanelli et al., 2004). GCs also express GABA_A receptors, including the $\alpha 2$, $\alpha 4$, $\alpha 5$, $\beta 3$, and δ receptor subunits (Laurie et al., 1992) and they exhibit inhibition (Mori and Kishi, 1982), which in a lower vertebrate (tiger salamander) has been shown to be GABAergic (Wellis and Kauer, 1994). In GABA_A $\beta 3$ receptor subunit-deficient mice, functional expression of GC GABA_A receptors was almost eliminated, amplitudes of mitral/tufted cell miniature IPSPs, theta-frequency oscillations, and γ -frequency oscillations were increased, and discrimination of closely related mixtures of alcohols after training was poor relative to normal mice (Nusser et al., 2001). These results provide functional evidence for the importance of both GC and mitral/tufted cell GABA_A receptors in olfaction.

GABA_B receptors are restricted to the GL in MOB (Bowerly et al., 1987; Chu et al., 1990). However, there are reports that GABA_B autoreceptors modulate GABA release at granule-to-mitral cell reciprocal synapses (Isaacson and Vitten, 2003). Activation of GABA_B receptors has been observed to reduce dendrodendritic inhibition via suppression of GABA release from GCs due to reduction of high voltage-activated Ca²⁺ currents (Isaacson and Vitten, 2003). In a lower vertebrate (frog), GABA_B and dopamine D₂ receptors both appear to strengthen mitral/tufted cell odor signaling by reducing spontaneous activity (Duchamp-Viret et al., 2000).

3.8.4 Glycine and Glycine Receptors

Cells in the EPL and GCs are also immunoreactive for glycine (Pourcho et al., 1992), and the MOB contains significant levels of both this amino acid and taurine, which inhibit cultured olfactory bulb cells (Trombley and Shepherd, 1994; Trombley et al., 1999) and mitral/tufted cells in situ (Belluzzi et al., 2004). The MOB exhibits strong labeling for glycine receptors (van den Pol and Gorcs, 1988), and the β receptor subunit has been localized to mitral/tufted cell synapses (Malioso et al., 1991).

3.8.5 Gap Junctions

Gap junctions are thought to play a role in synchronization of MOB activity (Schoppa and Westbrook, 2002; Christie et al., 2005; Hayar et al., 2005). Gap junction inhibitors block γ -frequency (20–70) Hz oscillations generated by chemically and electrically coupled networks of mitral and GCs (Friedman and Strowbridge, 2003), suggesting a role for both types of synapses. [Table 6-3](#) summarizes the distribution of gap junctions and the gap junction proteins, connexins, in the MOB. Gap junctions have been observed on mitral, tufted, and GC bodies (Reyher et al., 1991; Paternostro et al., 1995; Miragall et al., 1996; Kosaka and Kosaka, 2003). In the GL, mitral/tufted cell dendrites have also been observed to form gap junctions with each other, with PG cells, and with processes of unidentified cells (Kosaka and Kosaka, 2003, 2004, 2005; Kosaka et al., 2005; Rash et al., 2005). The junctions tend to occur as mixed synapses with glutamatergic synapses. Some of the unidentified processes involved in these synapses are not immunoreactive for either DA or GABA and could be mitral/tufted cell dendrites (Kosaka and Kosaka, 2003). In support of this, mitral/tufted cell dendrites of a lower vertebrate (tiger salamander) have been observed to form asymmetrical synapses with each other in the GL (Allen and Hamilton, 2000). In addition to their GL synapses, mammalian mitral/tufted cell dendrites form some gap junctions with dendrites of GCs (Landis et al., 1974; Kosaka et al., 2005) and other interneurons (Kosaka and Kosaka, 2003) in the EPL.

The predominant gap junction protein of MOB neurons appears to be connexin36 (Condorelli et al., 1998, 2000; Belluardo et al., 2000; Rash et al., 2000, 2005), although mitral/tufted cells and unidentified interneurons in the EPL and GCL have been reported to express other connexin mRNAs (connexin43, Miragall et al., 1996; connexin45, Zhang and Restrepo, 2002) or proteins (connexin43, Theis et al., 2003) that are mainly expressed by glial cells. It is noteworthy that expression of connexin36 mRNA is particularly high within (but not uniform among) the glomeruli (Teubner et al., 2000; Zhang and Restrepo, 2003), and

that expression of this connexin remains high into adulthood (Hormuzdi et al., 2001). This is in contrast to the hippocampus, where connexin36 mRNA expression peaks during early postnatal life and subsequently declines (Rozentel et al., 2000). In addition, γ -frequency electrical signaling is disrupted in connexin36 knockout mice (Hormuzdi et al., 2001), providing evidence that this connexin is involved in oscillatory signaling. Gap junction channels formed by connexin36 exhibit unitary conductances of ~ 15 pS, weak transjunctional voltage dependence, and are also permeable to cAMP and IP₃ (Rozentel et al., 2000; Teubner et al., 2000). Thus, gap junctions connecting MOB neurons could participate in both fast oscillatory signaling and slower, second messenger-mediated signaling, even in adults.

3.9 Neuromodulatory Inputs to MOB

MOB and most components of POC receive extrinsic inputs from cholinergic, noradrenergic, serotonergic, and dopaminergic cell groups in the brainstem and basal forebrain.

3.9.1 Cholinergic Inputs to MOB


3.9.1.1 Cholinergic Innervation Pattern and Receptor Distribution In the mouse, about 3.5% of all neurons that project to the bulb originate in the nucleus of the horizontal limb of the diagonal band (NDB); far fewer originate in the vertical limb of DB (Carson, 1984; Shipley and Adamek, 1984). At least two distinct transmitter-specific populations of NDB neurons project to the MOB (Zaborszky et al., 1986). About 20% of the NDB neurons that project to the bulb are cholinergic; most of these cells are concentrated in the rostromedial portion of the horizontal limb of NDB. Many NDB-MOB projection neurons are GABAergic and occur mainly in the caudo-lateral aspect of NDB (Zaborszky et al., 1986). AChE is concentrated in the IPL, GCL, inner third of the EPL, and the GL. Some glomeruli are more intensely stained for AChE and correspond to regions of leutenizing hormone releasing hormone (LHRH) innervation (Zheng et al., 1988), and may include the modified glomerular complex as defined by Greer and colleagues (Teicher et al., 1980; Greer et al., 1982). There are also AChE-positive neurons in the bulb (Nickell and Shipley, 1988b). Choline acetyltransferase (ChAT), the biosynthetic enzyme for ACh, is located in axons distributed in the same layers as described for AChE. As shown in [Figure 6-9](#), cholinergic fibers are distributed throughout the MOB, but are especially heavy in the GL and IPL. The GABAergic projection from NDB (Zaborszky et al., 1986) is more difficult to characterize than the cholinergic projection, because the intrinsic GABAergic PG cells and GCs provide intrinsic GABAergic innervation.

Cholinergic receptor localization in the MOB is in close agreement with terminal staining from the NDB ([Table 6-4](#)). Muscarinic receptors are found in all layers while nicotinic receptors are found in all layers except the IPL (see [Table 6-4](#)). The EPL exhibits intense staining for muscarinic-1 (M_1), M_3 , and M_4 receptors (Rotter et al., 1979; Spencer et al., 1986; Buckley et al., 1988; Fonseca et al., 1991b). M_2 receptors are immunocytochemically localized to PG cells in the GL and tufted cells in the EPL (Fonseca et al., 1991a). M_2 receptors are also present in the IPL and GCL. Nicotinic receptors show a different regional distribution throughout the MOB and are concentrated in the GL and EPL (Hunt and Schmidt, 1978; Sahin et al., 1992).

3.9.1.2 Cholinergic Actions in MOB Only limited information is available about cholinergic actions in MOB. Electrical activation of NDB has been reported to depress (Nickell and Shipley, 1988a) or increase (Kunze et al., 1991, 1992) mitral cell activity indirectly via primary effects on GABAergic GCs. NDB stimulation also reduced the field potential in the MOB caused by stimulation of the anterior commissure (Nickell and Shipley, 1993), an effect mediated by presynaptic inhibition of anterior commissure terminals via muscarinic receptors. One interpretation of these results is that cholinergic input to MOB may function to modulate interhemispheric transmission of olfactory information. In this regard, it is noteworthy that anterior commissural fibers are required for access and recall of olfactory memories between the two hemispheres. Infusion of ACh into MOB was reported to reduce paired-pulse depression of lateral olfactory tract (LOT)-evoked field potentials recorded in the GCL. This effect was attributed to

■ Figure 6-9

Neuromodulatory transmitter inputs to MOB. Darkfield photomicrographs showing the distribution of cholinergic (a), noradrenergic (b), and serotonergic (c) fibers revealed respectively with immunohistochemistry for choline acetyltransferase (ChAT), dopamine- β -hydroxylase (DBH), and serotonin (5-HT). Reprinted from *Handbook of Chem. Neuroanat. Integrated Sys. CNS, Vol. 12, Part III, Chapter III, The Olfactory System*, M. Shipley et al., pp. 469–573, 1996, with permission from Elsevier, Ltd



muscarinic receptor-mediated inhibition of GABA release from GCs (Elaagouby et al., 1991). In slice preparations, nicotinic but not muscarinic receptor agonists directly excited mitral cells, and this effect appeared to be due to an inward current with a reversal potential of -5 to $+10$ mV (Castillo et al., 1999). In slices, muscarinic receptor agonists inhibited GCs (Castillo et al., 1999), and paradoxically, also appeared to increase GABA release from these cells. The same study reported that in the GL only bipolar PG cells were sensitive to nicotine (Castillo et al., 1999). The morphological identity of these cells is unclear.

3.9.2 Dopaminergic (DA) Input to MOB

There is no known extrinsic DA innervation of the MOB. As noted earlier, however, the MOB contains several hundred thousand intrinsic JG DA neurons that mediate potent presynaptic inhibition of ON terminals via activation of D2 receptors. This presynaptic inhibition as well as DA receptor localization is discussed in [Section 3.2.3.2](#).


3.9.3 Noradrenergic (NE) Input to MOB

3.9.3.1 NE Innervation Pattern and Receptor Distribution A significant modulatory input to the MOB is from the pontine nucleus, locus coeruleus (LC). In the rat, all LC neurons contain the neurotransmitter, norepinephrine (NE); LC contains the largest population of NE neurons in the brain. It has been estimated that up to 40% of LC neurons (400–600 of a total of 1,600 LC neurons) project to the bulb in the rat (Shipley et al., 1985). A subset of LC neurons projecting to MOB contain NPY (Bouna et al., 1994). NE fibers terminate in MOB with laminar specificity. LC axons project mainly to the subglomerular layers of the bulb, particularly the IPL and GCL (McLean et al., 1989). The EPL and MCL are moderately innervated, whereas the GL is nearly devoid of NE input.

NE receptors occur in multiple layers of the MOB and they are expressed by multiple cell types, in general consistent with the pattern of NE fiber innervation ([Figure 6-10](#) and [Table 6-4](#)). Receptor

■ Figure 6-10

The distribution of NE receptors in the MOB. See Section 3.9.3 for details. CFF, centrifugal fibers



localization studies also suggest that individual MOB cells express multiple NE receptor subtypes. α receptors are abundantly expressed in MOB. Binding sites for $\alpha 1$ receptors are particularly dense in the EPL, and moderate in the MCL and IPL/GCL (Young and Kuhar 1980a; Sargent-Jones et al., 1985); the density of $\alpha 1$ receptors in the EPL was the highest in the brain (Young and Kuhar, 1980a). Consistent with these findings are observations that $\alpha 1$ receptor mRNA is expressed by mitral cells ($\alpha 1_A$, $\alpha 1_B$, $\alpha 1_D$ subtypes) and GCs ($\alpha 1_A$, $\alpha 1_D$ subtypes) (McCune et al., 1993; Pieribone et al., 1994; Domyancic and Morilak, 1997; Day et al., 1997). The MCL, IPL, and GCL express moderate-to-dense levels of $\alpha 2$ receptors (Young and Kuhar, 1979; Nicholas et al., 1993; Rosin et al., 1996; Talley et al., 1996; Winzer-Serhan et al., 1997a, b), and $\alpha 2_C$ receptor mRNA is expressed by both mitral cells and GCs (Nicholas et al., 1993; Rosin et al., 1996; Winzer-Serhan et al., 1997a, b). The high density of $\alpha 2$ receptor binding sites and NE fibers in the IPL/GCL suggests the possibility that these receptors may be present presynaptically on axon terminals of NE fibers themselves or on other centrifugal inputs to the soma and/or proximal dendrites of GCs (Figure 6-10). Unidentified JG neurons have been reported to express $\alpha 1$ and $\alpha 2$ receptors (Rosin et al., 1996; Winzer-Serhan et al., 1997a, b). By contrast, the level of β receptors is relatively low in the MOB, and they appear to

be localized preferentially to the GL, IPL, and GCL (Palacios and Kuhar, 1982; Booze et al., 1989; Wanaka et al., 1989; Woo and Leon, 1995). JG, mitral, and tufted cells, as well as subsets of GCs express the $\beta 1$ receptor subtype (Yuan et al., 2003).

3.9.3.2 Physiological Actions of NE Although, NE clearly plays significant roles in olfactory function, the effects of NE at the cellular and network levels are somewhat discrepant. For example, LC stimulation was reported to have no effect on LOT-evoked field-potentials recorded in the GCL (Perez et al., 1987). A subsequent study reported that LC stimulation initially decreased and then subsequently increased paired-pulse depression of GC field-potential responses to LOT stimulation (Okutani et al., 1998). These effects were attributed to activation of β receptors. Another field-potential study reported that NE infusion into MOB, acting at $\alpha 1$ receptors, increased the depolarization of GC dendrites elicited by LOT stimulation. Mitral cell responses to antidromic shocks were not affected, suggesting that NE excites GC (Mouly et al., 1995). In neonatal animals, β receptor stimulation in MOB decreased LOT-evoked, paired-pulse inhibition of GC field potentials (Wilson and Leon, 1988). It is unclear if this was mediated by presynaptic inhibition of transmitter release from MCs and/or increased excitability of GCs.

Cellular recording studies are also somewhat discrepant. In the rabbit and cat, Salmoiraghi et al. (1964) and McLennan (1971) found that iontophoretically applied NE inhibited mitral cells. This effect was blocked by the GABA_A receptor antagonist, bicuculline. In the isolated turtle bulb (Jahr and Nicoll, 1982), mitral cell spike activity increased, and GABAergic IPSPs decreased, following bath application of NE. In dissociated MOB cultures, NE decreased mitral cell-to-GC dendrodendritic synaptic transmission acting presynaptically at $\alpha 2$ receptors to decrease Ca²⁺ currents in both granule and mitral cells (Trombley, 1992; Trombley and Shepherd, 1992). NE release, evoked by selective chemical activation of LC in vivo, enhanced the response of mitral cells in response to weak (i.e., perithreshold) but not strong (i.e., suprathreshold) stimulation of the olfactory epithelium (Jiang et al., 1996). Interestingly, NE release from LC axon terminals is facilitated and suppressed by activation of presynaptic nicotinic and muscarinic cholinergic receptors, respectively (El-Etri et al., 1999). In rat MOB slices, application of NE or $\alpha 1$ receptor agonists, but not $\alpha 2$ or β receptor agonists, also selectively increased mitral cell responses to perithreshold intensity ON stimulation (Ciombor et al., 1999). Noradrenergic agonists had no effect on ON-evoked field-potentials recorded in the GL, or on ON-evoked postsynaptic currents in mitral cells (Hayar et al., 2001). This suggests that NE-evoked modulation of ON-evoked mitral cell spiking is mediated by postsynaptic actions on bulb neurons. In voltage clamp recordings, NE or $\alpha 1$ agonists directly evoked an inward current in mitral cells that appeared to be due to closure of K⁺ channels. In current clamp recordings from bistable mitral cells, $\alpha 1$ agonists shifted the membrane potential from the downstate (−52 mV) toward the upstate (−40 mV), and significantly increased spike generation in response to perithreshold ON input. Taken together, these findings suggest that NE release directly alters mitral cell excitability in a manner that could increase their sensitivity to weak ON input, perhaps to improve the detection of weak odorants.

NE inputs to the bulb are critical to olfactory function. Olfactory cues increase the discharge of LC neurons in behaving animals (Aston-Jones and Bloom, 1981) and increase NE levels in the MOB (Chanse and Kopin, 1968; Rosser and Keverne, 1985; Brennan et al., 1990). LC projections to the main and accessory olfactory bulb are pivotal to the formation and/or recall of specific olfactory memories, pheromonal regulation of pregnancy and postpartum maternal behavior (Pissonnier et al., 1985; Kaba and Keverne, 1988; Brennan et al., 1990; Kendrick et al., 1992). NE plays an important role in the so-called Bruce effect in mice: when impregnated female mice are exposed to the odor of a strange male, they abort; if exposed to the odor of the impregnating male they do not abort (Kaba et al., 1989). Systemic administration of adrenergic receptor antagonists or 6-OHDA lesions, selectively destroying only the NE inputs to the MOB, cause the female to abort when presented with the odor of the impregnating male (Rosser and Keverne, 1985; Kaba and Keverne, 1988; Brennan et al., 1990). Finally, NE has been shown to play a critical role in olfactory learning in young animals. In neonatal rats, NE release via tactile stimulation leads to a preference for an odor associatively paired with this stimulation (Sullivan et al., 1989). The conditioned preference is associated with odor-specific metabolic changes in the bulb (Coopersmith and Leon, 1984). Following the conditioning, there is an increased inhibition of mitral cells by the odor (Sullivan et al., 1989). Such

conditioning is abolished by eliminating NE input to the bulb or via β receptor antagonists (Sullivan et al., 1989, 1992, 2000; Wilson and Sullivan, 1991; Moriceau and Sullivan, 2004). Recent studies from McLean's laboratory demonstrate that this β receptor-dependent neonatal learning involves activation of intracellular cAMP and CREB pathways (see McLean and Harley, 2004 for review).

3.9.4 Serotonergic (5-HT) Inputs to MOB

3.9.4.1 5-HT Innervation Pattern, Receptor Distribution, and Physiological Actions The midbrain dorsal and median raphe provides strong inputs to the MOB. In the rat, about 1,000 dorsal and 300 median raphe neurons project to the bulb. These neurons are serotonergic and they do not contain TH (McLean and Shipley, 1987a, b) or substance P (Zaborszky et al., 1986). As shown in [Figure 6-9](#), 5-HT fibers are present in all layers of MOB, but with varying densities. Input to the GL is especially dense, while the EPL contains very low density. [^3H]5-HT-labeled boutons have been observed in close proximity to probable SA cells in the EPL (Halász et al., 1978). The MCL, IPL, and GCL have a fairly heavy and uniform innervation, but not as dense as that of the GL. Thick serotonergic fibers preferentially innervate the glomeruli of MOB, whereas thinner serotonergic axons preferentially innervate inframitral layers (McLean and Shipley, 1987a, b). In neocortex, thick axons arise from the median raphe and thin axons arise from dorsal raphe (McLean and Shipley, 1987a, b) and the same segregation occurs in MOB.

In agreement with the 5-HT fiber distribution, 5-HT receptors are localized in most layers of the MOB ([Table 6-4](#)). The 5-HT_{1A} subtype is present on processes in the EPL, MCL, and minimally in the GCL; mitral cells and GCs appear to express the 5-HT_{1A} receptor (Pompeiano et al., 1992). 5-HT_{1C} is present in the GCL. It is possible that many of the cells possessing 5-HT receptors are glia (Whitaker-Azmitia et al., 1993). The 5-HT₂ receptors are present in the GL, EPL, and MCL. The 5-HT_{2A} receptor subtype is found in the EPL and in mitral and tufted cells (McLean et al., 1993, 1995; McLean, 1994; Pompeiano et al., 1994; Hamada et al., 1998; Cornea-Hebert et al., 1999). In mitral and tufted cells, 5-HT_{2A} receptors colocalize with the β 1 receptor (Yuan et al., 2003). The 5-HT_{2A} receptors are also present in unidentified JG cells in the GL (Hamada et al., 1998; Cornea-Hebert et al., 1999). The 5-HT_{2C} receptor subtype is expressed at fairly high levels by GCs and at relatively low levels by unidentified JG neurons (Clemett et al., 2000). JG cells have also been reported to express 5-HT₃ receptors (Tecott et al., 1993).

At present, there is only limited data about the actions of serotonin in the MOB. In the GL, it was recently reported that serotonin depolarized 34% of JG cells *in vitro* via activation of 5HT_{2C} receptors (Hardy et al., 2005). The serotonin-induced depolarization was due to activation of a nonselective cation current with a reversal potential of -44mV . The heterogeneous electrophysiological properties of 5HT-responsive JG cells suggested that several types of JG cells could be targeted by 5HT centrifugal fibers. A subset of mitral cells was also depolarized by serotonin acting at 5HT_{2A} receptors. In contrast with these results, another subset of mitral cells was hyperpolarized by serotonin, an action that was indirectly mediated by GCs as it was blocked by GABA_A receptor antagonists (Hardy et al., 2005). This effect of serotonin was also thought to be mediated by 5HT_{2A} receptors. Behavioral studies indicated that lesion of serotonergic fibers reversed conditioned olfactory learning (Morizumi et al., 1994), and also induced glomerular atrophy. Behavioral work on neonates by McLean et al. (1996) showed that serotonin depletion or 5-HT₂ receptor antagonism compromised olfactory learning and that serotonin release or 5-HT₂ receptor activation promoted odor conditioning. 5-HT release appeared to facilitate β receptor-mediated, NE-induced olfactory learning by facilitating cAMP mechanisms (see McLean and Harley, 2004 for review).

4 Olfactory Input Regulates Neurochemistry of Specific MOB Neurons

ORNs turn over throughout life, most MOB interneurons are added after birth, and new interneurons are continually added during adulthood. The MOB is consequently a highly plastic neural system that is readily affected by both sensory experience and deprivation. A variety of studies showed that occlusion of one naris

during early postnatal life markedly affected development of the ipsilateral MOB, resulting in numerous cellular changes and ~25% smaller MOB (Brunjes, 1994). The studies also showed that naris occlusion resulted in reduced DA levels (Baker et al., 1983; Brunjes et al., 1985; Wilson and Wood, 1992). During development, the maturing dopaminergic JG neurons only exhibited TH immunoreactivity after they reached the GL, and they failed to do so after naris occlusion (McLean and Shipley, 1988; Baker, 1990; Baker and Farbman, 1993; Baker et al., 1993). In adults, both naris occlusion and destruction of the mature olfactory sensory neurons by nasal irrigation with ZnSO₄ or detergent, or by severing the ON, reduced transneuronal regulation of DA levels and TH expression in rats (Nadi et al., 1981; Kawano and Margolis, 1982; Baker et al., 1983, 1993; Erlich et al., 1990; Cho et al., 1996; Couper Leo et al., 2000), hamsters (Kream et al., 1984) and dogs (Nadi et al., 1981). In concert with the reduced TH expression, dopamine D2 receptor density was increased in the ONL and GL (Guthrie et al., 1991). TH expression was also downregulated in mice homozygous for a null mutation in the olfactory cyclic nucleotide gated channel subunit-1 (OCNC1), which rendered them functionally anosmic (Baker et al., 1999). The loss of TH expression following naris occlusion and deafferentation was not due to cell death, because JG neurons could be detected with antibodies to other DA enzymes (Baker et al., 1984). Moreover, both GAD and GABA, which are coexpressed by many dopaminergic JG neurons, were still detected (Kosaka et al., 1987a, b, c; Baker et al., 1988, 1993; Stone et al., 1990). Depolarizing stimuli could induce TH expression *in vitro* and *in vivo* (McMillian et al., 1994; Philpot et al., 1998), and the *in vitro* studies showed that the induction required Ca²⁺ influx into the bulb neurons (Cigola et al., 1998). In an MOB neuron/olfactory epithelium coculture, TH expression was abolished by NMDA receptor blockade (Puche and Shipley, 1999), suggesting that glutamate released by ORNs stimulates the dopaminergic JG neurons, resulting in Ca²⁺ influx and regulation of TH expression. Thus, both the developmental induction and maintenance of the DA phenotype appear to be transneuronally regulated and to depend on the presence and normal functioning of the ON. Although the intracellular mechanisms remain to be elucidated, the distributions of *c-fos* immediate early gene mRNA and Fos protein immunoreactivity were found partially to overlap with TH distribution and were downregulated following naris occlusion (Guthrie and Gall, 1995a, b; Jin et al., 1996). Fos-B was implicated in the regulation of TH expression through interactions with the AP-1 motif (Liu et al., 1999).

The expression of other neurotransmitters and receptors is also influenced by the afferent sensory input to the MOB. In the hamster, many JG neurons express substance P, and this peptide was found to be downregulated following chemical deafferentation (Kream et al., 1984). By contrast in rats, glomerular mGluR1 immunoreactivity was increased. Initially, the mGluR1a mRNA level increased, but it then declined (like TH expression), presumably due to product inhibition (Ferraris et al., 1997; Casabona et al., 1998). As with chemical deafferentation, neonatal naris occlusion of rats and mice had both negative and positive effects. It reduced the density of β 1- and β 2-adrenergic receptors as determined by quantitative autoradiography (Woo and Leon, 1995), although NE levels were not reduced following either naris occlusion (Brunjes et al., 1985) or chemical deafferentation (Nadi et al., 1981). Similarly, although levels of GluR1 were not reduced following chemical deafferentation (Ferraris et al., 1997), immunoreactivity of cell bodies in the EPL for both GluR1 (Hamilton and Coppola, 2003) and for PV were greatly reduced, as was immunoreactivity of nondopaminergic JG neurons for calbindin (Philpot et al., 1997). Focal surgical deafferentation had similar effects on PV and calbindin immunoreactivity (Couper Leo et al., 2000). CR immunoreactivity of other interneurons was not affected, however. In addition, although the number of synapses between mitral cell bodies and GC dendrites were reduced following naris occlusion, mitral/tufted cell inhibition appeared to be increased, possibly via enhanced NMDA receptor-mediated excitation of the GCs by the mitral/tufted cells at their reciprocal synapses (Wilson, 1995). Expression of voltage-sensitive Na⁺ channel α II and β 1 subunits was also downregulated in tufted cells, but not in mitral cells (Sashihara et al., 1997). The afferent sensory input to MOB therefore appears to influence the expression of a variety of neurotransmitters, receptors, and ion channel proteins, with different effects observed on different cell types.

Neurotrophic factors, receptors, and kinases were also shown to be affected by naris occlusion. These include insulin receptor (IR) kinase, brain-derived neurotrophic factor (BDNF), neurotrophin receptors TrkB and TrkC, and mitogen-activated protein kinase/extracellular signal-related kinase (MAPK/ERK)

(Mackay-Sim and Chuah, 2000). Expression of IR kinase was reduced, as was the insulin-induced suppression of current through the Kv1.3 channel, which is typically tyrosine phosphorylated in the N and C termini by IR kinase (Fadool et al., 2000). The Kv1.3 channel carries most of the outward current of cultured MOB neurons (Fadool and Levitan, 1998). Moreover, BDNF levels initially increased *in vivo* and subsequently decreased, without concomitant effects on TrkB levels (McLean et al., 2001; Tucker and Fadool, 2002). In cultured MOB neurons isolated from the occluded side, however, the BDNF-induced phosphorylation of Kv1.3 was increased, indicating that tyrosine kinase activity was increased. Activation of ERK pathway was also downregulated (Mirich et al., 2004). Levels of other kinases did not appear to be altered, however, at least as examined by immunocytochemical staining (Liu, 2000). These results suggest that altered levels of neurotrophic factors and altered kinase expression or activity may contribute to the developmental changes observed in MOB neurons following naris occlusion.

5 Primary Olfactory Cortex: Piriform Cortex

The MOB projects to a collection of structures referred to as POC (de Olmos et al., 1978). These structures may be divided into three groups: (1) the AON, (2) medial olfactory cortex (indusium griseum, anterior hippocampal continuation, taenia tecta, and the olfactory tubercle), and (3) lateral olfactory cortex comprising, from rostral to caudal, PC and the periamygdaloid, transitional and entorhinal cortices (▶ [Figure 6-7](#)). At least from an olfactory perspective PC is perhaps the most extensively studied and well understood member of POC. A thorough presentation of the cytoarchitecture, connections, and neurochemistry of each of the components of POC is beyond the scope of the present report, and readers are directed to more complete reviews of these subjects (Shipley et al., 1996, 2004). Here, we focus on PC (▶ [Figures 6-3](#) and ▶ [6-7](#)) as an example of POC.

5.1 Organization and Architecture

PC is the largest component of the lateral olfactory cortical areas. It extends rostrocaudally and is located in the cortical mantle ventral to the rhinal sulcus (▶ [Figure 6-3](#)). Haberly and Price (1978a) divided PC into three layers that were further subdivided on the basis of cytoarchitecture and afferent connections. Layer I, the superficial plexiform layer, is divided into Ia and Ib, which receive different afferents: layer Ia, from the ipsilateral MOB, and layer Ib, association fibers from AON and from other parts of POC. Layer II, the superficial compact cell layer, is divided into two zones; the more superficial zone has a lower cell density and the deeper zone has a higher cell density. Layer III is the thickest cell layer. The endopiriform nucleus lies deep to the PC (Loo, 1931; Haberly and Price, 1978a). Behan and Haberly (1999) reported that this nucleus, while similar in many regards to PC proper, has unique intrinsic and extrinsic connections that differ from PC. Additionally, electrophysiological properties of endopiriform neurons, such as certain K⁺ currents (e.g., A-current), differ from those of PC neurons (Banks et al., 1996). As reviewed later, there is considerable rostrocaudal heterogeneity in the structure and connections of anterior versus posterior PC (Haberly and Price, 1978a; Luskin and Price, 1983). There is evidence for distinct subdivisions within the anterior PC based on cytoarchitecture, connections, and neurochemistry (Ekstrand et al., 2001b). Candidate transmitters and neurotransmitter receptors in PC are summarized in ▶ [Tables 6-5](#) and ▶ [6-6](#).

5.2 PC Cell Types

Neurons in PC can be divided into two main classes: (1) principal or output neurons—the pyramidal cells and (2) intrinsic interneurons.

■ **Table 6-5**

Neuroactive substances in PC neurons

Neurotransmitter	Cell Location	Cell Type	References
Calbindin/ Calretinin	Mainly layer II	Pyramidal, basket, bi-multipolar	Celio (1990), Morales and Bloom (1997)
CCK	Mainly layer II and III	Pyramidal, multipolar, basket	Roberts et al. (1982), Westebroek et al. (1987)
Dyn B	Layer II and III	Nonpyramidal fusiform, multipolar	Fallon and Leslie (1986)
ENK	Mainly layer II	Fusiform, pyramidal, multipolar	Harlan et al. (1987)
GABA/GAD	All layers	Diverse interneurons	Ekstrand et al. (2001a), Haberly et al. (1987), Morales and Bloom (1997)
Glutamate	Layers II and III	Pyramidal cells	See Section 3.8.1
Neurotensin	NR	NR	Hara et al. (1982)
Parvalbumin	Mainly layers II and III	Multipolar, basket	Celio (1990)
VIP	Mainly layers II and III	Bipolar, basket	Ekstrand et al. (2001a), Roberts et al. (1982), Sanides-Kohlrausch and Wahle (1990b)

NR, not reported. Adapted from Handbook of Chem. Neuroanat. Integrated Sys. CNS, Vol. 12, Part III, Chapter III, The Olfactory System, M. Shipley et al., pp. 469–573, 1996, Elsevier, Ltd

5.2.1 Pyramidal Cells

PC has two principal layers of pyramidal cells, layers II and III, corresponding to superficial and deep pyramidal cells. Both superficial and deep pyramidal neurons share characteristic features similar to pyramidal cells in the hippocampus and other cortical areas, including: (1) a primary apical-dendritic trunk that extends radially toward the pial surface and arborizes in layer Ia and Ib (Haberly, 1983), (2) secondary or basal dendrites that extend from the soma into deeper parts of PC. Both the apical and basal dendrites are heavily invested with spines and varicosities, and (3) a myelinated axon that typically extends deep to the soma that terminates on other local pyramidal cells and interneurons (see later), or projects back to MOB. Reconstructions of intracellularly filled pyramidal cells show that they have extensive axonal projections covering almost an entire cerebral hemisphere, including local connections to anterior and posterior PC as well as arborizations in orbital cortex, insular cortex, olfactory tubercle, perirhinal cortex, entorhinal cortex, and amygdaloid cortex (Johnson et al., 2000). The collaterals of deep pyramidal axons travel into the endopiriform nucleus (Tseng and Haberly, 1989), whereas those of superficial pyramidal neurons do not appear to enter this structure. The apical dendritic tree of superficial pyramidal cells generally branches more extensively and is more densely invested with spines and varicosities than that of deep pyramidal cells. Pyramidal cells contain glutamate and participate in intrinsic (intra-PC) and extrinsic excitatory projections as discussed later.

5.2.2 Interneurons

PC possesses a variety of interneurons as evidenced by morphological and neurochemical heterogeneity (Haberly and Presto, 1986; Haberly et al., 1987; Ekstrand et al., 2001a). A thorough consideration of PC

Table 6-6

Neurotransmitter receptors in the PC

Receptor	Layer	References
Cholinergic		Buckley et al. (1988), Hill et al. (1993), Levey et al. (1991), Rotter et al. (1979), Sahin et al. (1992), Seguela et al. (1993), Spencer et al. (1986)
mAChR1	I, II	
mAChR2	II	
mAChR3	II	
mAChR4	II	
nAChR	II, III	
Noradrenergic		Day et al. (1997), Domyancic and Morilak (1997), Nicholas et al. (1993), Palacios and Kuhar (1980), Pieribone et al. (1994), Rosin et al. (1996), Sargent-Jones et al. (1985), Talley et al. (1996), Unnerstall et al. (1984), Wanaka et al. (1989), Winzer-Serhan et al. (1997a, b), Young and Kuhar (1979, 1980a)
α 1	II, III	
α 2	I, II, III	
β 1	II	
β 2	II	
Dopaminergic		Freneau et al. (1991), Huang et al. (1992), Mansour et al. (1990a, b)
D ₁	II	
D ₂	NR	
Serotonergic		Clemett et al. (2000), Cornea-Hebert et al. (1999), Hamada et al. (1998), Hoffman and Mezey (1989), Mengod et al. (1990a, b), Morales and Bloom (1997), Pompeiano et al. (1992, 1994), Tecott et al. (1993), Wright et al. (1995)
5-HT _{1A}	II	
5-HT _{1C}	II	
5-HT _{2A/C}	I, II, III	
5-HT ₃	NR	
Glutamatergic		Gall et al. (1990), Kinoshita et al. (1998), Kinzie et al. (1995), Monaghan et al. (1985), Ohishi et al. (1993a, b, 1995, 1998), Petralia and Wenthold (1992), Petralia et al. (1994a, b), Romano et al. (1995), Saugstad et al. (1997), Shigemoto et al. (1992), Sun et al. (2000), Wada et al. (1998), Wisden and Seeburg (1993)
KA	II	
NMDA	Ia, II	
AMPA	II	
mGluR		
Group I mGluR (mGluR1/5)	II, III	
Group II mGluR (mGluR2,3)	NR	
Group III mGluR (mGluR4,7,8)	Ia	
GABAergic		Bowery et al. (1987), Margeta-Mitrovic et al. (1999), Palacios et al. (1981b), Young and Kuhar (1980b)
GABA _A	I, III	
GABA _B		

NR, not reported

Adapted from Handbook of Chem. Neuroanat. Integrated Sys. CNS, Vol. 12, Part III, Chapter III, The Olfactory System, M. Shipley et al., pp. 469–573, 1996, with permission from Elsevier, Ltd

interneurons is beyond the scope of this review and detailed descriptions of these cells are available elsewhere (Shipley et al., 1996). Here, we present the distribution of the main interneuron types and some of their major neurochemical features. Interneurons are distributed throughout all layers and regions of this structure, and most, if not all, contain GABA (Haberly et al., 1987). Major interneuronal subtypes include the following:

1. Horizontal cells with large fusiform somata, found only in the superficial part of layer I (Haberly, 1983). These GABAergic cells produce dendrites that ramify within layer Ia and to a lesser extent within layer Ib (Haberly et al., 1987). Layer I interneurons are thought to function in feedforward inhibitory systems.
2. Spiny cells, found in layers I and III. Spiny actually refers to several distinct morphological subtypes (e.g., those with spherical or fusiform somata) that exhibit dendritic spines (Haberly, 1983). The dendrites of layer I cells extend into all parts of layer I and occasionally into deeper layers. Spiny cells give rise to unmyelinated axons that are horizontally oriented. Giant cells, another subcategory of spiny cell, have the largest soma size in PC. These cells are concentrated in the ventral parts of layer III.
3. Smooth cells, also found in layers I and III. Those in layer I exhibit a variety of cell shapes and sizes and they are distinguished by a lack of spines on the cell body or dendrites and by the presence of beaded varicosities on the distal dendrites. Smooth cells are the most numerous nonpyramidal cells in layer III, and they are preferentially distributed in the middle and deep parts of layer III. The dendrites emerge from numerous sites on the soma, usually branch once or twice near the soma, and then extend radially in all directions, with few subsequent bifurcations. The dendritic tree typically respects the boundaries of layer III.
4. Neurogliaform cells, found in layers I and II, have spherical somata and they are the most common cell type in layer I. The dendritic trees of these cells are usually restricted to a single sublamina in layer I. The axons are unmyelinated and branch extensively. Neurogliaform cells in layer II are essentially similar to those in layer I.
5. Semilunar cells, located in layer IIa, lack a basal dendritic tree, but they have several apical dendrites that arborize in layer I and axons that extend deep to the somata.
6. Multipolar cells, located throughout layer III and the endopiriform nucleus (Tseng and Haberly, 1989), have dendrites that are confined to layer III. The axons of these cells collateralize and synapse extensively within layer III and also extend to layer I. Many multipolar cells are GABAergic (Haberly et al., 1987).

In parallel with their diverse anatomical makeup, interneurons in PC are neurochemically heterogeneous, and they are thought to subservise multiple functions. CR and CCK also colocalize in GABAergic interneurons in PC (Morales and Bloom, 1997). Many GABAergic interneurons, described as “basket cells,” colocalize with Ca²⁺ binding proteins (PV, calbindin), VIP, or CCK (Kubota and Jones, 1993; Ekstrand et al., 2001a). Basket cells are primarily distributed in layers II/III and exhibit diverse molecular markers and morphological characteristics. These cells are thought to predominantly form axo-somatic or proximal axo-dendritic synapses with neurons in layers II and III, and to participate both in feedback and feedforward inhibitory circuits.

5.3 PC Circuitry

5.3.1 Input from MOB

The output of MOB is through mitral cells and tufted cells. Their main axons gather at the caudo-lateral extent of MOB forming the LOT. These caudally directed axons give off collaterals in the AON and in other regions of POC (► [Figure 6-7](#)). The MOB sends a projection to the entire extent of PC, terminating in the superficial half of layer I, which is layer Ia. Within layer Ia, mitral/tufted cell axons synapse on the apical dendrites of pyramidal cells and with certain types of interneurons (Haberly, 1983). At the ultrastructural

level, mitral/tufted cell axon terminals form asymmetrical synapses with dendritic spines. Some of the outputs of the MOB have a modest degree of topographical organization. For example, neurons in the dorsolateral quadrant of MOB project to the dorsal part of the external subdivision of AON, whereas output cells of the ventral half of MOB project to the lateral subdivision (Schoenfeld and Macrides, 1984). Intracellular HRP injections into mitral cells show that their axons form collateral terminal arbors within AON and PC (Ojima et al., 1984). The terminal arbors have a patchy anterior–posterior distribution in layer Ia of AON and PC in rabbit (Ojima et al., 1984). Some mitral cells branch and project to both PC and the olfactory tubercle. Mitral cells that are close together are reported to have similar patterns of axonal projections to the olfactory cortex (Buonviso et al., 1991). These findings, as well as observation that ORNs bearing the same odor receptor converge on single glomeruli, suggested that mitral cells of the same glomerulus might terminate with topographic specificity in PC. Studies by Buck and colleagues (Zou et al., 2001), using a genetic transneuronal tract tracing approach, reported that mitral/tufted cells associated with the same glomerulus projected to patches in PC. Such findings suggest that the representation of glomerular input to PC may have a higher degree of topographical organization than previously suspected. However, odor responses in PC, as determined by Fos mapping, do not show discrete patches, perhaps indicating that a restricted afferent input is distributed by intracortical processing (Illig and Haberly, 2003).

5.3.1.1 Neurochemistry of MOB Inputs The MOB projection to PC, via the mitral/tufted cells, is glutamatergic. Bulbectomy reduces evoked release of aspartate and NAG in PC (Collins and Probett, 1981a; Ffrench-Mullen et al., 1985) and stimulation of LOT induces glutamate and aspartate release in PC (Collins and Probett, 1981b). Receptor localization studies indicate that ionotropic glutamate and mGluRs are present in PC (🔗 [Table 6-6](#)). Stimulation of mitral/tufted cell axons in the LOT produces monosynaptic depolarization and spiking in superficial pyramidal cells. LOT-evoked excitation of PC is also blocked by DNQX (Collins and Buckley, 1989), a potent postsynaptic antagonist of kainate and AMPA receptors. By contrast, selective NMDA receptor antagonists do not reduce LOT-evoked monosynaptic excitation of PC cells. LOT-evoked responses in PC are attenuated by 2-amino-4-phosphonobutyric acid (AP4) (Collins, 1982; Hori et al., 1982; Ffrench-Mullen et al., 1985, 1986; Hasselmo and Bower, 1991), an agonist of Group III mGluRs. The AP4-induced suppression appears to be mediated by presynaptic inhibition of glutamate release from mitral/tufted cell axon terminals (Hasselmo and Bower, 1991), consistent with anatomical evidence that these cells express Group III mGluRs on their axon terminals. Collins and Richards (1990) reported that protein kinase inhibitors reduce LOT-evoked monosynaptic excitation of PC, further supporting a role for a metabotropic, second messenger-mediated regulation of excitatory transmission at this synapse.

5.3.2 Intrinsic and Association Connections

PC has extensive connections, including (1) intrinsic, local, short translaminal connections between neurons in different layers of the PC, and (2) associative, longer-range, rostrocaudally directed connections with different parts of the PC.

5.3.2.1 Intrinsic or Local Connections There are extensive translaminal connections from superficial to deeper layers and vice versa. Layer II pyramidal cell axon collaterals synapse with deeper layer III pyramidal cells and with local inhibitory interneurons in layers I and II. Local collaterals from deeper pyramidal cells synapse with local interneurons or with more superficial pyramidal cells. Local projections from pyramidal cells to GABAergic interneurons play an important role in regulating PC neuronal excitability, including the expression of LTP and seizures, via feedback and feedforward inhibition (Ekstrand et al., 2001a).

5.3.2.2 Association Connections Cortico-cortical projections within PC are extensive and exhibit laminar and regional organization (Haberly and Price, 1978a, b; Luskin and Price, 1983). Axons from pyramidal cells of layer IIb are primarily directed at more caudal sites in PC, whereas pyramidal cells in layer III project predominantly to rostral parts of PC. Overall, the rostral-to-caudal associational connections are

heavier than the caudal-to-rostral directed counterparts. Both systems terminate more heavily within the lateral than the medial parts of PC (Datiche et al., 1996). The contralateral PC also receives commissural fibers that arise from layer IIb of the anterior parts of PC. All of these association fiber systems terminate in a highly laminar fashion in layer Ib, immediately below the zone that contains the afferent input from the MOB; a lighter projection terminates in layer III.

5.3.2.3 Features and Neurochemistry of a Simple Model of PC Network Operation The orderly laminar structure of the PC and the segregation of MOB and association fiber inputs in layer I facilitate the analysis of PC's physiological organization. Haberly and others have provided a detailed analysis of the site and sequence of synaptic responses mediated by afferent inputs (i.e., LOT) and associative circuits (Haberly and Shepherd, 1973; Satou et al., 1983a, b, c; Haberly and Bower, 1984; Rodriguez and Haberly, 1989). LOT-shocks, or shocks applied to layer Ib to activate association fibers, result in a volley of activity that sweeps sequentially through PC. Field-potential recording studies suggest that there is little spatial topography to the pattern of activity following LOT or association fiber shocks. Thus, activity spreads diffusely from the site of activation, both rostrally and caudally. Single LOT shocks elicit a field potential exhibiting monosynaptic (A1) and disynaptic (B1) excitatory components (Haberly and Shepherd, 1973; Rodriguez and Haberly, 1989). The A1 component reflects monosynaptic excitation of pyramidal cell apical dendrites by LOT fibers. The B1 component reflects disynaptic excitation of pyramidal cells mediated by association fibers. The A1 and B1, mono- and disynaptic components are mediated by glutamate acting at both AMPA/kainate and NMDA receptors (Collins, 1982; Collins and Buckley, 1989). Baclofen, a GABA_B receptor agonist, inhibits responses elicited by association fibers, but not LOT fibers, via a presynaptic mechanism (Tang and Hasselmo, 1994). Feedback and feedforward inhibitory circuits within PC suppress repetitive firing from pyramidal cells, and hence, reactivation of intracortical excitatory circuits. Fast (GABA_A) and slow (GABA_B) feedforward inhibitory systems independently regulate the excitability of the pyramidal cell somata and apical dendrites (Tseng and Haberly, 1988; Kanter et al., 1996; Kapur et al., 1997). Selective blockade of the apical-dendritic inhibition enhances NMDA receptor-dependent afferent synaptic responses in pyramidal cells (Kanter et al., 1996). Computational modeling suggests that inhibitory circuits in PC operate to reiterate patterned oscillatory activity from the bulb. Slow GABA_B receptor-mediated inhibition is thought to tune PC activity to the 3–10 Hz θ range while fast GABA_A receptor-mediated inhibition provides tuning in the 40–50 Hz γ range (Wilson and Bower, 1992). Thus, at least part of the oscillatory activity emerges from intrinsic properties of intracortical inhibitory circuits.

5.4 Extrinsic Outputs of PC

Outputs of PC can be categorized into three main classes, which include: (1) MOB, (2) cortical structures, and (3) subcortical structures (▶ [Figure 6-7](#)).

5.4.1 Feedback to the MOB

As noted earlier, PC and other structures of POC project heavily back to the MOB (▶ [Figure 6-7](#)). These feedback projections are heavier from the rostral than the caudal parts of POC (Shipley and Adamek, 1984) and arise mainly from pyramidal neurons in layer II and to a lesser extent in layer III.

5.4.2 Cortical Projections

There are also direct projections from PC to insular and orbital cortex. Insular and orbital cortices are also the primary cortical targets of ascending pathways arising in the nucleus of the solitary tract in the medulla and appear to contain the primary cortical representations for both gustatory and visceral sensations. Thus,

olfactory projections to insular and orbital cortex may form a part of the circuitry that integrates olfactory and gustatory signals to generate the integrated perception of flavor.

5.4.3 Subcortical Projections

There are direct olfactory projections to the hypothalamus from neurons in the deepest layers of PC that terminate most heavily in the lateral hypothalamic area. Other components of PC project to medial and anterior parts of the hypothalamus and to the thalamus (Benjamin et al., 1982; Price and Slotnick, 1983).

5.5 Amino Acid Receptor Distribution in PC

5.5.1 Glutamate Receptors

Receptor localization studies indicate that layer II of PC contains an extensive amount of AMPA and kainate receptor subtypes, whereas layers Ia and II stain for NMDA receptors (Monaghan et al., 1985; Gall et al., 1990; Petralia and Wenthold, 1992; Wisden and Seeburg, 1993; Molnar et al., 1993; Petralia et al., 1994a, b). Receptors for mGluR1, mGluR2/3, mGluR5, mGluR7, and mGluR8 are present in PC (Shigemoto et al., 1992; Kinzie et al., 1995, 1997; Romano et al., 1995; Saugstad et al., 1997; Kinoshita et al., 1998; Wada et al., 1998). Pyramidal cells express mGluR1 (Shigemoto et al., 1992). Consistent with evidence that mGluR7-8 are presynaptically located on mitral/tufted cell axons, these receptors are densely expressed in layer Ia where these axons terminate (Kinzie et al., 1997; Kinoshita et al., 1998; Wada et al., 1998), and expression of mGluR7-8 was markedly decreased after bulbectomy or LOT transection. At the EM level, mGluR7 and 8 were found to be colocalized on axon terminals in layer Ia (Wada et al., 1998) (▶ [Table 6-6](#)).

5.5.2 GABA Receptors

GABA_A receptors are located in layers I and III of PC (Young and Kuhar, 1980b; Palacios et al., 1981b; Bowery et al., 1987). Although studies have shown intense staining for GABA_A receptors, there is only a very weak signal GABA_B receptors (Bowery et al., 1987).

5.6 Neuromodulatory Inputs to PC

5.6.1 Cholinergic Inputs to PC

5.6.1.1 Cholinergic Innervation Pattern and Receptor Distribution Cholinergic inputs to PC arise from NDB cells that are codistributed among, but are distinct from, those that project to MOB. Thus, the cholinergic inputs to MOB and PC originate from separate populations of NDB neurons. The distribution of cholinergic inputs along the rostrocaudal axis of PC is fairly homogenous (Lysakowski et al., 1989). Layer I receives a sparse cholinergic innervation, whereas layers II and III receive a moderate and fairly uniform cholinergic innervation; the density of fibers is somewhat heavier in layer II than III. All four muscarinic subtypes (M₁, M₂, M₃, and M₄) are present in layer II (Rotter et al., 1979; Spencer et al., 1986; Buckley et al., 1988). The M₁ receptor subtype has also been localized on dendritic spines in layer I, but the cell type was not specified (Levey et al., 1991). Nicotinic cholinergic receptors are located in layers II and III, but as with the muscarinic receptors, the cellular and dendritic locations of these receptors are not known (Sahin et al., 1992; Hill et al., 1993; Seguella et al., 1993) (▶ [Table 6-6](#)).

5.6.1.2 Cholinergic Actions In PC slices, ACh and muscarinic cholinergic agonists suppress intrinsic (layer Ib) fiber transmission without affecting transmission at afferent (layer Ia) fiber synapses (Hasselmo and

Bower, 1992; Hasselmo et al., 1992b; Hasselmo et al., 1997). This suppression is presynaptically mediated by the M1 muscarinic subtype. Cholinergic agonists also appear to directly increase the excitability of pyramidal neurons, increasing neuronal bursts induced by intracellular depolarization. This effect is mediated, in part, by blockade of the slow afterhyperpolarization mediated by a Ca^{2+} -dependent K^{+} current. ACh, acting at muscarinic receptors, has been shown to reduce the M-current in PC pyramidal neurons (Constanti and Galvan, 1983), and to depolarize and increase the firing rate of pyramidal cells (Libri et al., 1994; Postlewaithe et al., 1998). In the later study, muscarinic receptor stimulation was also reported to induce a slow poststimulus afterdepolarization and to depress EPSPs elicited by association fibers. Activation of the NDB, which is the source of cholinergic projections to PC, in vivo increased the spontaneous firing rate of PC cells, and it also increased the disynaptic excitatory (B1), and decreased the disynaptic inhibitory (P2), field-potential components evoked by LOT stimulation (Zimmer et al., 1999). NDB stimulation decreased the P2 component following activation of association fibers in caudal PC, and also reduced the paired-pulse inhibition of the P2 component following LOT and caudal PC shocks. These effects were reversed by scopolamine, suggesting the involvement of muscarinic receptors. These results suggest that activation of cholinergic inputs to PC increases the excitability of pyramidal cells, probably by a disinhibitory mechanism. A subsequent in vivo study reported similar findings: NDB stimulation enhanced the late, disynaptic component of the evoked potential elicited by LOT stimulation, but caused a suppression of the synaptic potential elicited by stimulation of the posterior PC; both effects were antagonized by muscarinic receptor blockers (Linster et al., 1999). Other in vivo experiments utilizing optical imaging reported contrasting results suggesting that NDB stimulation reduced both mono- and disynaptic responses elicited by LOT stimulation; the mechanism and transmitter involved in these effects were not studied (Rosin et al., 1999). More recent in vivo experiments showed that topical application of muscarinic receptor antagonists did not alter spontaneous or odor-evoked activity of anterior PC cells (Wilson, 2001). However, these antagonists enhanced odor-evoked cross-habituation, perhaps indicating that ACh acts to enhance generalization between odor representations in PC.

5.6.2 Dopaminergic Inputs to PC


5.6.2.1 DA Innervation Pattern, Receptor Distribution, and Physiological Actions DA inputs to PC arise from several sources, including the substantia nigra, ventral tegmental area and, to a lesser extent, the rostromedial part of dorsal raphe nucleus (Datiche and Cattarelli, 1996). The dopaminergic innervation of PC exhibits a marked rostrocaudal gradient and laminar specificity (Datiche and Cattarelli, 1996). Rostrally, DA fibers are relatively sparse and primarily confined to layers II/III (Figure 6-11). Along the rostral-to-caudal axis of PC, the density of innervation progressively increases and DA fibers invade more superficial layers of PC. A moderately dense plexus of DA fibers extends from the deep part of layer I through layer III, to the caudal limit of PC. Autoradiographic, in situ hybridization and immunocytochemical studies have revealed the presence of D1 receptors in layer II (Palacios et al., 1981a; Mansour et al., 1990a, b; Fremeau et al., 1991; Huang et al., 1992). To date, there is little definitive evidence concerning the presence of D2 receptors in PC. Relatively little is known about DA functions in PC. DA was reported to excite PC interneurons, leading to increased IPSC activity in pyramidal cells (Gellman and Aghajanian, 1993, 1994).

5.6.3 Noradrenergic Inputs to PC

5.6.3.1 Noradrenergic Innervation Pattern and Receptor Distribution Noradrenergic LC neurons project heavily to PC and represent the major source of NE input to this structure (Fallon and Loughlin, 1982; Datiche and Cattarelli, 1996). LC lesions decrease NE levels in PC by 77% (Fallon and Moore, 1978). Like the projection to the neocortex, the LC projection to PC is primarily ipsilateral with a small contralateral component. Early histofluorescence studies (Fallon and Moore, 1978) of monoaminergic innervation of PC

■ Figure 6-11

Neuromodulatory transmitter inputs to PC. Darkfield photomicrographs showing the distribution of dopaminergic (a), noradrenergic (b), and serotonergic (c) fibers revealed respectively with immunohistochemistry for tyrosine hydroxylase (TH), dopamine- β -hydroxylase (DBH), and serotonin (5-HT). Abbreviations: Endo, endopiriform nucleus; lot, lateral olfactory tract. Reprinted from Handbook of Chem. Neuroanat. Integrated Sys. CNS, Vol. 12, Part III, Chapter III, The Olfactory System, M. Shipley et al., pp. 469–573, 1996, with permission from Elsevier, Ltd



could not unambiguously distinguish among 5-HT, DA, and NE fibers. Immunohistochemistry for the NE biosynthetic enzyme dopamine- β -hydroxylase (DBH) showed that layers Ia and III of PC contained a moderate plexus of NE fibers; layer II was sparsely innervated (Figure 6-11) (Datiche and Cattarelli, 1996). A distinctive feature of NE innervation of PC is the long fibers oriented primarily parallel to the pial surface in layer Ia. The density and laminar distribution of NE fibers are relatively uniform along the rostrocaudal axis of PC (Datiche and Cattarelli, 1996).

Both α and β receptors are present in PC (Table 6-6). α 1-receptors are highly expressed in layers II/III (Sargent-Jones et al., 1985) and layer II pyramidal cells express high levels of α 1_A, α 1_B, and low levels of α 1_D (Pieribone et al., 1994; Day et al., 1997; Domyancic and Morilak, 1997). High levels of α 2 receptor binding sites are present in PC (Young and Kuhar, 1979, 1980a; Unnerstall et al., 1984; Winzer-Serhan et al., 1997a), and α 2_A and α 2_C appear to be expressed by layer II pyramidal cells (Nicholas et al., 1993; Rosin et al., 1996; Talley et al., 1996; Winzer-Serhan et al., 1997a, b); α 2_C receptors appear to be expressed by neurons in all layers of PC (Rosin et al., 1996). Weak-to-moderate expression of β receptor subtypes has also been reported (Palacios and Kuhar, 1982; Wanaka et al., 1989).

5.6.3.2 Physiological Actions of NE Early field-potential studies reported that low concentrations of NE (0.1–5 μ M) enhanced, and high concentrations (20–250 μ M) suppressed, LOT-evoked responses in PC (Collins et al., 1984). These effects were blocked by both α and β receptor antagonists. A subsequent study reported uniform suppression of LOT-evoked field potentials by NE (1–100 μ M), although the effects were greater in layer Ib than Ia (Hasselmo et al., 1997). These effects were similar to those elicited by carbachol. The cellular actions of NE on PC neurons have received relatively little attention. NE was shown to block the slow afterhyperpolarization in guinea-pig pyramidal cells (Constanti and Sim, 1987). NE was reported to excite putative interneurons located at the layer II/III border, which in turn, increased IPSC activity in pyramidal cells in vitro (Sheldon and Aghajanian, 1990; Gellman and Aghajanian, 1993, 1994). Other studies from the same laboratory indicated that the NE-induced interneuronal excitation was mediated by α 1_B receptors (Marek and Aghajanian, 1996a). LC activation in vivo was found to enhance odor-evoked responses in anterior and posterior PC (Bouret and Sara, 2002).

5.6.4 Serotonergic Inputs to PC

5.6.4.1 5-HT Innervation Pattern and Receptor Distribution Anterograde and retrograde tracing studies have demonstrated a rich projection from the dorsal raphe nucleus to PC (De Olmos and Heimer, 1980; Vertes, 1991; Datiche et al., 1995). Ascending serotonergic fibers from the dorsal raphe terminating in PC are believed to travel in the ventrolateral aspect of the medial forebrain bundle (Azmitia and Segal, 1978). Anterograde labeling of ascending dorsal raphe axons demonstrated that the entire PC was targeted by raphe projections (Vertes, 1991; Datiche et al., 1996). The projection was reported to be heavier to rostral than caudal PC, and heavier to the deeper than the superficial layers. The transmitter of labeled fibers could not be identified in these anterograde tracing results. The serotonergic innervation of PC is very heavy by comparison to DA and NE. 5-HT fibers are especially heavy in layers I and III and in the endopiriform nucleus (Figure 6-11) (Datiche et al., 1996). The density of 5-HT fibers progressively decreases in the deeper parts of layer III. Layer II by contrast, is sparsely innervated.

Receptor binding and in situ hybridization studies suggest that 5-HT_{1A} receptors are located on the dendrites of pyramidal cells (layer II) and also on intrinsic cells of layer III (Pompeiano et al., 1992) (Table 6-6). 5-HT_{2A} and 5-HT_{2C} receptors are strongly expressed by layer II pyramidal cells, as well as by interneurons in layers II and III (Pompeiano et al., 1994; Hamada et al., 1998; Cornea-Hebert et al., 1999). The 5-HT₃ subtype is present in PC and colocalizes with diverse interneuronal subtypes (Tecott et al., 1993; Morales and Bloom, 1997). 5-HT₃ colocalizes with approximately 50% of CCK-positive interneurons, and it is also expressed by interneurons containing GABA or CR, but not PV (Morales and Bloom, 1997).

5.6.4.2 Physiological Actions of 5-HT 5-HT has several actions on PC pyramidal cells as well as on putative interneurons that are mediated by distinct 5-HT receptor subtypes (Sheldon and Aghajanian, 1990, 1991; Gellman and Aghajanian, 1994). 5-HT has mixed, but predominately excitatory actions on layer II pyramidal cells. This excitation appears to be mediated by a reduction of the M-current, a noninactivating voltage-dependent outward K⁺ current, as well as a decrease in a Ca²⁺-activated K⁺ current. These actions appear to be mediated via 5-HT_{1C} receptors. 5-HT also inhibits and increases the frequency of IPSPs in other pyramidal cells. This is mediated by excitation of layer III GABAergic interneurons that are directly activated (i.e., depolarized) by 5-HT acting at the 5-HT_{2A} receptor subtype (Gellman and Aghajanian, 1994; Marek and Aghajanian, 1994, 1996b). Interneurons excited by 5-HT appear to be a distinct subset distinguished by a relatively depolarized resting membrane potential, higher input resistance, and short action potential duration. The 5-HT_{2A} receptor-mediated effects on interneurons were not blocked, but instead were enhanced, by inhibitors of PKC, suggesting that PKC negatively modulates the excitation of PC interneurons (Marek and Aghajanian, 1995). In vivo, layer III putative interneurons were inhibited by 5-HT_{2A} agonists (Bloms-Funke et al., 1999). It is unclear if and how these cells correspond to those studied in vitro by Aghajanian and colleagues.

Acknowledgments

We acknowledge grant support for portions of this chapter that refer to research of the authors: PHS Grants DC03195, DC06356, DC07123, and RR020146.

References

- Allen DM, Hamilton KA. 2000. Ultrastructural identification of synapses between mitral/tufted cell dendrites. *Brain Res* 860: 170-173.
- Allison A. 1949. Quantitative observations on the olfactory system of the rabbit. *Brain* 72: 186-197.
- Allison A. 1953. The morphology of the olfactory system in the vertebrates. *Biol Rev Camb Philos Soc* 28: 195-244.
- Alonso JR, Arévalo R, Garcia-Ojeda E, Porteros A, Briñón JG, et al. 1995. NADPH-diaphorase active and calbindin D-28k-immunoreactive neurons and fibers in the olfactory

- bulb of the hedgehog (*Erinaceus europaeus*). *J Comp Neurol* 351: 307-327.
- Alonso JR, Arévalo R, Porteros A, Briñón JG, Lara J, et al. 1993. Calbindin D-28K and NADPH-diaphorase activity are localized in different populations of periglomerular cells in the rat olfactory bulb. *J Chem Neuroanat* 6: 1-6.
- Alonso JR, Briñón JG, Crespo C, Bravo IG, Arévalo R, et al. 1998. Chemical organization of the macaque monkey olfactory bulb: II. Calretinin, calbindin D-28k, parvalbumin, and neurocalcin immunoreactivity. *J Comp Neurol* 432: 389-407.
- Altman J. 1969. Autoradiographic and histological studies of postnatal neurogenesis. IV. Cell proliferation and migration in the anterior forebrain, with special reference to persisting neurogenesis in the olfactory bulb. *J Comp Neurol* 137: 433-457.
- Aroniadou-Anderjaska V, Ennis M, Shipley MT. 1997. Glomerular synaptic responses to olfactory nerve input in rat olfactory bulb slices. *Neuroscience* 79: 425-434.
- Aroniadou-Anderjaska V, Ennis M, Shipley MT. 1999a. Current-source density analysis in the rat olfactory bulb: Laminar distribution of kainate/AMPA and NMDA receptor-mediated currents. *J Neurophysiol* 81: 15-28.
- Aroniadou-Anderjaska V, Ennis M, Shipley MT. 1999b. Dendrodendritic recurrent excitation in mitral cells of the rat olfactory bulb. *J Neurophysiol* 82: 489-494.
- Aroniadou-Anderjaska V, Zhou FM, Priest CA, Ennis M, Shipley MT. 2000. Tonic and synaptically evoked presynaptic inhibition of sensory input to the rat olfactory bulb via GABA (B) heteroreceptors. *J Neurophysiol* 84: 1194-1203.
- Astic L, Saucier D. 1981. Metabolic mapping of functional activity in the olfactory projections of the rat: Ontogenetic study. *Brain Res* 254: 141-156.
- Astic L, Saucier D. 1986. Anatomical mapping of the neuroepithelial projection to the olfactory bulb in the rat. *Brain Res Bull* 16: 445-454.
- Astic L, Saucier D, Holley A. 1987. Topographical relationships between olfactory receptor cells and glomerular foci in the rat olfactory bulb. *Brain Res* 424: 144-152.
- Aston-Jones G, Bloom FE. 1981. Norepinephrine-containing locus coeruleus neurons in behaving rats exhibit pronounced responses to non-noxious environmental stimuli. *J Neurosci* 1: 887-900.
- Aungst JL, Heyward PM, Puche AC, Karnup SV, Hayar A, et al. 2003. Center-surround inhibition among olfactory bulb glomeruli. *Nature* 426: 623-629.
- Azmitia EC, Segal M. 1978. An autoradiographic analysis of the differential ascending projections of the dorsal and median raphe nuclei in the rat. *J Comp Neurol* 179: 641-667.
- Bailey MS, Shipley MT. 1993. Astrocyte subtypes in the rat olfactory bulb: Morphological heterogeneity and differential laminar distribution. *J Comp Neurol* 328: 501-526.
- Baimbridge KG, Miller JJ. 1982. Immunohistochemical localization of calcium-binding protein in the cerebellum, hippocampal formation and olfactory bulb of the rat. *Brain Res* 245: 223-229.
- Baker H. 1986. Species differences in the distribution of substance P and tyrosine hydroxylase immunoreactivity in the olfactory bulb. *J Comp Neurol* 252: 206-226.
- Baker H. 1990. Unilateral, neonatal olfactory deprivation alters tyrosine hydroxylase expression but not aromatic amino acid decarboxylase or GABA immunoreactivity. *Neuroscience* 36: 761-771.
- Baker H, Farbman AI. 1993. Olfactory afferent regulation of the dopamine phenotype in the fetal rat olfactory system. *Neuroscience* 52: 115-134.
- Baker H, Kawano T, Margolis FL, Joh TH. 1983. Trans-neuronal regulation of tyrosine hydroxylase expression in olfactory bulb of mouse and rat. *J Neurosci* 3: 69-78.
- Baker H, Morel K, Stone DM, Maruniak JA. 1993. Adult naris closure profoundly reduces tyrosine hydroxylase expression in mouse olfactory bulb. *Brain Res* 614: 109-116.
- Baker H, Cummings DM, Munger SD, Margolis JW, Franzen L, et al. 1999. Targeted deletion of a cyclic nucleotide-gated channel subunit (OCNC1): Biochemical and morphological consequences in adult mice. *J Neurosci* 19: 9313-9321.
- Baker H, Kawano T, Albert V, Joh TH, Reis DJ, et al. 1984. Olfactory bulb dopamine neurons survive deafferentation-induced loss of tyrosine hydroxylase. *Neuroscience* 11: 605-615.
- Baker H, Towle AC, Margolis FL. 1988. Differential afferent regulation of dopaminergic and GABAergic neurons in the mouse main olfactory bulb. *Brain Res* 450: 69-80.
- Baker H, Liu N, Chun HS, Saino S, Berlin R, et al. 2001. Phenotypic differentiation during migration of dopaminergic progenitor cells to the olfactory bulb. *J Neurosci* 21: 8505-8513.
- Banay-Schwartz M, Lajtha A, Palkovits M. 1989a. Changes with aging in the levels of amino acids in rat CNS structural elements: I. glutamate and related amino acids. *Neurochem Res* 14: 555-562.
- Banay-Schwartz M, Lajtha A, Palkovits M. 1989b. Changes with aging in the levels of amino acids in rat CNS structural elements: II. taurine and small neutral amino acids. *Neurochem Res* 14: 563-570.
- Banks MI, Haberly LB, Jackson MB. 1996. Layer-specific properties of the transient K current (Ia) in piriform cortex. *J Neurosci* 16: 3862-3876.

- Bardoni R, Magherini PC, Belluzzi O. 1996. Excitatory synapses in the glomerular triad of frog olfactory bulb in vitro. *Neuroreport* 7: 1851-1855.
- Bassett JL, Shipley MT, Foote SL. 1992. Localization of corticotropin-releasing factor-like immunoreactivity in monkey olfactory bulb and secondary olfactory areas. *J Comp Neurol* 316: 348-362.
- Bastianelli E, Okazaki K, Hidaka H, Pochet R. 1993. Neurocalcin immunoreactivity in rat olfactory bulb. *Neurosci Lett* 161: 165-168.
- Bayer SA. 1983. 3H-Thymidine-radiographic studies of neurogenesis in the rat olfactory bulb. *Exp Brain Res* 50: 329-340.
- Behan M, Haberly LB. 1999. Intrinsic and efferent connections of the endopiriform nucleus in rat. *J Comp Neurol* 408: 532-548.
- Belluardo N, Mudò G, Trovato-Salinaro A, Le Gurun S, Charollais A, et al. 2000. Expression of connexin36 in the adult and developing rat brain. *Brain Res* 865: 121-138.
- Belluscio L, Katz LC. 2001. Symmetry, stereotypy, and topography of odorant representations in mouse olfactory bulbs. *J Neurosci* 21: 2113-2122.
- Belluscio L, Gold GH, Nemes A, Axel R. 1998. Mice deficient in G(olf) are anosmic. *Neuron* 20: 69-81.
- Belluscio L, Lodovichi C, Feinstein P, Mombaerts P, Katz LC. 2002. Odorant receptors instruct functional circuitry in the mouse olfactory bulb. *Nature* 419: 296-300.
- Belluzzi O, Benedusi M, Ackman J, Lo Turco J. 2003. Electrophysiological identification of new neurons in the olfactory bulb. *J Neurosci* 23: 10411-10418.
- Belluzzi O, Puopolo M, Benedusi M, Kratskin I. 2004. Selective neuroinhibitory effects of taurine in slices of rat main olfactory bulb. *Neuroscience* 124: 929-944.
- Benjamin RM, Jackson JC, Golden GT, West CH. 1982. Sources of olfactory inputs to opossum mediodorsal nucleus identified by horseradish peroxidase and autoradiographic methods. *J Comp Neurol* 207: 358-368.
- Benson TE, Burd GD, Greer CA, Landis DM, Shepherd GM. 1985. High-resolution 2-deoxyglucose autoradiography in quick-frozen slabs of neonatal rat olfactory bulb. *Brain Res* 339: 67-78.
- Berkowicz DA, Trombley PQ. 2000. Dopaminergic modulation at the olfactory nerve synapse. *Brain Res* 855: 90-99.
- Biffo S, Grillo M, Margolis FL. 1990. Cellular localization of carnosine-like and anserine-like immunoreactivities in rodent and avian central nervous system. *Neuroscience* 35: 637-651.
- Blakely RD, Ory-Lavallee L, Grzanna R, Koller KJ, Coyle JT. 1987. Selective immunocytochemical staining of mitral cells in rat olfactory bulb with affinity purified antibodies against N-acetyl-aspartyl-glutamate. *Brain Res* 402: 373-378.
- Blakemore LJ, Trombley PQ. 2003. Kinetic variability of AMPA receptors among olfactory bulb neurons in culture. *Neuroreport* 14: 965-970.
- Blakemore LJ, Trombley PQ. 2005. Slice electrophysiology and histological evidence of calcium-fluxing AMPA receptors in the olfactory bulb. *ACHEM S Abstr* 75.
- Blanes T. 1898. Sobre algunos puntos dudosos de la estructura del bulbo olfactorio. *Revista Trimestral micrografica* 3: 99-127.
- Bloms-Funke P, Gernert M, Ebert U, Loscher W. 1999. Extracellular single-unit recordings of piriform cortex neurons in rats: Influence of different types of anesthesia and characterization of neurons by pharmacological manipulation of serotonin receptors. *J Neurosci Res* 55: 608-619.
- Bogan N, Brecha N, Gall C, Karten HJ. 1982. Distribution of enkephalin-like immunoreactivity in the rat main olfactory bulb. *Neuroscience* 7: 895-906.
- Bonino M, Cantino D, Sassoe-Pognetto M. 1999. Cellular and subcellular localization of gamma-aminobutyric acidB receptors in the rat olfactory bulb. *Neurosci Lett* 274: 195-198.
- Bonnemann C, Holland A, Meyer DK. 1989. Cholecystokinin in rat olfactory bulb is mainly contained in intrinsic neurons. *Neurosci Lett* 103: 281-286.
- Booze RM, Crisostomo EA, Davis JN. 1989. Species differences in the localization and number of CNS beta adrenergic receptors: Rat versus guinea pig. *J Pharmacol Exp Ther* 249: 911-920.
- Bouna S, Gysling K, Calas A, Araneda S. 1994. Some noradrenergic neurons of locus ceruleus-olfactory pathways contain neuropeptide-Y. *Brain Res Bull* 34: 413-417.
- Bouret S, Sara SJ. 2002. Locus coeruleus activation modulates firing rate and temporal organization of odour-induced single-cell responses in rat piriform cortex. *Eur J Neurosci* 16: 2371-2382.
- Bowery NG, Hudson AL, Price GW. 1987. GABAA and GABAB receptor site distribution in the rat central nervous system. *Neuroscience* 20: 365-383.
- Boyd JD, Delaney KR. 2002. Tyrosine hydroxylase-immunoreactive interneurons in the olfactory bulb of the frogs *Rana pipiens* and *Xenopus laevis*. *J Comp Neurol* 454: 42-57.
- Bradley J, Reuter D, Frings S. 2001. Facilitation of calmodulin-mediated odor adaptation by cAMP-gated channel subunits. *Science* 294: 2176-2178.
- Breer H. 2003. Olfactory receptors: Molecular basis for recognition and discrimination of odors. *Anal Bioanal Chem* 377: 427-433.
- Brennan P, Kaba H, Keverne EB. 1990. Olfactory recognition: A simple memory system. *Science* 250: 1223-1226.
- Briñón JG, Alonso JR, Arévalo R, García-Ojeda E, Lara J, et al. 1992. Calbindin D-28k-positive neurons in the rat olfactory

- bulb. An immunohistochemical study. *Cell Tissue Res* 269: 289-297.
- Briñón JG, Martínez-Guijarro FJ, Bravo IG, Arévalo R, Crespo C, et al. 1999. Coexpression of neurocalcin with other calcium-binding proteins in the rat main olfactory bulb. *J Comp Neurol* 407: 404-414.
- Briñón JG, Weruaga E, Crespo C, Porteros Á, Arévalo R, et al. 2001. Calretinin-, neurocalcin-, and parvalbumin-immunoreactive elements in the olfactory bulb of the hedgehog (*Erinaceus europaeus*). *J Comp Neurol* 429: 554-570.
- Brunet LJ, Gold GH, Ngai J. 1996. General anosmia caused by a targeted disruption of the mouse olfactory cyclic nucleotide-gated cation channel. *Neuron* 17: 681-693.
- Brüning I, Sommer M, Hatt H, Bormann J. 1999. Dopamine receptor subtypes modulate olfactory bulb gamma-aminobutyric acid type A receptors. *Proc Natl Acad Sci USA* 96: 2456-2460.
- Brunjes PC. 1994. Unilateral naris closure and olfactory system development. *Brain Res Rev* 19: 146-160.
- Brunjes PC, Armstrong AM. 1996. Apoptosis in the rostral migratory stream of the developing rat. *Dev Brain Res* 92: 219-222.
- Brunjes PC, Smith-Crafts LK, McCarty R. 1985. Unilateral odor deprivation: Effects on the development of olfactory bulb catecholamines and behavior. *Dev Brain Res* 22: 1-6.
- Buck L, Axel R. 1991. A novel multigene family may encode odorant receptors: A molecular basis for odor recognition. *Cell* 65: 175-187.
- Buckley NJ, Bonner TI, Brann MR. 1988. Localization of a family of muscarinic receptor mRNAs in rat brain. *J Neurosci* 8: 4646-4652.
- Buonviso N, Reval MF, Jourdan F. 1991. The projections of mitral cells from small local regions of the olfactory bulb: An anterograde tracing study using PHA-L (*Phaseolus vulgaris leucoagglutinin*). *Eur J Neurosci* 3: 493-500.
- Burd GD, Davis BJ, Macrides F, Grillo M, Margolis F. 1982. Carnosine in primary afferents of the olfactory system: An autoradiographic and biochemical study. *J Neurosci* 2: 244-255.
- Cajal RSy. 1890. The origins and terminations of the olfactory, optic, and acoustic nerves in vertebrates. In studies on the cerebral cortex [Limbic structures]. Translated by Kraft LM (1955). Chicago, IL: The Year Book Publishers; pp. 1-27.
- Cajal RSy. 1911a. Olfactory apparatus: Olfactory mucosa and olfactory bulb or first-order olfactory center. *Histology of the nervous system*, Vol. II. Translated by Swanson N, Swanson L (1995). New York: Oxford Univ. Press; pp. 532-554.
- Cajal RSy. 1911b. *Histologie du Systeme Nerveux de l'Homme et des Vertebres*. Maloine: Paris.
- Camps M, Kelly PH, Palacios JM. 1990. Autoradiographic localization of D1 and D2 receptors in the brain of several mammalian species. *J Neural Trans* 80: 105-127.
- Carleton A, Petreanu LT, Lansford R, Alvarez-Buylla A, Lledo P-M. 2003. Becoming a new neuron in the adult olfactory bulb. *Nat Neurosci* 6: 607-518.
- Carlson GC, Shipley MT, Keller A. 2000. Long-lasting depolarizations in mitral cells of the rat olfactory bulb. *J Neurosci* 20: 2011-2021.
- Carlson GC, Slawewski ML, Lancaster E, Keller A. 1997. Distribution and activation of intracellular Ca^{2+} stores in cultured olfactory bulb neurons. *J Neurophysiol* 78: 2176-2185.
- Carson KA. 1984. Quantitative localization of neurons projecting to the mouse main olfactory bulb. *Brain Res Bull* 12: 629-634.
- Carson KA, Burd GD. 1980. Localization of acetylcholinesterase in the main and accessory olfactory bulbs of the mouse by light and electron microscopic histochemistry. *J Comp Neurol* 191: 353-371.
- Casabona G, Catania MV, Storto M, Ferraris N, Perroteau I, et al. 1998. Deafferentation up-regulates the expression of the mGlu1a metabotropic glutamate receptor protein in the olfactory bulb. *Eur J Neurosci* 10: 771-776.
- Castillo PE, Carleton A, Vincent JD, Lledo PM. 1999. Multiple and opposing roles of cholinergic transmission in the main olfactory bulb. *J Neurosci* 19: 9180-9191.
- Celio MR. 1990. Calbindin D-28k and parvalbumin in the rat nervous system. *Neuroscience* 35: 375-475.
- Chanse NT, Kopin IJ. 1968. Stimulus-induced release of substances from olfactory bulb using the push-pull cannula. *Nature* 217: 466-498.
- Chao TI, Kasa P, Wolff JR. 1997. Distribution of astroglia in glomeruli of the rat main olfactory bulb: Exclusion from the sensory subcompartment of neuropil. *J Comp Neurol* 388: 191-210.
- Chen WR, Shepherd GM. 1997. Membrane and synaptic properties of mitral cells in slices of rat olfactory bulb. *Brain Res* 745: 189-196.
- Chen WR, Xiong W, Shepherd GM. 2000. Analysis of relations between NMDA receptors and GABA release at olfactory bulb reciprocal synapses. *Neuron* 25: 625-633.
- Chen AM, Perrin MH, Digruccio MR, Vaughan JM, Brar BK, et al. 2005. A soluble mouse brain spice variant of type 2 alpha corticotropin-releasing factory (CRF) binds ligands and modulates their activity. *Proc Natl Acad Sci USA* 102: 2620-2625.
- Christie JM, Westbrook GL. 2003. Regulation of back-propagating action potential in mitral cell lateral dendrites by A-type potassium currents. *J Neurophysiol* 89: 2466-2472.
- Christie JM, Schoppa NE, Westbrook GL. 2001. Tufted cell dendrodendritic inhibition in the olfactory bulb is

- dependent on NMDA receptor activity. *J Neurophysiol* 85: 169-173.
- Christie JM, Bark C, Hormuzdi SG, Helbig I, Monyer H, et al. 2005. Connexin36 mediates spike synchrony in olfactory bulb glomeruli. *Neuron* 46: 761-772.
- Cho JY, Min N, Franzen L, Baker H. 1996. Rapid down-regulation of tyrosine hydroxylase expression in the olfactory bulb of naris-occluded adult rats. *J Comp Neurol* 369: 264-276.
- Chu DCM, Albin RL, Young AB, Penney JB. 1990. Distribution and kinetics of GABAB binding sites in rat central nervous system: A quantitative autoradiographic study. *Neuroscience* 34: 341-357.
- Cielo MR. 1990. Calbindin D-28k and parvalbumin in the rat nervous system. *Neuroscience* 35: 375-475.
- Cigola E, Volpe BT, Lee JW, Franzen L, Baker H. 1998. Tyrosine hydroxylase expression in primary cultures of olfactory bulb: Role of L-type calcium channels. *J Neurosci* 18: 7638-7649.
- Cinelli AR, Hamilton KA, Kauer JS. 1995. Salamander olfactory bulb neuronal activity observed by video rate, voltage-sensitive dye imaging. III. Spatial and temporal properties of responses evoked by odorant stimulation. *J Neurophysiol* 73: 2053-2071.
- Ciombor KJ, Ennis M, Shipley MT. 1999. Norepinephrine increases rat mitral cell excitatory responses to weak olfactory nerve input via alpha-1 receptors in vitro. *Neuroscience* 90: 595-606.
- Clancy AN, Schoenfeld TA, Forbes WB, Macrides F. 1994. The spatial organization of the peripheral olfactory system of the hamster. Part II: Receptor surfaces and odorant passageways within the nasal cavity. *Brain Res Bull* 34: 211-241.
- Clemett DA, Punhani T, Duxon MS, Blackburn TP, Fone KC. 2000. Immunohistochemical localisation of the 5-HT_{2C} receptor protein in the rat CNS. *Neuropharmacol* 39: 123-132.
- Collins GG. 1974. The rates of synthesis, uptake and disappearance of [¹⁴C]-taurine in eight areas of the rat central nervous system. *Brain Res* 76: 447-459.
- Collins GG. 1982. Some effects of excitatory amino acid receptor antagonists on synaptic transmission in the rat olfactory cortex slice. *Brain Res* 244: 311-318.
- Collins GG, Buckley KS. 1989. Antagonism of monosynaptic excitations in the mouse olfactory cortex slice by 6,7-dinitroquinoxaline-2,3-dione. *Neuropharmacol* 28: 1123-1128.
- Collins GG, Probett GA. 1981a. Aspartate and not glutamate is the likely transmitter of the rat lateral olfactory tract fibres. *Brain Res* 209: 231-234.
- Collins GG, Probett GA. 1981b. Patterns of endogenous amino acid release from slices of rat guinea-pig olfactory cortex. *Brain Res* 204: 103-120.
- Collins GG, Richards WJ. 1990. Pharmacological evidence that protein kinase C modulates monosynaptic excitations in the olfactory cortex. *Naunyn Schmiedebergs Arch Pharmacol* 341: 114-122.
- Collins GG, Probett GA, Anson J, McLaughlin NJ. 1984. Excitatory and inhibitory effects of noradrenaline on synaptic transmission in the rat olfactory cortex slice. *Brain Res* 294: 211-223.
- Condorelli DF, Belluardo N, Trovato-Salimaro A, Mudò G. 2000. Expression of C_X36 in mammalian neurons. *Brain Res Rev* 32: 72-85.
- Condorelli DF, Parenti R, Spinella F, Salinaro AT, Belluardo N, et al. 1998. Cloning of a new gap junction gene (C_X36) highly expressed in mammalian neurons. *Eur J Neurosci* 10: 1202-1208.
- Conn PJ, Pin JP. 1997. Pharmacology and functions of metabotropic glutamate receptors. *Ann Rev Pharmacol Toxicol* 37: 205-237.
- Constanti A, Galvan M. 1983. M-current in voltage-clamped olfactory cortex neurons. *Neurosci Lett* 39: 65-70.
- Constanti A, Sim JA. 1987. Calcium-dependent potassium conductance in guinea-pig olfactory cortex neurons in vitro. *J Physiol* 387: 173-194.
- Cooper-Kuhn CM, Winkler J, Kuhn HG. 2004. Decreased neurogenesis after cholinergic forebrain lesion in the adult rat. *J Neurosci Res* 77: 155-165.
- Coopersmith R, Leon M. 1984. Enhanced neural response to familiar olfactory cues. *Science* 225: 849-851.
- Cornea-Hebert V, Riad M, Wu C, Singh SK, Descarries L. 1999. Cellular and subcellular distribution of the serotonin 5-HT_{2A} receptor in the central nervous system of adult rat. *J Comp Neurol* 409: 187-209.
- Coronas V, Srivastava LK, Liang JJ, Jourdan F, Moysé E. 1997. Identification and localization of dopamine receptor subtypes in rat olfactory mucosa and bulb: A combined in situ hybridization and ligand binding radioautographic approach. *J Chem Neuroanat* 12: 243-257.
- Corotto FS, Henegar JR, Maruniak JA. 1994. Odor deprivation leads to reduced neurogenesis and reduced neuronal survival in the olfactory bulb of the adult mouse. *Neuroscience* 61: 739-744.
- Couper Leo JM, Devine AH, Brunjes PC. 2000. Focal denervation alters cellular phenotypes and survival in the developing rat olfactory bulb. *J Comp Neurol* 417: 325-336.
- Crespo C, Blasco-Ibáñez JM, Marqués-Mari AI. 2001. Parvalbumin-containing interneurons do not innervate granule cells in the olfactory bulb. *Neuroreport* 12: 2553-2556.
- Crespo C, Blasco-Ibáñez JM, Marqués-Mari AI, Alonso JR, Brinon JG, et al. 2002. Vasoactive intestinal polypeptide-containing elements in the olfactory bulb of the hedgehog (*Erinaceus europaeus*). *J Chem Neuroanat* 24: 49-63.

- Cuschieri A, Bannister LH. 1975a. The development of the olfactory mucosa in the mouse: Electron microscopy. *J Anat* 119: 471-498.
- Cuschieri A, Bannister LH. 1975b. The development of the olfactory mucosa in the mouse: Light microscopy. *J Anat* 119: 277-286.
- Datiche F, Cattarelli M. 1996. Catecholamine innervation of the piriform cortex: A tracing and immunohistochemical study in the rat. *Brain Res* 671: 69-78
- Datiche F, Litaudon P, Cattarelli M. 1996. Intrinsic association fiber system of the piriform cortex: a quantitative study based on a cholera toxin B subunit tracing in the rat. *J Comp Neurol* 376: 265-277
- Datiche F, Luppi P-H, Cattarelli M. 1995. Serotonergic and non-serotonergic projections from the raphe nuclei to the piriform cortex in the rat: A cholera toxin B subunit (CTb) and 5-HT immunohistochemical study. *Brain Res* 671: 27-37.
- Davila NG, Blakemore LJ, Trombley PQ. 2003. Dopamine modulates synaptic transmission between rat olfactory bulb neurons in culture. *J Neurophysiol* 90: 395-404.
- Davis BJ. 1991. NADPH-diaphorase activity in the olfactory system of the hamster and rat. *J Comp Neurol* 314: 493-511.
- Davis BJ, Macrides F. 1983. Tyrosine hydroxylase immunoreactive neurons and fibers in the olfactory system of the hamster. *J Comp Neurol* 214: 427-440.
- Davis BJ, Burd GD, Macrides F. 1982. Localization of methionine-enkephalin, substance P, and somatostatin immunoreactivities in the main olfactory bulb of the hamster. *J Comp Neurol* 204: 377-383.
- Davison IG, Boyd JD, Delaney KR. 2004. Dopamine inhibits mitral/tufted-granule cell synapses in the frog olfactory bulb. *J Neurosci* 24: 8057-8067.
- Day HE, Campeau S, Watson SJ Jr, Akil H. 1997. Distribution of alpha 1a-, alpha 1b- and alpha 1d-adrenergic receptor mRNA in the rat brain and spinal cord. *J Chem Neuroanat* 13: 115-139.
- De Lorenzo AJ. 1957. Electron microscope observations of the olfactory mucosa and olfactory nerve. *J Biophys Biochem Cytol* 3: 839-850.
- de Olmos J, Hardy H, Heimer L. 1978. The afferent connections of the main and the accessory olfactory bulb formations in the rat: An experimental HRP-study. *J Comp Neurol* 181: 213-244.
- De Olmos J, Heimer L. 1980. Double and triple labeling of neurons with fluorescent substances; the study of collateral pathways in the ascending raphe system. *Neurosci Lett* 19: 7-12.
- De Saint Jan D, Westbrook GL. 2005. Detecting activity in olfactory bulb glomeruli with astrocyte recording. *J Neurosci* 25: 2917-2924.
- De Souza EB, Insel TR, Perrin MH, Rivier J, Vale WW, et al. 1985. Corticotropin-releasing factor receptors are widely distributed within the rat central nervous system: An autoradiographic study. *J Neurosci* 5: 3189-3203.
- Didier A, Ottersen OP, Storm-Mathisen J. 1994. Differential subcellular distribution of glutamate and taurine in primary olfactory neurons. *Neuroreport* 6: 145-148.
- Didier A, Carleton A, Bjaalie JG, Vincent JD, Ottersen OP, et al. 2001. A dendrodendritic reciprocal synapse provides a recurrent excitatory connection in the olfactory bulb. *Proc Natl Acad Sci USA* 98: 6441-6446.
- Domyancic AV, Morilak DA. 1997. Distribution of alpha1A adrenergic receptor mRNA in the rat brain visualized by in situ hybridization. *J Comp Neurol* 386: 358-378.
- Doty RL, Rissler JM. 1989. Influence of the D-2 dopamine receptor agonist quinpirole on the odor detection performance of rats before and after spiperone administration. *Psychopharmacol* 98: 310-315.
- Doucette JR. 1984. The glial cells in the nerve fiber layer of the rat olfactory bulb. *Anat Rec* 210: 385-391.
- Doucette R. 1989. Development of the nerve fiber layer in the olfactory bulb of mouse embryos. *J Comp Neurol* 285: 514-527.
- Duchamp-Viret P, Chaput MA, Duchamp A. 1999. Odor response properties of rat olfactory receptor neurons. *Science* 284: 2171-2174.
- Duchamp-Viret P, Delaleu J-C, Duchamp A. 2000. GABA_B-mediated action in the frog olfactory bulb makes odor responses more salient. *Neuroscience* 97: 771-777.
- Duchamp-Viret P, Coronas V, Delaleu J-C, Moysse E, Duchamp A. 1997. Dopaminergic modulation of mitral cell activity in the frog olfactory bulb: A combined radioligand binding-electrophysiological study. *Neuroscience* 79: 203-216.
- Duvoisin RM, Zhang C, Ramonell K. 1995. A novel metabotropic receptor expressed in the retina and olfactory bulb. *J Neurosci* 15: 3075-3083.
- Ekstrand JJ, Domroese ME, Feig SL, Illig KR, Haberly LB. 2001a. Immunocytochemical analysis of basket cells in rat piriform cortex. *J Comp Neurol* 434: 308-328.
- Ekstrand JJ, Domroese ME, Johnson DMG, Feig SL, Knodel SM, et al. 2001b. A new subdivision of anterior piriform cortex and associated deep nucleus with novel features of interest for olfaction and epilepsy. *J Comp Neurol* 434: 289-307.
- Elaagouby A, Ravel N, Gervais R. 1991. Cholinergic modulation of excitability in the rat olfactory bulb: Effect of local application of cholinergic agents on evoked field potentials. *Neuroscience* 45: 653-662.
- El-Etri MM, Griff ER, Ennis M, Shipley MT. 1999. Evidence for cholinergic regulation of basal norepinephrine release in the rat olfactory bulb. *Neuroscience* 93: 611-617.

- Ennis M, Zimmer LA, Shipley MT. 1996. Olfactory nerve stimulation activates rat mitral cells via NMDA and non-NMDA receptors in vitro. *Neuroreport* 7: 989-992.
- Ennis M, Zhou FM, Ciombor KJ, Aroniadou-Anderjaska V, Hayar A, et al. 2001. Dopamine D2 receptor-mediated presynaptic inhibition of olfactory nerve terminals. *J Neurophysiol* 86: 2986-2997.
- Ennis M, Zhu M, Heinbockel T, Hayar A. 2006. Olfactory nerve-evoked metabotropic glutamate receptor-mediated responses in olfactory bulb mitral cells. *J Neurophysiol* 95: 2233-2241.
- Erlich ME, Grillo M, Joh TH, Margolis FL, Baker H. 1990. Transneuronal regulation of neuronal specific gene expression in the mouse olfactory bulb. *Mol Brain Res* 7: 115-122.
- Ezeh PI, Wellis DP, Scott JW. 1993. Organization of inhibition in the rat olfactory bulb external plexiform layer. *J Neurophysiol* 70: 263-274.
- Fadool DA, Levitan IB. 1998. Modulation of olfactory bulb potassium current by tyrosine phosphorylation. *J Neurosci* 18: 6126-6137.
- Fadool DA, Tucker K, Phillips JJ, Simmen JA. 2000. Brain insulin receptor causes activity-dependent current suppression in olfactory bulb through multiple phosphorylation of Kv1.3. *J Neurophysiol* 83: 2332-2348.
- Fallon JH, Leslie FM. 1986. Distribution of dynorphin and enkephalin peptides in the rat brain. *J Comp Neurol* 249: 293-336.
- Fallon JH, Loughlin SE. 1982. Monoamine innervation of the forebrain: Collateralization. *Brain Res Bull* 9: 295-307.
- Fallon JH, Moore RY. 1978. Catecholamine innervation of the basal forebrain. IV. Topography of the dopamine projection to the basal forebrain and neostriatum. *J Comp Neurol* 180: 545-580.
- Ferraris N, Perroteau I, De Marchis S, Fasolo A, Bovolenta P. 1997. Glutamatergic deafferentation of olfactory bulb modulates the expression of mGluR1a mRNA. *Neuroreport* 8: 1949-1953.
- Ferriero D, Margolis FL. 1975. Denervation in the primary olfactory pathway of mice. II. Effects on carnosine and other amine compounds. *Brain Res* 94: 75-86.
- Ffrench-Mullen JMH, Hori N, Carpenter DO. 1986. Receptors for excitatory amino acids on neurons in rat pyriform cortex. *J Neurophysiol* 55: 1283-1294.
- Ffrench-Mullen JM, Koller K, Zaczek R, Coyle JT, Hori N, et al. 1985. N-Acetylaspartylglutamate: Possible role as the neurotransmitter of the lateral olfactory tract. *Proc Natl Acad Sci USA* 82: 3897-3900.
- Firestein S. 2001. How the olfactory system makes sense of scents. *Nature* 413: 211-218.
- Fiske BK, Brunjes PC. 2001. Cell death in the developing and sensory-deprived rat olfactory bulb. *J Comp Neurol* 431: 311-319.
- Fonseca MI, Lunt GG, Aguilar JS. 1991b. Inhibition of muscarinic cholinergic receptors by disulfide reducing agents and arsenicals. Differential effect on locust and rat. *Biochem Pharmacol* 41: 735-742.
- Fonseca MI, Aguilar JS, Skorupa AF, Klein WL. 1991a. Cellular mapping of m2 muscarinic receptors in the rat olfactory bulb using antiserum raised against a cytoplasmic loop peptide. *Brain Res* 563: 163-170.
- Frazier LL, Brunjes PC. 1988. Unilateral odor deprivation: Early postnatal changes in olfactory bulb cell density and number. *J Comp Neurol* 269: 355-370.
- Frazier-Cierpial L, Brunjes PC. 1989. Early postnatal cellular proliferation and survival in the olfactory bulb and rostral migratory stream of normal and unilaterally odor-deprived rats. *J Comp Neurol* 289: 481-492.
- Fremeau RT Jr, Duncan GE, Fornaretto MG, Dearry A, Gingrich JA, et al. 1991. Localization of D1 dopamine receptor mRNA in brain supports a role in cognitive, affective, and neuroendocrine aspects of dopaminergic neurotransmission. *Proc Natl Acad Sci USA* 88: 3772-3776.
- Friedman D, Strowbridge BW. 2000. Functional role of NMDA autoreceptors in olfactory mitral cells. *J Neurophysiol* 84: 39-50.
- Friedman D, Strowbridge BW. 2003. Both electrical and chemical synapses mediate fast network oscillations in the olfactory bulb. *J Neurophysiol* 89: 2601-2610.
- Friedrich RW, Korsching SI. 1997. Combinatorial and chemotopic odorant coding in the zebrafish olfactory bulb visualized by optical imaging. *Neuron* 18: 737-752.
- Frings S, Seifert R, Godde M, Kaupp UB. 1995. Profoundly different calcium permeation and blockage determine the specific function of distinct cyclic nucleotide-gated channels. *Neuron* 15: 169-179.
- Fritschy JM, Benke D, Mertens S, Oertel WH, Bachi T, et al. 1992. Five subtypes of type A gamma-aminobutyric acid receptors identified in neurons by double and triple immunofluorescence staining with subunit-specific antibodies. *Proc Natl Acad Sci USA* 89: 6726-6730.
- Frosch MP, Dichter MA. 1982. Physiology and pharmacology of olfactory bulb neurons in dissociated cell culture. *Brain Res* 290: 321-332.
- Fuller TA, Price JL. 1988. Putative glutamatergic and/or aspartatergic cells in the main and accessory olfactory bulbs of the rat. *J Comp Neurol* 276: 209-218.
- Gall C, Seroogy KB, Brecha N. 1986. Distribution of VIP- and NPY-like immunoreactivities in rat main olfactory bulb. *Brain Res* 374: 389-394.
- Gall C, Sumikawa K, Lynch G. 1990. Regional distribution of mRNA for a putative kainate receptor in rat brain. *Eur J Pharmacol* 189: 217-221.
- Gall CM, Hendry SHC, Seroogy KB, Jones EG, Haycock JW. 1987. Evidence for coexistence of GABA and dopamine

- in neurons of the rat olfactory bulb. *J Comp Neurol* 266: 307-318.
- Geiling H, Schild D. 1996. Glutamate-mediated release of Ca^{2+} in mitral cells of the olfactory bulb. *J Neurophysiol* 76: 563-570.
- Gellman RL, Aghajanian GK. 1993. Pyramidal cells in piriform cortex receive a convergence of inputs from monoamine-activated GABAergic interneurons. *Brain Res* 600: 63-73.
- Gellman RL, Aghajanian GK. 1994. Serotonin₂ receptor-mediated excitation of interneurons in piriform cortex: Antagonism by atypical antipsychotic drugs. *Neuroscience* 58: 515-525.
- Getchell ML, Getchell TV. 1992. Fine structural aspects of secretion and extrinsic innervation in the olfactory mucosa. *Microsc Res Tech* 23:111-127.
- Getchell TV, Shepherd GM. 1975. Synaptic actions on mitral and tufted cells elicited by olfactory nerve volleys in the rabbit. *J Physiol* 251: 497-522.
- Getchell TV, Margolis FL, Getchell ML. 1984. Perireceptor and receptor events in vertebrate olfaction. *Prog Neurobiol* 23: 317-345.
- Gheusi G, Cremer H, McLean H, Chazal G, Vincent J-D, et al. 2000. Importance of newly generated neurons in the adult olfactory bulb for odor discrimination. *Proc Natl Acad Sci USA* 97: 1823-1828.
- Giustetto M, Kirsch J, Fritschy JM, Cantino D, Sassoè-Pognetto M. 1998. Localization of the clustering protein gephyrin at GABAergic synapses in the main olfactory bulb of the cat. *J Comp Neurol* 395: 231-244.
- Giustetto M, Bovolín P, Fasolo A, Bonino M, Cantino D, et al. 1997. Glutamate receptors in the olfactory bulb synaptic circuitry: Heterogeneity and synaptic localization of *N*-methyl-D-aspartate receptor subunit 1 and AMPA receptor subunit 1. *Neuroscience* 76: 787-798.
- Golgi C. 1875. Sulla Fina Structura del Bulbi Olfattori. *Rev Sper Feniatr* 1: 405-425.
- Gracia-Llanes FJ, Crespo C, Blasco-Ibáñez JM, Marqués-Mari AI, Martínez-Guijarro FJ. 2003. VIP-containing deep short-axon cells of the olfactory bulb innervate interneurons different from granule cells. *Eur J Neurosci* 18: 1751-1763.
- Graziadei PP, Graziadei GA. 1979. Neurogenesis and neuron regeneration in the olfactory system of mammals. I. Morphological aspects of differentiation and structural organization of the olfactory sensory neurons. *J Neurocytol* 8: 1-18.
- Greer CA. 1987. Golgi analyses of dendritic organization among denervated olfactory bulb granule cells. *J Comp Neurol* 256: 442-452.
- Greer CA, Stewart WB, Teicher MH, Shepherd GM. 1982. Functional development of the olfactory bulb and a unique glomerular complex in the neonatal rat. *J Neurosci* 2: 1744-1759.
- Guthrie KM, Gall CM. 1995a. Functional mapping of odor-activated neurons in the olfactory bulb. *Chem Senses* 20: 271-282.
- Guthrie KM, Gall CM. 1995b. Odors increase fos in olfactory bulb neurons including dopaminergic cells. *Neuroreport* 6: 2145-2149.
- Guthrie KM, Pullara JM, Marshall JF, Leon M. 1991. Olfactory deprivation increases dopamine D2 receptor density in the rat olfactory bulb. *Synapse* 8: 61-70.
- Haberly L. 1983. Structure of the piriform cortex of the opossum. I. Description of neuron types with Golgi methods. *J Comp Neurol* 213: 163-187.
- Haberly LB, Bower JM. 1984. Analysis of association fiber system in piriform cortex with intracellular recording and staining techniques. *J Neurophysiol* 51: 90-112.
- Haberly LB, Presto S. 1986. Ultrastructural analysis of synaptic relationships of intracellularly stained pyramidal cell axons in piriform cortex. *J Comp Neurol* 248: 464-474.
- Haberly LB, Price JL. 1978a. Association and commissural fiber systems of the olfactory cortex of the rat. I. Systems originating in the piriform cortex and adjacent areas. *J Comp Neurol* 178: 711-740.
- Haberly LB, Price JL. 1978b. Association and commissural fiber systems of the olfactory cortex of the rat. II. Systems originating in the olfactory peduncle. *J Comp Neurol* 181: 781-807.
- Haberly LB, Shepherd GM. 1973. Current-density analysis of summed evoked potentials in opossum prepiriform cortex. *J Neurophysiol* 36: 789-802.
- Haberly LB, Hansen DJ, Feig SL, Presto S. 1987. Distribution and ultrastructure of neurons in opossum piriform cortex displaying immunoreactivity to GABA and GAD and high-affinity tritiated GABA uptake. *J Comp Neurol* 266: 269-290.
- Hack MA, Saghatelian A, de Chevigny A, Pfeifer A, Ashery-Padan R, et al. 2005. Neuronal fate determinants of adult olfactory bulb neurogenesis. *Nat Neurosci* 8: 865-872.
- Halabisky B, Strowbridge BW. 2003. γ -frequency excitatory input to granule cells facilitates dendrodendritic inhibition in the rat olfactory bulb. *J Neurophysiol* 90: 644-654.
- Halabisky B, Friedman D, Radojicic M, Strowbridge BW. 2000. Calcium influx through NMDA receptors directly evokes GABA release in olfactory bulb granule cells. *J Neurosci* 20: 5124-5134.
- Hadley WM, Dahl AR. 1982. Cytochrome P-450 dependent monooxygenase activity in rat nasal epithelial membranes. *Toxicol Lett* 10: 417-422.
- Halász N. 1987. Autoradiographic localization of high affinity uptake sites for 3H-D-aspartate in the rat olfactory bulb. *Acta Biol Hung* 38: 383-392.

- Halász N. 1990. The vertebrate olfactory system. Chemical neuroanatomy, function and development. Akadémiai Kiadó, Budapest, 281 pp.
- Halász N, Shepherd GM. 1983. Neurochemistry of the vertebrate olfactory bulb. *Neuroscience* 10: 579-619.
- Halász N, Ljungdahl A, Hökfelt T. 1978. Transmitter histochemistry of the rat olfactory bulb. II. Fluorescence histochemical, autoradiographic and electron microscopic localization of monoamines. *Brain Res* 154: 253-271.
- Halász N, Hokfelt T, Norman AW, Goldstein M. 1985. Tyrosine hydroxylase and 28K-vitamin D-dependent calcium binding protein are localized in different subpopulations of periglomerular cells of the rat olfactory bulb. *Neurosci Lett* 61: 103-107.
- Halász N, Johansson O, Hökfelt T, Ljungdahl A, Goldstein M. 1981. Immunohistochemical identification of two types of dopamine neuron in the rat olfactory bulb as seen by serial sectioning. *J Neurocytol* 10: 251-259.
- Halász N, Ljungdahl A, Hokfelt T, Johansson O, Goldstein M, et al. 1977. Transmitter histochemistry of the rat olfactory bulb. I. Immunohistochemical localization of monoamine synthesizing enzymes. Support for intrabulbar, peri-glomerular dopamine neurons. *Brain Res* 126: 455-474.
- Hamada S, Senzaki K, Hamaguchi-Hamada K, Tabuchi K, Yamamoto H, et al. 1998. Localization of 5-HT_{2A} receptor in rat cerebral cortex and olfactory system revealed by immunohistochemistry using two antibodies raised in rabbit and chicken. *Brain Res Mol Brain Res* 54: 199-211.
- Hamana H, Hirono J, Kizumi M, Sato T. 2003. Sensitivity-dependent hierarchical receptor codes for odors. *Chem Senses* 28: 87-104.
- Hamilton KA, Coppola DM. 2003. Distribution of GluR1 is altered in the olfactory bulb following neonatal naris occlusion. *J Neurobiol* 54: 326-336.
- Hamilton KA, Heinbockel T, Ennis M, Szabó G, Erdélyi F, et al. 2005. Properties of external plexiform layer interneurons in mouse olfactory bulb slices. *Neuroscience* 133: 819-829.
- Hara Y, Shiosaka S, Senba E, Sakanaka M, Inagaki S, et al. 1982. Ontogeny of the neurotensin-containing neuron system of the rat: Immunohistochemical analysis. I. Forebrain and diencephalon. *J Comp Neurol* 208: 177-195.
- Hardy A, Palouzier-Paulignan B, Duchamp A, Royet JB, Duchamp-Viret P. 2005. 5-hydroxytryptamine action in the rat olfactory bulb: In vitro electrophysiological patch-clamp recordings of juxtaglomerular and mitral cells. *Neuroscience* 131: 717-731.
- Harlan RE, Shivers BD, Romano GJ, Howells RD, Pfaff DW. 1987. Localization of preproenkephalin mRNA in the rat brain and spinal cord by in situ hybridization. *J Comp Neurol* 258: 159-184.
- Hasselmo ME, Bower JM. 1991. Selective suppression of afferent but not intrinsic fiber synaptic transmission by 2-amino-4-phosphonobutyric acid (AP4) in piriform cortex. *Brain Res* 548: 248-255.
- Hasselmo ME, Bower JM. 1992. Cholinergic suppression specific to intrinsic not afferent fiber synapses in rat piriform (olfactory) cortex. *J Neurophysiol* 67: 1222-1229.
- Hasselmo ME, Anderson BP, Bower JM. 1992. Cholinergic modulation of cortical associative memory function. *J Neurophysiol* 67: 1230-1246.
- Hasselmo ME, Linster C, Ma D, Cekic M. 1997. Noradrenergic suppression of synaptic transmission may influence cortical signal-to-noise ratio. *J Neurophysiol* 77: 3326-3339.
- Hayar A, Shipley MT, Ennis M. 2005. Olfactory bulb external tufted cells are synchronized by multiple intraglomerular mechanisms. *J Neurosci* 25: 8197-8208.
- Hayar A, Karnup S, Ennis M, Shipley MT. 2004b. External tufted cells: A major excitatory element that coordinates glomerular activity. *J Neurosci* 24: 6676-6685.
- Hayar A, Karnup S, Shipley MT, Ennis M. 2004a. Olfactory bulb glomeruli: External tufted cells intrinsically burst at theta frequency and are entrained by patterned olfactory input. *J Neurosci* 24: 1190-1199.
- Hayar A, Heyward PM, Heinbockel T, Shipley MT, Ennis M. 2001. Direct excitation of mitral cells via activation of alpha1-noradrenergic receptors in rat olfactory bulb slices. *J Neurophysiol* 86: 2173-2182.
- Heinbockel T, Ennis M. 2003. Granule cell-mediated lateral inhibition can be driven by metabotropic glutamate receptors. *AchemS Abstr XXV*: 201.
- Heinbockel T, Heyward P, Conquet F, Ennis M. 2004. Regulation of main olfactory bulb mitral cell excitability by metabotropic glutamate receptor mGluR1. *J Neurophysiol* 92: 3085-3096.
- Hill JAJ, Zoli M, Bourgeois J, Changeux J. 1993. Immunocytochemical localization of a neuronal nicotinic receptor: The b2-subunit. *J Neurosci* 13: 1551-1568.
- Hinds JW. 1968. Autoradiographic study of histogenesis in the mouse olfactory bulb. I. Time of origin of neurons and neuroglia. *J Comp Neurol* 134: 287-304.
- Hirata Y. 1964. Some observations on the fine structure of the synapses in the olfactory bulb of the mouse, with particular reference to the atypical synaptic configurations. *Arch Histol Jap* 24: 293-302.
- Hoffman BJ, Mezey E. 1989. Distribution of serotonin 5-HT_{1C} receptor mRNA in adult rat brain. *FEBS Lett* 247: 453-462.
- Hori N, Auker CR, Braitman DJ, Carpenter DO. 1982. Pharmacologic sensitivity of amino acid responses and synaptic activation of in vitro prepyriform neurons. *J Neurophysiol* 1289-1301.
- Hormuzdi SG, Pais I, Le Bleu FEN, Towers SK, Rozov A, et al. 2001. Impaired electrical signaling disrupts gamma

- frequency oscillations in connexin 36-deficient mice. *Neuron* 31: 487-495.
- Horning MS, Blakemore LJ, Trombley PQ. 2000. Endogenous mechanisms of neuroprotection: Role of zinc, copper, and carnosine. *Brain Res* 852: 56-61.
- Horning MS, Kwon B, Blakemore LJ, Spencer CM, Goltz M, et al. 2004. Alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionate receptor subunit expression in rat olfactory bulb. *Neurosci Lett* 372: 230-234.
- Hsia AY, Vincent JD, Lledo PM. 1999. Dopamine depresses synaptic inputs into the olfactory bulb. *J Neurophysiol* 82: 1082-1085.
- Huang Q, Zhou D, Chase K, Gusella JF, Aronin N, et al. 1992. Immunohistochemical localization of the D1 dopamine receptor in rat brain reveals its axonal transport, pre- and postsynaptic localization, and prevalence in the basal ganglia, limbic system, and thalamic reticular nucleus. *Proc Natl Acad Sci USA* 89: 11988-11992.
- Hunt S, Schmidt J. 1978. Some observations on the binding patterns of alpha-bungarotoxin in the central nervous system of the rat. *Brain Res* 157: 213-232.
- Illig KR, Haberly LB. 2003. Odor-evoked activity is spatially distributed in piriform cortex. *J Comp Neurol* 457: 361-373.
- Imaki T, Nahon JL, Sawchenko PE, Vale W. 1989. Widespread expression of corticotropin-releasing factor messenger RNA and immunoreactivity in the rat olfactory bulb. *Brain Res* 496: 35-44.
- Inagaki S, Sakanaka M, Shiosaka S, Senba E, Takatsuki K, et al. 1982. Ontogeny of substance P-containing neuron system of the rat: Immunohistochemical analysis—I. Forebrain and upper brain stem. *Neuroscience* 7: 251-277.
- Isaacson JS. 1999. Glutamate spillover mediates excitatory transmission in the rat olfactory bulb. *Neuron* 23: 377-384.
- Isaacson JS, Murphy GJ. 2001. Glutamate-mediated extrasynaptic inhibition: Direct coupling of NMDA receptors to Ca^{2+} -activated K^+ channels. *Neuron* 31: 1027-1034.
- Isaacson JS, Strowbridge BW. 1998. Olfactory reciprocal synapses: Dendritic signaling in the CNS. *Neuron* 20: 749-761.
- Isaacson JS, Vitten H. 2003. GABAB receptors inhibit dendrodendritic transmission in the rat olfactory bulb. *J Neurosci* 23: 2032-2039.
- Ivic L, Pyrski MM, Margolis JW, Richards LJ, Firestein S, et al. 2000. Adenoviral vector-mediated rescue of the OMP-null phenotype in vivo. *Nat Neurosci* 3: 1113-1120.
- Jackowski A, Parnavelas JG, Lieberman AR. 1978. The reciprocal synapse in the external plexiform layer of the mammalian olfactory bulb. *Brain Res* 159: 17-28.
- Jacobowitz DM, Winsky L. 1991. Immunocytochemical localization of calretinin in the forebrain of the rat. *J Comp Neurol* 304: 198-218.
- Jahr CE, Nicoll RA. 1982. Noradrenergic modulation of dendrodendritic inhibition in the olfactory bulb. *Nature* 297: 227-229.
- Jardemark K, Nilsson M, Muyderman H, Jacobson I. 1997. Ca^{2+} ion permeability properties of (RS) α -amino-3-hydroxy-5-methyl-4-isoxazolepropionate (AMPA) receptors in isolated interneurons from the olfactory bulb of the rat. *J Neurophysiol* 77: 702-708.
- Jin BK, Franzen L, Baker H. 1996. Regulation of c-Fos mRNA and fos protein expression in olfactory bulb from unilaterally odor-deprived mice. *Int J Dev Neurosci* 14: 971-982.
- Jia C, Halpern M. 2004. Calbindin D28k, parvalbumin, and calretinin immunoreactivity in the main and accessory olfactory bulbs of the gray short-tailed opossum, *Monodelphis domestica*. *J Morphol* 259: 271-280.
- Jiang MR, Griff ER, Ennis M, Zimmer LA, Shipley MT. 1996. Activation of locus coeruleus enhances the responses of olfactory bulb mitral cells to weak olfactory nerve input. *J Neurosci* 16: 6319-6329.
- Johnson BA, Leon M. 2000a. Modular representations of odorants in the glomerular layer of the rat olfactory bulb and the effects of stimulus concentration. *J Comp Neurol* 422: 496-509.
- Johnson BA, Leon M. 2000b. Odorant molecular length: One aspect of the olfactory code. *J Comp Neurol* 426: 330-338.
- Johnson BA, Woo CC, Leon M. 1998. Spatial coding of odorant features in the glomerular layer of the rat olfactory bulb. *J Comp Neurol* 393: 457-471.
- Johnson DM, Illig KR, Behan M, Haberly LB. 2000. New features of connectivity in piriform cortex visualized by intracellular injection of pyramidal cells suggest that "primary" olfactory cortex functions like "association" cortex in other sensory systems. *J Neurosci* 20: 6974-6982.
- Jourdan F, Duveau A, Astic L, Holley A. 1980. Spatial distribution of [^{14}C]2-deoxyglucose uptake in the olfactory bulbs of rats stimulated with two different odours. *Brain Res* 188: 139-154.
- Kaba H, Keverne EB. 1988. The effect of microinfusions of drugs into the accessory olfactory bulb on the olfactory block to pregnancy. *Neuroscience* 25: 1007-1011.
- Kaba H, Rosser A, Keverne B. 1989. Neural basis of olfactory memory in the context of pregnancy block. *Neuroscience* 32: 657-662.
- Kakuta S, Oda S, Takayanagi M, Kishi K. 1998. Parvalbumin immunoreactive neurons in the main olfactory bulb of the house mouse shrew, *Suncus murinus*. *Brain Behav Evol* 52: 285-291.
- Kamisaki Y, Wada K, Nakamoto K, Itoh T. 1996. Effects of taurine on GABA release from synaptosomes of rat olfactory bulb. *Amino Acids* 10: 49-57.

- Kanter ED, Kapur A, Haberly LB. 1996. A dendritic GABA-mediated IPSP regulates facilitation of NMDA-mediated responses to burst stimulation of afferent fibers in piriform cortex. *J Neurosci* 16: 307-312.
- Kaplan MS, McNelly NA, Hinds JW. 1985. Population dynamics of adult-formed granule neurons of the rat olfactory bulb. *J Comp Neurol* 239: 117-125.
- Kapur A, Pearce RA, Lytton WW, Haberly LB. 1997. GABA-mediated IPSCs in piriform cortex have fast and slow components with different properties and locations on pyramidal cells. *J Neurophysiol* 78: 2531-2545.
- Kasowski HJ, Kim H, Greer CA. 1999. Compartmental organization of the olfactory bulb glomerulus. *J Comp Neurol* 407: 261-274.
- Kato T, Yokouchi K, Kawagishi K, Fukushima N, Miwa T, et al. 2000. Fate of newly formed periglomerular cells in the olfactory bulb. *Acta Otolaryngol* 120: 876-879.
- Kawano T, Margolis FL. 1982. Transsynaptic regulation of olfactory bulb catecholamines in mice and rats. *J Neurochem* 39: 342-348.
- Keller A, Margolis FL. 1975. Immunological studies of the rat olfactory marker protein. *J Neurochem* 24: 1101-1106.
- Keller A, Yagodin S, Aroniadou-Anderjaska V, Zimmer LA, Ennis M, et al. 1998. Functional organization of rat olfactory bulb glomeruli revealed by optical imaging. *J Neurosci* 18: 2602-2612.
- Kendrick KM, Levy F, Keverne EB. 1992. Changes in the sensory processing of olfactory signals induced by birth in sheep. *Science* 256: 833-836.
- Kerr DIB, Hagbarth KE. 1955. An investigation of the olfactory centrifugal system. *J Neurophysiol* 18: 362-374.
- Kinoshita A, Shigemoto R, Ohishi H, van der Putten H, Mizuno N. 1998. Immunohistochemical localization of metabotropic glutamate receptors, mGluR7a and mGluR7b, in the central nervous system of the adult rat and mouse: A light and electron microscopic study. *J Comp Neurol* 393: 332-352.
- Kinoshita A, Ohishi H, Neki A, Nomura S, Shigemoto R, et al. 1996. Presynaptic localization of metabotropic glutamate receptor, mGluR8, in the rhinencephalic areas: A light and electron microscope study in the rat. *Neurosci Lett* 207: 61-64.
- Kinzie JM, Saugstad JA, Westbrook GL, Segerson TP. 1995. Distribution of metabotropic glutamate receptor 7 messenger RNA in the developing and adult rat brain. *Neuroscience* 69: 167-176.
- Kinzie JM, Shinohara MM, Pol van den AN, Westbrook GL, Segerson TP. 1997. Immunolocalization of metabotropic glutamate receptor 7 in the rat olfactory bulb. *J Comp Neurol* 385: 372-384.
- Kirschenbaum B, Doetsch F, Lois C, Alvarez-Buylla A. 1999. Adult subventricular zone neuronal precursors continue to proliferate and migrate in the absence of the olfactory bulb. *J Neurosci* 19: 2171-2180.
- Kiyoshi K, Mori K, Ojima H. 1984. Distribution of local axon collaterals of mitral, displaced mitral, and tufted cells in the rabbit olfactory bulb. *J Comp Neurol* 225: 511-525.
- Kiyoshi K, Mori K, Tazawa Y. 1982. Three-dimensional analysis of dendritic trees of mitral cells in the rabbit olfactory bulb. *Neurosci Lett* 28: 127-132.
- Kleene SJ, Gesteland RC. 1991. Calcium-activated chloride conductance in frog olfactory cilia. *J Neurosci* 11: 3624-3629.
- Kosaka T, Kosaka K. 2003. Neuronal gap junctions in the rat main olfactory bulb, with special reference to intraglomerular gap junction. *Neurosci Res* 45: 189-209.
- Kosaka T, Kosaka K. 2004. Neuronal gap junctions between intraglomerular mitral/tufted cell dendrites in the mouse main olfactory bulb. *Neurosci Res* 49: 373-378.
- Kosaka T, Kosaka K. 2005. Intraglomerular dendritic link connected by gap junctions and chemical synapses in the mouse main olfactory bulb: Electron microscopic serial section analyses. *Neuroscience* 131: 611-625.
- Kosaka K, Heizmann CW, Kosaka T. 1994a. Calcium-binding protein parvalbumin-immunoreactive neurons in the rat olfactory bulb. 1. Distribution and structural features in adult rat. *Exp Brain Res* 99: 191-204.
- Kosaka K, Heizmann CW, Kosaka T. 1994b. Calcium-binding protein parvalbumin-immunoreactive neurons in the rat olfactory bulb. 2. Postnatal development. *Exp Brain Res* 99: 205-213.
- Kosaka K, Aika Y, Toida K, Kosaka T. 2001. Structure of intraglomerular dendritic tufts of mitral cells and their contacts with olfactory nerve terminals and calbindin-immunoreactive type 2 periglomerular neurons. *J Comp Neurol* 440: 219-235.
- Kosaka T, Deans MR, Paul DL, Kosaka K. 2005. Neuronal gap junctions in the mouse main olfactory bulb: Morphological analyses on transgenic mice. *Neuroscience* 134: 757-769.
- Kosaka K, Toida K, Margolis FL, Kosaka T. 1997. Chemically defined neuron groups and their subpopulations in the glomerular layer of the rat main olfactory bulb—II. Prominent differences in the intraglomerular dendritic arborization and their relationship to olfactory nerve terminals. *Neuroscience* 76: 775-786.
- Kosaka K, Toida K, Aika Y, Kosaka T. 1998. How simple is the organization of the olfactory glomerulus?: The heterogeneity of so-called periglomerular cells. *Neurosci Res* 30: 101-110.
- Kosaka K, Hama K, Nagatsu I, Wu JY, Kosaka T. 1988. Possible coexistence of amino acid (gamma-aminobutyric acid), amine (dopamine) and peptide (substance P); neurons containing immunoreactivities for glutamic acid decarboxylase,

- tyrosine hydroxylase and substance P in the hamster main olfactory bulb. *Exp Brain Res* 71: 633-642.
- Kosaka T, Hataguchi Y, Hama K, Nagatsu I, Wu JY. 1985. Coexistence of immunoreactivities for glutamate decarboxylase and tyrosine hydroxylase in some neurons in the periglomerular region of the rat main olfactory bulb: Possible coexistence of gamma-aminobutyric acid (GABA) and dopamine. *Brain Res* 343: 166-171.
- Kosaka T, Kosaka K, Hama K, Wu J-Y, Nagatsy I. 1987b. Differential effect of functional olfactory deprivation on the GABAergic and catecholaminergic traits in the rat main olfactory bulb. *Brain Res* 413: 197-203.
- Kosaka K, Aika Y, Toida K, Heizmann CW, Hunziker W, et al. 1995. Chemically defined neuron groups and their subpopulations in the glomerular layer of the rat main olfactory bulb. *Neurosci Res* 23: 73-88.
- Kosaka K, Hama K, Nagatsu I, Wu JY, Ottersen OP, et al. 1987a. Postnatal development of neurons containing both catecholaminergic and GABAergic traits in the rat main olfactory bulb. *Brain Res* 403: 355-360.
- Kosaka T, Kosaka K, Hataguchi Y, Nagatsu I, Wu JY, et al. 1987c. Catecholaminergic neurons containing GABA-like and/or glutamic acid decarboxylase-like immunoreactivities in various brain regions of the rat. *Exp Brain Res* 66: 191-210.
- Kosaka T, Kosaka K, Heizmann CW, Nagatsu I, Wu JY, et al. 1987d. An aspect of the organization of the GABAergic system in the rat main olfactory bulb: Laminar distribution of immunohistochemically defined subpopulations of GABAergic neurons. *Brain Res* 411: 373-378.
- Koster NL, Norman AB, Richtand NM, Nickell WT, Puche AC, et al. 1999. Olfactory receptor neurons express D2 dopamine receptors. *J Comp Neurol* 411: 666-673.
- Kratskin I, Belluzzi O. 2002. Immunohistochemical localization of the taurine-synthesizing enzyme in the rat olfactory bulb. *Chem Senses* 27: 46.
- Kratskin IL, Rio JP, Kenigfest NB, Doty RL, Repérant J. 2000. A light and electron micro scopic study of taurine-like immunoreactivity in the main olfactory bulb of frogs. *J Chem Neuroanat* 18: 87-101.
- Krautwurst D, Yau KW, Reed RR. 1998. Identification of ligands for olfactory receptors by functional expression of a receptor library. *Cell* 95: 917-926.
- Kream RM, Davis BJ, Kawano T, Margolis FL, Macrides F. 1984. Substance P and catecholaminergic expression in neurons of the hamster main olfactory bulb. *J Comp Neurol* 222: 140-154.
- Kubota Y, Jones EG. 1993. Co-localization of two calcium binding proteins in GABA cells of rat piriform cortex. *Brain Res* 600: 339-344.
- Kunze WAA, Shafton AD, Kemm RE, McKenzie JS. 1991. Effect of stimulating the nucleus of the horizontal limb of the diagonal band on single unit activity in the olfactory bulb. *Neuroscience* 40: 21-27.
- Kunze WAA, Shafton AD, Kemm RE, McKenzie JS. 1992. Intracellular responses of olfactory bulb granule cells to stimulating the horizontal diagonal band nucleus. *Neuroscience* 48: 363-369.
- Laaris N, Ennis M. 2002. Distinct activity patterns evoked by activation of mitral/tufted cell and centrifugal fiber inputs to main olfactory bulb (MOB) granule cells. *Soc Neurosci Abstr* 561: 14.
- Landis DMD, Reese TS, Raviola E. 1974. Differences in membrane structure between excitatory and inhibitory components of the reciprocal synapse in the olfactory bulb. *J Comp Neurol* 155: 67-92.
- Laurie DJ, Seeburg PH, Wisden W. 1992. The distribution of 13 GABAA receptor subunit mRNAs in the rat brain. II. Olfactory bulb and cerebellum. *J Neurosci* 12: 1063-1076.
- Le Jeune H, Jourdan F. 1994. Acetylcholinesterase-containing intrinsic neurons in the rat main olfactory bulb: Cytological and neurochemical features. *Eur J Neurosci* 6: 1432-1444.
- Le Jeune H, Aubert I, Jourdan F, Quirion R. 1996. Developmental profiles of various cholinergic markers in the rat main olfactory bulb using quantitative autoradiography. *J Comp Neurol* 373: 433-450.
- Leinders-Zufall T, Ma M, Zufall F. 1999. Impaired odor adaptation in olfactory receptor neurons after inhibition of Ca^{2+} /calmodulin kinase II. *J Neurosci* 19: 1-6.
- Leinders-Zufall T, Greer CA, Shepherd GM, Zufall F. 1998. Imaging odor-induced calcium transients in single olfactory cilia: Specificity of activation and role in transduction. *J Neurosci* 18: 5630-5639.
- Leinders-Zufall T, Rand MN, Shepherd GM, Greer CA, Zufall F. 1997. Calcium entry through cyclic nucleotide-gated channels in individual cilia of olfactory receptor cells: Spatiotemporal dynamics. *J Neurosci* 17: 4136-4148.
- Levey AI, Kitt CA, Simonds WF, Price DL, Brann MR. 1991. Identification and localization of muscarinic acetylcholine receptor proteins in brain with subtype-specific antibodies. *J Neurosci* 11: 3218-3226.
- Levey AI, Hersch SM, Rye DB, Sunahara RK, Niznik HB, et al. 1993. Localization of D1 and D2 dopamine receptors in brain with subtype-specific antibodies. *Proc Natl Acad Sci USA* 90: 8861-8865.
- Libri V, Constanti A, Calaminci M, Nistico G. 1994. A comparison of the muscarinic response and morphological properties of identified cells in the guinea-pig olfactory cortex in vitro. *Neuroscience* 59: 331-347.
- Linster C, Wyble BP, Hasselmo ME. 1999. Electrical stimulation of the horizontal limb of the diagonal band of Broca modulates population EPSPs in piriform cortex. *J Neurophysiol* 8: 2737-2742.

- Liu N. 2000. Regional distribution of protein kinases in normal and odor-deprived mouse olfactory bulbs. *Chem Sens* 25: 401-406.
- Liu G, Rao Y. 2003. Neuronal migration from the forebrain to the olfactory bulb requires a new attractant persistent in the olfactory bulb. *J Neurosci* 23: 6651-6659.
- Liu WL, Shipley MT. 1994. Intrabulbar associational system in the rat olfactory bulb comprises cholecystokinin-containing tufted cells that synapse onto the dendrites of GABAergic granule cells. *J Comp Neurol* 346: 541-558.
- Liu CJ, Grandes P, Matute C, Cuenod M, Streit P. 1989. Glutamate-like immunoreactivity revealed in rat olfactory bulb, hippocampus and cerebellum by monoclonal antibody and sensitive staining method. *Histochemistry* 90: 427-445.
- Liu N, Cigola E, Tinti C, Jin BK, Conti B, et al. 1999. Unique regulation of immediate early gene and tyrosine hydroxylase expression in the odor-deprived mouse olfactory bulb. *J Biol Chem* 274: 3042-3047.
- Lodovichi C, Belluscio L, Katz LC. 2003. Functional topography of connections linking mirror-symmetric maps in the mouse olfactory bulb. *Neuron* 38: 265-276.
- Lois C, Alvarez-Buylla A. 1994. Long-distance neuronal migration in the adult mammalian brain. *Science* 264: 1145-1148.
- Loo YT. 1931. The forebrain of the opossum, *Dedelphys virginiana*. II. Histology. *J Comp Neurol* 52: 1-48.
- López-Mascaraque L, de Carlos JA, Valverde F. 1990. Structure of the olfactory bulb of the hedgehog (*Erinaceus europaeus*): A Golgi study of the intrinsic organization of the superficial layers. *J Comp Neurol* 301: 243-261.
- López-Mascaraque L, Villalba RM, de Carlos JA. 1989. Vasoactive intestinal polypeptide-immunoreactive neurons in the main olfactory bulb of the hedgehog (*Erinaceus europaeus*). *Neurosci Lett* 98: 19-21.
- Lowe G. 2003. Flash photolysis reveals a diversity of ionotropic glutamate receptors on the mitral cell somatodendritic membrane. *J Neurophysiol* 90: 1737-1746.
- Lowe G, Gold GH. 1993. Contribution of the ciliary cyclic nucleotide-gated conductance to olfactory transduction in the salamander. *J Physiol* 462: 175-96.
- Luo M, Katz L. 2001. Response correlation maps of neurons in the mammalian olfactory bulb. *Neuron* 32: 1165-1179.
- Luskin MB. 1993. Restricted proliferation and migration of postnatally generated neurons derived from the forebrain subventricular zone. *Neuron* 11: 173-189.
- Luskin MB, Price JL. 1983. The topographic organization of associational fibers of the olfactory system in the rat, including centrifugal fibers to the olfactory bulb. *J Comp Neurol* 216: 264-291.
- Lysakowski A, Wainer BH, Bruce G, Hersh LB. 1989. An atlas of the regional and laminar distribution of choline acetyltransferase immunoreactivity in rat cerebral cortex. *Neuroscience* 28: 291-336.
- Ma J, Lowe G. 2005. Dendrites of juxtglomerular neurons in rat olfactory bulb express Ca²⁺-permeable AMPA receptors. *ACHEM S Abstr* 86.
- Mackay-Sim A, Chuah MI. 2000. Neurotrophic factors in the primary olfactory pathway. *Prog Neurobiol* 62: 527-559.
- MacLeod NK, Straughan DW. 1979. Responses of olfactory bulb neurones to the dipeptide carnosine. *Exp Brain Res* 34: 183-188.
- Macrides F, Schneider SP. 1982. Laminar organization of mitral and tufted cells in the main olfactory bulb of the adult hamster. *J Comp Neurol* 208: 419-430.
- Macrides F, Schoenfeld TA, Marchand JE, Clancy AN. 1985. Evidence for morphologically, neurochemically and functionally heterogeneous classes of mitral and tufted cells in the olfactory bulb. *Chem Sens* 2: 175-202.
- Maliosio M-L, Marqueze-Pouey B, Kuhse J, Betz H. 1991. Widespread expression of glycine receptor subunit mRNAs in the adult and developing rat brain. *EMBO J* 10: 2401-2409.
- Malnic B, Hirono J, Sato T, Buck LB. 1999. Combinatorial receptor codes for odors. *Cell* 96: 713-723.
- Mansour A, Meador-Woodruff JH, Bunzow JR, Civelli O, Akil H, et al. 1990a. Localization of dopamine D2 receptor mRNA and D1 and D2 receptor binding in the rat brain and pituitary: An in situ hybridization-receptor autoradiographic analysis. *J Neurosci* 10: 2587-2600.
- Mansour A, Meador-Woodruff JH, Camp DM, Robinson TE, Bunzow J, et al. 1990b. The effects of nigrostriatal 6-hydroxydopamine lesions on dopamine D2 receptor mRNA and opioid systems. *Prog Clin Biol Res* 328: 227-230.
- Marek GJ, Aghajanian GK. 1994. Excitation of interneurons in piriform cortex by 5-hydroxytryptamine: Blockade by MDL 100,907, a highly selective 5-HT_{2A} receptor antagonist. *Eur J Pharmacol* 259: 137-141.
- Marek GJ, Aghajanian GK. 1995. Protein kinase C inhibitors enhance the 5-HT_{2A} receptor-mediated excitatory effects of serotonin on interneurons in rat piriform cortex. *Synapse* 21: 123-130.
- Marek GJ, Aghajanian GK. 1996a. Alpha 1B-adrenoceptor-mediated excitation of piriform cortical interneurons. *Eur J Pharmacol* 305: 95-100.
- Marek GJ, Aghajanian GK. 1996b. LSD and the phenethylamine hallucinogen DOI are potent partial agonists at 5-HT_{2A} receptors on interneurons in rat piriform cortex. *J Pharmacol Exp Ther* 278: 1373-1382.
- Margeta-Mitrovic M, Mitrovic I, Riley RC, Jan LY, Basbaum AI. 1999. Immunohistochemical localization of GABA(B) receptors in the rat central nervous system. *J Comp Neurol* 405: 299-321.

- Margolis FL. 1972. A brain protein unique to the olfactory bulb. *Proc Natl Acad Sci USA* 69: 1221-1224.
- Margolis FL. 1974. Carnosine in the primary olfactory pathway. *Science* 184: 909-911.
- Margolis FL. 1980. Carnosine: An olfactory neuropeptide. The role of peptides in neuronal function. Barker JL, Smith TGJ, editors. New York: Marcel Dekker, Inc.; pp. 545-572.
- Margolis FL, Grillo M. 1984. Carnosine, homocarnosine and anserine in vertebrate retinas. *Neurochem Int'l* 6: 207-209.
- Margolis FL, Grillo M, Grannot-Reisfeld N, Farbman AI. 1983. Purification, characterization and immunocytochemical localization of mouse kidney carnosinase. *Biochimica et Biophysica Acta* 744: 237-248.
- Margolis FL, Grillo M, Hempstead J, Morgan JJ. 1987. Monoclonal antibodies to mammalian carnosine synthetase. *J Neurochem* 48: 593-600.
- Margolis FL, Grillo M, Kawano T, Farbman AI. 1985. Carnosine synthesis in olfactory tissue during ontogeny: Effect of exogenous β -alanine. *J Neurochem* 44: 1459-1464.
- Martin LJ, Blackstone CD, Haganir RL, Price DL. 1992. Cellular localization of a metabotropic glutamate receptor in rat brain. *Neuron* 9: 259-270.
- Martin LJ, Blackstone CD, Levey AI, Haganir RL, Price DL. 1993. Cellular localizations of AMPA glutamate receptors within the basal forebrain magnocellular complex of rat and monkey. *J Neurosci* 13: 2249-2263.
- Masu M, Tanabe Y, Tsuchida K, Shigemoto R, Nakanishi S. 1991. Sequence and expression of a metabotropic glutamate receptor. *Nature* 349: 760-765.
- Matsutani S, Senba E, Tohyama M. 1988. Neuropeptide- and neurotransmitter-related immunoreactivities in the developing rat olfactory bulb. *J Comp Neurol* 272: 331-342.
- McCune SK, Voigt MM, Hill JM. 1993. Expression of multiple alpha adrenergic receptor subtype messenger RNAs in the adult rat brain. *Neuroscience* 57: 143-151.
- McLean JH, Harley CW. 2004. Olfactory learning in the rat pup: A model that may permit visualization of a mammalian memory trace. *Neuroreport* 15: 1691-1697.
- McLean JH, Shipley MT. 1987a. Serotonergic afferents to the rat olfactory bulb: I. Origins and laminar specificity of serotonergic inputs in the adult rat. *J Neurosci* 7: 3016-3028.
- McLean JH, Shipley MT. 1987b. Serotonergic afferents to the rat olfactory bulb: II. Changes in fiber distribution during development. *J Neurosci* 7: 3029-3039.
- McLean JH, Shipley MT. 1988. Postmitotic, postmigrational expression of tyrosine hydroxylase in olfactory bulb dopaminergic neurons. *J Neurosci* 8: 3658-3669.
- McLean JH, Darby-King A, Bonnell WS. 2001. Neonatal olfactory sensory deprivation decreases BDNF in the olfactory bulb of the rat. *Dev Brain Res* 128: 17-24.
- McLean JH, Darby-King A, Hodge E. 1996. 5-HT₂ receptor involvement in conditioned olfactory learning in the neonatal rat pup. *Behav Neurosci* 110: 1426-1434.
- McLean JH, Darby-King A, Paterno GD. 1995. Localization of 5-HT_{2A} receptor mRNA by in situ hybridization in the olfactory bulb of the postnatal rat. *J Comp Neurol* 353: 371-378.
- McLean JH, Darby-King A, Sullivan RM, King SR. 1993. Serotonergic influence on olfactory learning in the neonate rat. *Behav Neural Biol* 60: 152-162.
- McLean JH, Shipley MT, Nickell WT, Aston-Jones G, Reyher CK. 1989. Chemoanatomical organization of the noradrenergic input from locus coeruleus to the olfactory bulb of the adult rat. *J Comp Neurol* 285: 339-349.
- McLennan H. 1971. The pharmacology of inhibition of mitral cells in the olfactory bulb. *Brain Res* 29: 177-184.
- McMillian MK, Mullis SB, Wu G-C, Hudson PM, Penny-packer KR, et al. 1994. Regulation of tyrosine hydroxylase in olfactory bulb cultures: Selective inhibition of depolarization-induced increase by endogenous opioids. *Brain Res* 658: 105-111.
- Mechawar N, Saghatelany A, Grailhe R, Scoriels L, Gheusi G, et al. 2004. Nicotinic receptors regulate the survival of newborn neurons in the adult olfactory bulb. *Proc Natl Acad Sci USA* 101: 9822-9826.
- Meisami E. 1989. A proposed relationship between increases in the number of olfactory receptor neurons, convergence ratio and sensitivity in the developing rat. *Brain Res Dev Brain Res* 46: 9-19.
- Meisami E, Safari L. 1981. A quantitative study of the effects of early unilateral olfactory deprivation on the number and distribution of mitral and tufted cells and of glomeruli in the rat olfactory bulb. *Brain Res* 221: 81-107.
- Meisami E, Sendera TJ. 1993. Morphometry of rat olfactory bulbs stained for cytochrome oxidase reveals that the entire population of glomeruli forms early in the neonatal period. *Brain Res Dev Brain Res* 71: 253-257.
- Menco BP. 1984. Ciliated and microvillous structures of rat olfactory and nasal respiratory epithelia. A study using ultra-rapid cryo-fixation followed by freeze-substitution or freeze-etching. *Cell Tissue Res* 235: 225-241.
- Menco BP, Farbman AI. 1985a. Genesis of cilia and microvilli of rat nasal epithelia during pre-natal development. I. Olfactory epithelium, qualitative studies. *J Cell Sci* 78: 283-310.
- Menco BP, Farbman AI. 1985b. Genesis of cilia and microvilli of rat nasal epithelia during pre-natal development. II. Olfactory epithelium, a morphometric analysis. *J Cell Sci* 78: 311-336.
- Mengod G, Nguyen H, Le H, Waeber C, Lubbert H, et al. 1990a. The distribution and cellular localization of the serotonin 1C receptor mRNA in the rodent brain examined by in situ hybridization histochemistry. Comparison with receptor binding distribution. *Neuroscience* 35: 577-591.

- Mengod G, Pompeiano M, Martinez-Mir MI, Palacios JM. 1990b. Localization of the mRNA for the 5-HT₂ receptor by in situ hybridization histochemistry. Correlation with the distribution of receptor sites. *Brain Res* 524: 139-143.
- Merchenthaler I, Csernus V, Csontos C, Petrusz P, Mess B. 1988. New data on the immunocytochemical localization of thyrotropin-releasing hormone in the rat central nervous system. *Am J Anat* 181: 359-376.
- Miller LP, Johnson AE, Gelhard RE, Insel TR. 1990. The ontogeny of excitatory amino acid receptors in the rat forebrain—II. Kainic acid receptors. *Neuroscience* 35: 45-51.
- Miragall F, Simburger E, Dermietzel R. 1996. Mitral and tufted cells of the mouse olfactory bulb possess gap junctions and express connexin43 mRNA. *Neurosci Lett* 216: 199-202.
- Mirich JM, Illig KR, Brunjes PC. 2004. Experience-dependent activation of extracellular signal-related kinase (ERK) in the olfactory bulb. *J Comp Neurol* 479: 234-241.
- Molnar E, Baude A, Richmond SA, Patel PB, Somogyi P, et al. 1993. Biochemical and immunocytochemical characterization of antipeptide antibodies to a cloned GluR1 glutamate receptor subunit: Cellular and subcellular distribution in the rat forebrain. *Neuroscience* 53: 307-326.
- Mombaerts P, Wang F, Dulac C, Chao SK, Nemes A, et al. 1996. Visualizing an olfactory sensory map. *Cell* 87: 675-686.
- Monaghan DT, Cotman CW. 1982. The distribution of [³H] kainic acid binding sites in rat CNS as determined by autoradiography. *Brain Res* 252: 91-100.
- Monaghan DT, Yao D, Cotman CW. 1985. L-[³H]Glutamate binds to kainate-, NMDA- and AMPA-sensitive binding sites: An autoradiographic analysis. *Brain Res* 340: 378-383.
- Montague AA, Greer CA. 1999. Differential distribution of ionotropic glutamate receptor subunits in the rat olfactory bulb. *J Comp Neurol* 405: 233-246.
- Monyer H, Burnashev N, Laurie DJ, Sakmann B, Seeburg PH. 1994. Developmental and regional expression in the rat brain and functional properties of four NMDA receptors. *Neuron* 12: 529-540.
- Morales M, Bloom FE. 1997. The 5-HT₃ receptor is present in different subpopulations of GABAergic neurons in the rat telencephalon. *J Neurosci* 17: 3157-3167.
- Moran DT, Rowley JC III, Jafek BW. 1982. Electron microscopy of human olfactory epithelium reveals a new cell type: The microvillar cell. *Brain Res* 253: 39-46.
- Mori K. 1987a. Membrane and synaptic properties of identified neurons in the olfactory bulb. *Prog Neurobiol* 29: 275-320.
- Mori K. 1987b. Monoclonal antibodies (2C5 and 4C9) against lactoseries carbohydrates identify subsets of olfactory and vomeronasal receptor cells and their axons in the rabbit. *Brain Res* 408: 215-221.
- Mori K, Kishi K. 1982. The morphology and physiology of the granule cells in the rabbit olfactory bulb revealed by intracellular recording and HRP injection. *Brain Res* 247: 129-133.
- Mori K, Takagi SF. 1978. Activation and inhibition of olfactory bulb neurones by anterior commissure volleys in the rabbit. *J Physiol (Lond)* 279: 589-604.
- Mori K, Kishi K, Ojima H. 1983. Distribution of dendrites of mitral, displaced mitral, tufted, and granule cells in the rabbit olfactory bulb. *J Comp Neurol* 219: 339-355.
- Moriceau S, Sullivan RM. 2004. Unique neural circuitry for neonatal olfactory learning. *J Neurosci* 24: 1182-1189.
- Morizumi T, Tsukatani H, Miwa T. 1994. Olfactory disturbance induced by deafferentation of serotonergic fibers in the olfactory bulb. *Neuroscience* 61: 733-738.
- Morrison EE, Costanzo RM. 1990. Morphology of the human olfactory epithelium. *J Comp Neurol* 297: 1-13.
- Mouly AM, Elaagouby A, Ravel N. 1995. A study of the effects of noradrenaline in the rat olfactory bulb using evoked field potential response. *Brain Res* 681: 47-57.
- Mouradian LE, Scott JW. 1988. Cytochrome oxidase staining marks dendritic zones in the rat olfactory bulb external plexiform layer. *J Comp Neurol* 271: 507-518.
- Mugnaini E, Oertel WH, Wouterlood FF. 1984a. Immunocytochemical localization of GABA neurons and dopamine neurons in the rat main and accessory olfactory bulbs. *Neurosci Lett* 47: 221-226.
- Mugnaini E, Wouterlood FG, Dahl A-L, Oertel WH. 1984b. Immunocytochemical identification of GABAergic neurons in the main olfactory bulb of the rat. *Archives Italiennes de Biologie* 122: 83-113.
- Muller JF, Marc RE. 1984. Three distinct morphological classes of receptors in fish olfactory organs. *J Comp Neurol* 222: 482-495.
- Munger SD, Lane AP, Zhong H, Leinders-Zufall T, Yau KW, et al. 2001. Central role of the CNGA4 channel subunit in Ca²⁺-calmodulin-dependent odor adaptation. *Science* 294: 2172-2175.
- Murphy GJ, Darcy DP, Isaacson JS. 2005. Intraglomerular inhibition: Signaling mechanisms of an olfactory microcircuit. *Nat Neurosci* 8: 354-364.
- Nadi NS, Head R, Grillo M, Hempstead J, Grannot-Reisfeld N, et al. 1981. Chemical deafferentation of the olfactory bulb: Plasticity of the levels of tyrosine hydroxylase, dopamine and norepinephrine. *Brain Res* 213: 365-377.
- Nagayama S, Takahashi YK, Yoshihara Y, Mori K. 2004. Mitral and tufted cells differ in the decoding manner of odor maps in the rat olfactory bulb. *J Neurophysiol* 91: 2532-2540.
- Najbauer J, Leon M. 1995. Olfactory experience modulated apoptosis in the developing olfactory bulb. *Brain Res* 674: 245-251.

- Nakajima T, Okamura M, Ogawa K, Taniguchi K. 1996. Immunohistochemical and enzyme histochemical characteristics of short axon cells in the olfactory bulb of the golden hamster. *J Vet Med Sci* 58: 903-908.
- Nakashima M, Mori K, Takagi SF. 1978. Centrifugal influence on olfactory bulb activity in the rabbit. *Brain Res* 154: 301-316.
- Nicholas AP, Pieribone V, Hokfelt T. 1993. Distributions of mRNAs for alpha-2 adrenergic receptor subtypes in rat brain: An in situ hybridization study. *J Comp Neurol* 328: 575-594.
- Nickell WT, Shipley MT. 1988a. Neurophysiology of magnocellular forebrain inputs to the olfactory bulb in the rat: Frequency potentiation of field potentials and inhibition of output neurons. *J Neurosci* 8: 4492-4502.
- Nickell WT, Shipley MT. 1988b. Two anatomically specific classes of candidate cholinceptive neurons in the rat olfactory bulb. *J Neurosci* 8: 4482-4491.
- Nickell WT, Shipley MT. 1993. Evidence for presynaptic inhibition of the olfactory commissural pathway by cholinergic agonists and stimulation of the nucleus of the diagonal band. *J Neurosci* 13: 650-659.
- Nickell WT, Norman AB, Wyatt LM, Shipley MT. 1991. Olfactory bulb DA receptors may be located on terminals of the olfactory nerve. *Neuroreport* 2: 9-12.
- Nicoll RA. 1971. Pharmacological evidence for GABA as the transmitter in granule cell inhibition in the olfactory bulb. *Brain Res* 35: 137-149.
- Nicoll RA, Alger BE. 1980. Peptides as putative excitatory neurotransmitters: Carnosine, enkephalin, substance P and TRH. *Proc R Soc Lond* 210: 133-149.
- Nowycky MC, Halasz N, Shepherd GM. 1983. Evoked field potential analysis of dopaminergic mechanisms in the isolated turtle olfactory bulb. *Neuroscience* 8: 717-722.
- Nusser Z, Kay LM, Laurent G, Homanics GE, Mody I. 2001. Disruption of GABA_A receptors on GABAergic interneurons leads to increased oscillatory power in the olfactory bulb network. *J Neurophysiol* 86: 2823-2833.
- Ohishi H, Neki A, Mizuno N. 1998. Distribution of a metabotropic glutamate receptor, mGluR2, in the central nervous system of the rat and mouse: An immunohistochemical study with a monoclonal antibody. *Neurosci Res* 30: 65-82.
- Ohishi H, Shigemoto R, Nakanishi S, Mizuno N. 1993a. Distribution of the messenger RNA for a metabotropic glutamate receptor, mGluR2, in the central nervous system of the rat. *Neuroscience* 53: 1009-1018.
- Ohishi H, Shigemoto R, Nakanishi S, Mizuno N. 1993b. Distribution of the mRNA for a metabotropic glutamate receptor (mGluR3) in the rat brain: An in situ hybridization study. *J Comp Neurol* 335: 252-266.
- Ohishi H, Akazawa C, Shigemoto R, Nakanishi S, Mizuno N. 1995. Distributions of mRNAs for L-2-amino-4-phosphono-butylate-sensitive metabotropic glutamate receptors mGluR4 and mGluR7, in the rat brain. *J Comp Neurol* 360: 555-570.
- Ohm TG, Müller H, Braak E. 1991. Calbindin-D-28k-like immunoreactive structures in the olfactory bulb and anterior olfactory nucleus of the human adult: Distribution and cell typology-partial complementarity with parvalbumin. *Neuroscience* 42: 823-840.
- Ohm TG, Braak E, Probst A, Weindl A. 1988. Neuropeptide Y-like immunoreactive neurons in the human olfactory bulb. *Brain Res* 451: 295-300.
- Ohm TG, Müller H, Ulfing N, Braak E. 1990. Glutamic-acid-decarboxylase- and parvalbumin-like-immunoreactive structures in the olfactory bulb of the human adult. *J Comp Neurol* 291: 1-8.
- Ojima H, Mori K, Kishi K. 1984. The trajectory of mitral cell axons in the rabbit olfactory cortex revealed by intracellular HRP injection. *J Comp Neurol* 230: 77-87.
- Okutani F, Kaba H, Takahashi S, Seto K. 1998. The biphasic effect of locus coeruleus noradrenergic activation on dendrodendritic inhibition in the rat olfactory bulb. *Brain Res* 783: 272-279.
- Onoda N. 1992. Odor-induced fos-like immunoreactivity in the rat olfactory bulb. *Neurosci Lett* 137: 157-160.
- Orona E, Rainer EC, Scott JW. 1984. Dendritic and axonal organization of mitral and tufted cells in the rat olfactory bulb. *J Comp Neurol* 226: 346-356.
- Orona E, Scott JW, Rainer EC. 1983. Different granule cell populations innervate superficial and deep regions of the external plexiform layer in rat olfactory bulb. *J Comp Neurol* 217: 227-237.
- Palacios JM, Kuhar MJ. 1980. Beta-adrenergic-receptor localization by light microscopic autoradiography. *Science* 208: 1378-1380.
- Palacios JM, Kuhar MJ. 1982. Beta adrenergic receptor localization in rat brain by light microscopic autoradiography. *Neurochem Int* 4: 473-490.
- Palacios JM, Niehoff DL, Kuhar MJ. 1981a. [³H]Spiperone binding sites in brain: Autoradiographic localization of multiple receptors. *Brain Res* 213: 277-289.
- Palacios JM, Wamsley JK, Kuhar MJ. 1981b. High affinity GABA receptors-autoradiographic localization. *Brain Res* 222: 285-307.
- Palouzier-Paulignan B, Duchamp-Viret P, Hardy AB, Duchamp A. 2002. GABA(B) receptor-mediated inhibition of mitral/tufted cell activity in the rat olfactory bulb: A whole-cell patch-clamp study in vitro. *Neuroscience* 111: 241-250.
- Panzanelli P, Perazzini A-Z, Fritschy J-M, Sassoè-Pognetto M. 2005. Heterogeneity of γ -aminobutyric acid type A receptors in mitral and tufted cells of the rat main olfactory bulb. *J Comp Neurol* 484: 121-131.

- Panzanelli P, Homanics GE, Ottersen OP, Fritschy J-M, Sassoè-Pognetto M. 2004. Pre- and postsynaptic GABAA receptors at reciprocal dendrodendritic synapses in the olfactory bulb. *Eur J Neurosci* 20: 2945-2952.
- Paternostro MA, Reyher CKH, Brunjes PC. 1995. Intracellular injections of Lucifer Yellow into lightly fixed mitral cells reveal neuronal dye-coupling in the developing rat olfactory bulb. *Dev Brain Res* 84: 1-10.
- Perez H, Hernandez A, Almlí CR. 1987. Locus coeruleus stimulation modulates olfactory bulb evoked potentials. *Brain Res Bull* 18: 767-770.
- Persohn E, Malherbe P, Richards JG. 1992. Comparative molecular neuroanatomy of cloned GABAA receptor subunits in the rat CNS. *J Comp Neurol* 326: 193-216.
- Petralia RS, Wenthold RJ. 1992. Light and electron immunocytochemical localization of AMPA-selective glutamate receptors in the rat brain. *J Comp Neurol* 318: 329-354.
- Petralia RS, Wang YX, Wenthold RJ. 1994a. Histological and ultrastructural localization of the kainate receptor subunits, KA2 and GluR6/7, in the rat nervous system using selective antipeptide antibodies. *J Comp Neurol* 349: 85-110.
- Petralia RS, Yokotani N, Wenthold RJ. 1994b. Light and electron microscope distribution of the NMDA receptor subunit NMDAR1 in the rat nervous system using a selective anti-peptide antibody. *J Neurosci* 14: 667-696.
- Petreanu L, Alvarez-Buylla A. 2002. Maturation and death of adult-born granule neurons: Role of olfaction. *J Neurosci* 22: 6106-6113.
- Philpot BD, Lim JH, Brunjes PC. 1997. Activity-dependent regulation of calcium-binding proteins in the developing rat olfactory bulb. *J Comp Neurol* 387: 12-26.
- Philpot BD, Men D, McCarty R, Brunjes. 1998. Activity-dependent regulation of dopamine content in the olfactory bulbs of naris-occluded rats. *Neuroscience* 85: 969-977.
- Pieribone VA, Nicholas AP, Dagerlind A, Hokfelt T. 1994. Distribution of $\alpha 1$ adrenoreceptors in rat brain revealed by in situ hybridization experiments utilizing subtype-specific probes. *J Neurosci* 14: 4252-4268.
- Pinching AJ, Powell TP. 1971a. The neuron types of the glomerular layer of the olfactory bulb. *J Cell Sci* 9: 305-345.
- Pinching AJ, Powell TP. 1971b. The neuropil of the glomeruli of the olfactory bulb. *J Cell Sci* 9: 347-377.
- Pinching AJ, Powell TP. 1971c. The neuropil of the periglomerular region of the olfactory bulb. *J Cell Sci* 9: 379-409.
- Pinching AJ, Powell TP. 1972a. A study of terminal degeneration in the olfactory bulb of the rat. *J Cell Sci* 10: 585-619.
- Pinching AJ, Powell TP. 1972b. Experimental studies on the axons intrinsic to the glomerular layer of the olfactory bulb. *J Cell Sci* 10: 637-655.
- Pinching AJ, Powell TP. 1972c. The termination of centrifugal fibres in the glomerular layer of the olfactory bulb. *J Cell Sci* 10: 621-635.
- Pissonnier D, Thiery JC, Fabre-Nys C, Poindron P, Keverne EB. 1985. The importance of olfactory bulb noradrenalin for maternal recognition in sheep. *Physiol Behav* 35: 361-363.
- Pompeiano M, Palacios JM, Mengod G. 1992. Distribution and cellular localization of mRNA coding for 5-HT1A receptor in the rat brain: Correlation with receptor binding. *J Neurosci* 12: 440-453.
- Pompeiano M, Palacios JM, Mengod G. 1994. Distribution of the serotonin 5-HT2 receptor family mRNAs: Comparison between 5-HT2A and 5-HT2C receptors. *Brain Res Mol Brain Res* 23: 163-178.
- Porteros A, Brinon JG, Crespo C, Okazaki K, Hidaka H, et al. 1996. Neurocalcin immunoreactivity in the rat accessory olfactory bulb. *Brain Res* 729: 82-89.
- Postlewaither M, Constanti A, Libri A. 1998. Muscarinic agonist-induced burst firing in immature olfactory cortex neurons in vitro. *J Neurophysiol* 79: 2003-2012.
- Potter SM, Zheng C, Koos DS, Feinstein P, Fraser SE, et al. 2001. Structure and emergence of specific olfactory glomeruli in the mouse. *J Neurosci* 21: 9713-9723.
- Pourcho RG, Goebel DJ, Jojich L, Hazlett JC. 1992. Immunocytochemical evidence for the involvement of glycine in sensory centers of the rat brain. *Neuroscience* 46: 643-656.
- Pow DV, Sullivan R, Reye P, Hermanussen S. 2002. Localization of taurine transporters, taurine, and 3H taurine accumulation in the rat retina, pituitary, and brain. *Glia* 37: 153-168.
- Pressler RT, Strowbridge BW. 2006. Blanes cells mediate persistent feedforward inhibition onto granule cells in the olfactory bulb. *Neuron* 49: 889-904.
- Price JL, Powell TPS. 1970a. An electron-microscopic study of the termination of the afferent fibres to the olfactory bulb from the cerebral hemisphere. *J Cell Sci* 7: 157-187.
- Price JL, Powell TPS. 1970b. The mitral and short axon cells of the olfactory bulb. *J Cell Sci* 7: 631-651.
- Price JL, Powell TPS. 1970c. The morphology of the granule cells of the olfactory bulb. *J Cell Sci* 9: 91-123.
- Price JL, Powell TPS. 1970d. The synaptology of the granule cells of the olfactory bulb. *J Cell Sci* 7: 125-155.
- Price JL, Slotnick BM. 1983. Dual olfactory representation in the rat thalamus: An anatomical and electrophysiological study. *J Comp Neurol* 215: 63-77.
- Puche AC, Shipley MT. 1999. Odor-induced, activity-dependent transneuronal gene induction in vitro: Mediation by NMDA receptors. *J Neurosci* 19: 1359-1370.

- Puopolo M, Bean BP, Raviola E. 2005. Spontaneous activity of isolated dopaminergic periglomerular cells of the main olfactory bulb. *J Neurophysiol* 94: 3618-3627.
- Rall W, Shepherd GM, Reese TS, Brightman MW. 1966. Dendrodendritic synaptic pathway for inhibition in the olfactory bulb. *Exp Neurol* 14: 44-56.
- Rash JE, Staines WA, Yasumura T, Patel D, Furman DS, et al. 2000. Immunogold evidence that the neuronal gap junctions in adult rat brain and spinal cord contain connexin-36 but not connexin-32 or connexin-43. *Proc Natl Acad Sci USA* 97: 7573-7578.
- Rash JE, Davidson KG, Kamasawa N, Yasumura T, Kamasawa M, Zhang C, Michaels R, Restrepo D, Ottersen OP, Olson CO, Nagy JI. 2005. Ultrastructural localization of connexins (Cx36, Cx43, Cx45), glutamate receptors and aquaporin-4 in rodent olfactory mucosa, olfactory nerve and olfactory bulb. *J Neurocytol* 34: 307-341.
- Rawson NE, Eberwine J, Dotson R, Jackson J, Ulrich P, et al. 2000. Expression of mRNAs encoding for two different olfactory receptors in a subset of olfactory receptor neurons. *J Neurochem* 75: 185-195.
- Reed RR. 1992. Signaling pathways in odorant detection. *Neuron* 8: 205-209.
- Ressler KJ, Sullivan SL, Buck LB. 1993. A zonal organization of odorant receptor gene expression in the olfactory epithelium. *Cell* 73: 597-609.
- Ressler KJ, Sullivan SL, Buck LB. 1994. Information coding in the olfactory system: Evidence for a stereotyped and highly organized epitope map in the olfactory bulb. *Cell* 79: 1245-1255.
- Reyher CK, Lubke J, Larsen WJ, Hendrix GM, Shipley MT, et al. 1991. Olfactory bulb granule cell aggregates: Morphological evidence for interperikaryal electrotonic coupling via gap junctions. *J Neurosci* 11: 1485-1495.
- Ribak CE, Vaughn JE, Saito K, Barber R, Roberts E. 1977. Glutamate decarboxylase localization in neurons of the olfactory bulb. *Brain Res* 126: 1-18.
- Richards JG, Schoch P, Haring P, Takacs B, Mohler H. 1987. Resolving GABAA/benzodiazepine receptors: Cellular and subcellular localization in the CNS with monoclonal antibodies. *J Neurosci* 7: 1866-1886.
- Roberts GW, Woodhams PL, Polak JM, Crow TJ. 1982. Distribution of neuropeptides in the limbic system of the rat: The amygdaloid complex. *Neuroscience* 7: 99-131.
- Rocheffort C, Gheusi G, Vincent J-D, Lledo P-M. 2002. Enriched odor exposure increases the number of newborn neurons in the adult olfactory bulb and improves odor memory. *J Neurosci* 22: 2679-2689.
- Rochel S, Margolis FL. 1982. Carnosine release from olfactory bulb synaptosomes is calcium-dependent and depolarization-stimulated. *J Neurochem* 38: 505-1514.
- Rodriguez R, Haberly LB. 1989. Analysis of synaptic events in the opossum piriform cortex with improved current source-density techniques. *J Neurophysiol* 61: 702-718.
- Rogers JH. 1992. Immunohistochemical markers in rat brain: Colocalization of calretinin and calbindin-D28k with tyrosine hydroxylase. *Brain Res* 587: 203-210.
- Romano C, Sesma MA, McDonald CT, O'Malley K, van den Pol AN, et al. 1995. Distribution of metabotropic glutamate receptor mGluR5 immunoreactivity in rat brain. *J Comp Neurol* 355: 455-469.
- Rosin DL, Talley EM, Lee A, Stornetta RL, Gaylenn BD, et al. 1996. Distribution of alpha 2C-adrenergic receptor-like immunoreactivity in the rat central nervous system. *J Comp Neurol* 372: 135-165.
- Rosin J-F, Datiche F, Cattarelli M. 1999. Modulation of the piriform cortex activity by the basal forebrain: An optical recording study in the rat. *Brain Res* 820: 105-111.
- Ross CD, Godfrey DA, Parli JA. 1995. Amino acid concentrations and selected enzyme activities in rat auditory, olfactory, and visual systems. *Neurochem Res* 20: 1483-1490.
- Rosser AE, Keverne EB. 1985. The importance of central noradrenergic neurones in the formation of an olfactory memory in the prevention of pregnancy block. *Neuroscience* 15: 1141-1147.
- Rotter A, Birdsall NJ, Field PM, Raisman G. 1979. Muscarinic receptors in the central nervous system of the rat. II. Distribution of binding of [3H]propylbenzylcholine mustard in the midbrain and hindbrain. *Brain Res* 180: 167-183.
- Rowley JC III, Moran DT, Jafek BW. 1989. Peroxidase backfills suggest the mammalian olfactory epithelium contains a second morphologically distinct class of bipolar sensory neuron: The microvillar cell. *Brain Res* 502: 387-400.
- Royet J, Distel H, Hudson R, Gervais R. 1998. A re-estimation of the number of glomeruli and mitral cells in the olfactory bulb of rabbit. *Brain Res* 788: 35-42.
- Royet JB, Souchier C, Jourdan F, Ploye H. 1988. Morphometric study of the glomerular population in the mouse olfactory bulb: Numerical density and size distribution along the rostrocaudal axis. *J Comp Neurol* 270: 559-568.
- Rozental R, Srinivas M, Gökhan S, Urban M, Dermietzel R, et al. 2000. Temporal expression of neuronal connexins during hippocampal ontogeny. *Brain Res Rev* 32: 57-71.
- Rubin BD, Katz LC. 1999. Optical imaging of odorant representations in the mammalian olfactory bulb. *Neuron* 23: 499-511.
- Rubin BD, Katz LC. 2001. Spatial coding of enantiomers in the rat olfactory bulb. *Nat Neurosci* 4: 355-356.
- Saghatelian A, de Chevigny A, Schachner M, Lledo P-M. 2004. Tenascin-R mediates activity-dependent recruitment of neuroblasts in the adult mouse forebrain. *Nat Neurosci* 7: 347-356.

- Sahara Y, Kubota T, Ichikawa M. 2001. Cellular localization of metabotropic glutamate receptors mGluR1, 2/3, 5 and 7 in the main and accessory olfactory bulb of the rat. *Neurosci Lett* 312: 59-62.
- Sahin M, Bowen WD, Donoghue JP. 1992. Location of nicotinic and muscarinic cholinergic and mu-opiate receptors in rat cerebral neocortex: Evidence from thalamic and cortical lesions. *Brain Res* 579: 135-147.
- Saito N, Kikkawa U, Nishizuka Y, Tanaka C. 1988. Distribution of protein kinase C-like immunoreactive neurons in rat brain. *J Neurosci* 8: 369-382.
- Salin PA, Lledo PM, Vincent JD, Chrapak S. 2001. Dendritic glutamate autoreceptors modulate signal processing in rat mitral cells. *J Neurophysiol* 85: 1275-1282.
- Sallaz M, Jourdan F. 1992. Apomorphine disrupts odour-induced patterns of glomerular activation in the olfactory bulb. *Neuroreport* 3: 833-836.
- Salmoiraghi GC, Bloom FE, Costa E. 1964. Adrenergic mechanisms in rabbit olfactory bulb. *Am J Physiol* 207: 1417-1424.
- Sanides-Kohlrausch C, Wahle P. 1990a. Morphology of neuropeptide Y-immunoreactive neurons in the cat olfactory bulb and olfactory peduncle: Postnatal development and species comparison. *J Comp Neurol* 291: 468-489.
- Sanides-Kohlrausch C, Wahle P. 1990b. VIP- and PHI-immunoreactivity in olfactory centers of the adult cat. *J Comp Neurol* 294: 325-339.
- Sargent-Jones L, Gauger LL, Davis JN. 1985. Anatomy of brain alpha1-adrenergic receptors: In vitro autoradiography with [¹²⁵I]-heat. *J Comp Neurol* 231: 190-208.
- Sashihara S, Waxman SG, Greer CA. 1997. Downregulation of Na⁺ channel mRNA in olfactory tufted cells following deafferentation. *Neuroreport* 8: 1289-1293.
- Sassoè-Pognetto M, Ottersen OP. 2000. Organization of ionotropic glutamate receptors at dendrodendritic synapses in the rat olfactory bulb. *J Neurosci* 20: 2192-2201.
- Sassoè-Pognetto M, Cantino D, Panzanelli P. 1993. Presynaptic co-localization of carnosine and glutamate in olfactory neurons. *Neuroreport* 5: 7-10.
- Sassoè-Pognetto M, Panzanelli P, Siegert W, Fritschy J-M. 2000. Colocalization of multiple GABAA receptor subtypes with gephyrin at postsynaptic sites. *J Comp Neurol* 420: 481-498.
- Sassoè-Pognetto M, Utvik JK, Camoletto P, Watanabe M, Stephenson FA, et al. 2003. Organization of postsynaptic density proteins and glutamate receptors in axodendritic dendrodendritic synapses of the rat olfactory bulb. *J Comp Neurol* 463: 237-248.
- Satou M, Mori K, Tazawa Y, Takagi SF. 1983a. Interneurons mediating fast postsynaptic inhibition in pyriform cortex of the rabbit. *J Neurophysiol* 50: 89-101.
- Satou M, Mori K, Tazawa Y, Takagi SF. 1983b. Monosynaptic and disynaptic activation of pyriform cortex neurons by synchronous lateral olfactory tract volleys in the rabbit. *Exp Neurol* 81: 571-585.
- Satou M, Mori K, Tazawa Y, Takagi SF. 1983c. Neuronal pathways for activation of inhibitory interneurons in pyriform cortex of the rabbit. *J Neurophysiol* 50: 74-88.
- Saucier D, Astic L. 1986. Analysis of the topographical organization of olfactory epithelium projections in the rat. *Brain Res Bull* 16: 455-462.
- Saugstad JA, Kinzie JM, Shinohara MM, Segerson TP, Westbrook GL. 1997. Cloning and expression of rat metabotropic glutamate receptor 8 reveals a distinct pharmacological profile. *Mol Pharmacol* 51: 119-125.
- Schild D, Restrepo D. 1998. Transduction mechanisms in vertebrate olfactory receptor cells. *Physiol Rev* 78: 429-466.
- Schneider SP, Macrides F. 1978. Laminar distributions of interneurons in the main olfactory bulb of the adult hamster. *Brain Res Bull* 3: 73-82.
- Schneider SP, Scott JW. 1983. Orthodromic response properties of rat olfactory mitral and tufted cells correlate with their projection patterns. *J Neurophysiol* 50: 358-378.
- Schoenfeld TA, Macrides F. 1984. Topographic organization of connections between the main olfactory bulb and pars externa of the anterior olfactory nucleus in the hamster. *J Comp Neurol* 227: 121-135.
- Schoenfeld TA, Marchand JE, Macrides F. 1985. Topographic organization of tufted cell axonal projections in the hamster main olfactory bulb: An intrabulbar associational system. *J Comp Neurol* 235: 503-518.
- Schoenfeld TA, Clancy AN, Forbes WB, Macrides F. 1994. The spatial organization of the peripheral olfactory system of the hamster. Part I: Receptor neuron projections to the main olfactory bulb. *Brain Res Bull* 34: 183-210.
- Schoppa NE, Westbrook GL. 1997. Modulation of mEPSCs in olfactory bulb mitral cells by metabotropic glutamate receptors. *J Neurophysiol* 78: 1468-1475.
- Schoppa NE, Westbrook GL. 1999. Regulation of synaptic timing in the olfactory bulb by an A-type potassium current. *Nat Neurosci* 2: 1106-13.
- Schoppa NE, Westbrook GL. 2001. Glomerulus-specific synchronization of mitral cells in the olfactory bulb. *Neuron* 31: 639-651.
- Schoppa NE, Westbrook GL. 2002. AMPA receptors drive correlated spiking in olfactory bulb glomeruli. *Nat Neurosci* 5: 1194-1202.
- Schoppa NE, Kinzie JM, Sahara Y, Segerson TP, Westbrook GL. 1998. Dendrodendritic inhibition in the olfactory bulb is driven by NMDA receptors. *J Neurosci* 18: 6790-6802.
- Scott JW. 1986. The olfactory bulb and central pathways. *Experientia* 42: 223-232.

- Scott JW, Harrison TA. 1987. The olfactory bulb: Anatomy and physiology. *Neurobiology of taste and smell*. Finger TE, Silver WF, editors. NY: Wiley; pp. 151-178.
- Scott JW, McDonald JK, Pemberton JL. 1987. Short axon cells of the rat olfactory bulb display NADPH-diaphorase activity, neuropeptide Y-like immunoreactivity, and somatostatin-like immunoreactivity. *J Comp Neurol* 260: 378-391.
- Seguela P, Datiche J, Donnelley-Miller K, Danni JA, Patrick JW. 1993. Molecular cloning, functional properties, and distribution of rat brain alpha 7: a nicotinic cation channel highly permeable to calcium. *J Neurosci* 13: 596-604.
- Seroogy KB, Brecha N, Gall C. 1985. Distribution of cholecystokinin-like immunoreactivity in the rat main olfactory bulb. *J Comp Neurol* 239: 373-383.
- Seroogy K, Hokfelt T, Buchan A, Brown JC, Terenius L, et al. 1989. Somatostatin-like immunoreactivity in rat main olfactory bulb: Extent of coexistence with neuropeptide Y-, tyrosine hydroxylase- and vitamin D-dependent calcium binding protein-like immunoreactivities. *Brain Res* 496: 389-396.
- Sheldon PW, Aghajanian GK. 1990. Serotonin (5-HT) induces IPSPs in pyramidal layer cells of rat piriform cortex: Evidence for the involvement of a 5-HT₂-activated interneuron. *Brain Res* 506: 62-69.
- Sheldon PW, Aghajanian GK. 1991. Excitatory responses to serotonin (5-HT) in neurons of the rat piriform cortex: Evidence for mediation by 5-HT_{1C} receptors in pyramidal cells and 5-HT₂ receptors in interneurons. *Synapse* 9: 208-218.
- Shepherd GM. 1972. Synaptic organization of the mammalian olfactory bulb. *Physiol Rev* 52: 864-917.
- Shepherd GM, Greer CA. 1998. The olfactory bulb. The synaptic organization of the brain. Shepherd GM, editor. New York: Oxford UP; pp. 159-203.
- Shepherd GM, Chen WR, Greer CA. 2004. Olfactory bulb. The synaptic organization of the brain, 5th Edn. Shepherd GM, editor. New York: Oxford Univ. Press; pp. 165-216.
- Shigemoto R, Nakanishi S, Mizuno N. 1992. Distribution of the mRNA for a metabotropic glutamate receptor (mGluR1) in the central nervous system: An in situ hybridization study in adult and developing rat. *J Comp Neurol* 322: 121-135.
- Shigemoto R, Nomura S, Ohishi H, Sugihara H, Nakanishi S, et al. 1993. Immunohistochemical localization of a metabotropic glutamate receptor, mGluR5, in the rat brain. *Neurosci Lett* 163: 53-57.
- Shibley MT, Adamek GD. 1984. The connections of the mouse olfactory bulb: A study using orthograde and retrograde transport of wheat germ agglutinin conjugated to horseradish peroxidase. *Brain Res Bull* 12: 669-688.
- Shibley MT, Ennis M, Puche A. 2004. The olfactory system. The rat nervous system. Paxinos G, editor. New York: Academic Press; pp. 923-964.
- Shibley MT, Halloran FJ, de la Torre J. 1985. Surprisingly rich projection from locus coeruleus to the olfactory bulb in the rat. *Brain Res* 329: 294-299.
- Shibley MT, McLean JH, Zimmer LA, Ennis M. 1996. The olfactory system. Integrated systems of the CNS, part III. Swanson LW, Bjorklund A, Hokfelt T, editors. New York: Elsevier; pp. 469-575.
- Shults CW, Quirion R, Chronwall B, Chase TN, O'Donohue TL. 1984. A comparison of the anatomical distribution of substance P and substance P receptors in the rat central nervous system. *Peptides* 5: 1097-1128.
- Siklos L, Rickmann M, Joo F, Freeman WJ, Wolff JR. 1995. Chloride is preferentially accumulated in a subpopulation of dendrites and periglomerular cells of the main olfactory bulb in adult rats. *Neuroscience* 64: 165-172.
- Smith C, Luskin MB. 1998. Cell cycle length of olfactory bulb neuronal progenitors in the rostral migratory stream. *Dev Dyn* 213: 220-227.
- Smith TC, Jahr CE. 2002. Self-inhibition of olfactory bulb neurons. *Nat Neurosci* 5: 760-766.
- Smith RL, Baker H, Kolstad K, Spencer DD, Greer CA. 1991. Localization of tyrosine hydroxylase and olfactory marker protein immunoreactivities in the human and cacaque olfactory bulb. *Brain Res* 548: 140-148.
- Spencer DG Jr, Horvath E, Traber J. 1986. Direct autoradiographic determination of M1 and M2 muscarinic acetylcholine receptor distribution in the rat brain: Relation to cholinergic nuclei and projections. *Brain Res* 380: 59-68.
- Stenman J, Toresson H, Campbell K. 2003. Identification of two distinct progenitor populations in the lateral ganglionic eminence: Implications for striatal and olfactory bulb neurogenesis. *J Neurosci* 23: 167-174.
- Stone DM, Wessel T, Joh TH, Baker H. 1990. Decrease in tyrosine hydroxylase, but not aromatic L-amino acid decarboxylase, messenger RNA in rat olfactory bulb following neonatal, unilateral odor deprivation. *Brain Res Mol Brain Res* 8: 291-300.
- Strotmann J, Wanner I, Helfrich T, Beck A, Breer H. 1994. Rostro-caudal patterning of receptor-expressing olfactory neurones in the rat nasal cavity. *Cell Tissue Res* 278: 11-20.
- Sullivan SL, Ressler KJ, Buck LB. 1995. Spatial patterning and information coding in the olfactory system. *Curr Opin Genet Dev* 5: 516-523.
- Sullivan RM, Wilson DA, Leon M. 1989. Norepinephrine and learning-induced plasticity in infant rat olfactory system. *J Neurosci* 9: 3998-4006.
- Sullivan RM, Zyzak DR, Skierkowski P, Wilson DA. 1992. The role of olfactory bulb norepinephrine in early olfactory learning. *Brain Res Dev* 70: 279-282.
- Sullivan RM, Stackenwalt G, Nasr F, Lemon C, Wilson DA. 2000. Association of an odor with activation of olfactory bulb noradrenergic beta-receptors or locus coeruleus

- stimulation is sufficient to produce learned approach responses to that odor in neonatal rats. *Behav Neurosci* 114: 957-962.
- Sun L, Shipley MT, Lidow MS. 2000. Expression of NR1, NR2A-D, and NR3 subunits of the NMDA receptor in the cerebral cortex and olfactory bulb of adult rat. *Synapse* 35: 212-221.
- Talley EM, Rosin DL, Lee A, Guyenet PG, Lynch KR. 1996. Distribution of α 2a-adrenergic receptor like immunoreactivity in the rat central nervous system. *J Comp Neurol* 372: 111-134.
- Tanabe Y, Masu M, Ishii T, Shigemoto R, Nakanishi S. 1992. A family of metabotropic glutamate receptors. *Neuron* 8: 169-179.
- Tang AC, Hasselmo ME. 1994. Selective suppression of intrinsic but not afferent fiber synaptic transmission by baclofen in the piriform (olfactory) cortex. *Brain Res* 659: 75-81.
- Tecott LH, Maricq AV, Julius D. 1993. Nervous system distribution of the serotonin 5-HT₃ receptor mRNA. *Proc Natl Acad Sci USA* 90: 1430-1434.
- Teicher MH, Stewart WB, Kauer JS, Shepherd GM. 1980. Suckling pheromone stimulation of a modified glomerular region in the developing rat olfactory bulb revealed by the 2-deoxyglucose method. *Brain Res* 194: 530-535.
- Teubner B, Degen J, Söhl G, Güldenagel M, Bukauskas FF, et al. 2000. Functional expression of the murine connexin 36 gene coding for a neuron-specific gap junctional protein. *J Membrane Biol* 176: 249-263.
- Theis M, Söhl G, Speidel D, Willecke K. 2003. Connexin43 is not expressed in principal cells of mouse cortex and hippocampus. *Eur J Neurosci* 18: 267-274.
- Toida K, Kosaka K, Aika Y, Kosaka T. 2000. Chemically defined neuron groups and their subpopulations in the glomerular layer of the rat main olfactory bulb-IV. Intraglomerular synapses of tyrosine hydroxylase-immunoreactive neurons. *Neuroscience* 101: 11-17.
- Toida K, Kosaka K, Heizmann CW, Kosaka T. 1994. Synaptic contacts between mitral/tufted cells and GABAergic neurons containing calcium-binding protein parvalbumin in the rat olfactory bulb, with special reference to reciprocal synapses between them. *Brain Res* 650: 347-352.
- Toida K, Kosaka K, Heizmann CW, Kosaka T. 1996. Electron microscopic serial-sectioning/reconstruction study of parvalbumin-containing neurons in the external plexiform layer of the rat olfactory bulb. *Neuroscience* 72: 449-466.
- Toida K, Kosaka K, Heizmann CW, Kosaka T. 1998. Chemically defined neuron groups and their subpopulations in the glomerular layer of the rat main olfactory bulb: III. Structural features of calbindin D28K-immunoreactive neurons. *J Comp Neurol* 392: 179-198.
- Treloar HB, Feinstein P, Mombaerts P, Greer CA. 2002. Specificity of glomerular targeting by olfactory sensory axons. *J Neurosci* 22: 2469-2477.
- Trombley PQ. 1992. Norepinephrine inhibits calcium currents and EPSPs via a G-protein-coupled mechanism in olfactory bulb neurons. *J Neurosci* 12: 3992-3998.
- Trombley PQ, Shepherd GM. 1992. Noradrenergic inhibition of synaptic transmission between mitral and granule cells in mammalian olfactory bulb cultures. *J Neurosci* 12: 3985-3991.
- Trombley PQ, Shepherd GM. 1994. Glycine exerts potent inhibitory actions on mammalian olfactory bulb neurons. *J Neurophysiol* 71: 761-767.
- Trombley PQ, Shepherd GM. 1996. Differential modulation of zinc and copper of amino acid receptors from rat olfactory bulb neurons. *J Neurophysiol* 76: 2536-2546.
- Trombley PQ, Westbrook GL. 1992. L-AP4 inhibits calcium currents and synaptic transmission via a G-protein-coupled glutamate receptor. *J Neurosci* 12: 2043-2050.
- Trombley PQ, Horning MS, Blakemore LJ. 1998. Carnosine modulates zinc and copper effects on amino acid receptors and synaptic transmission. *Neuroreport* 9: 3503-3507.
- Trombley PQ, Hill BJ, Horning MS. 1999. Interactions between GABA and glycine at inhibitory amino acid receptors on rat olfactory bulb neurons. *J Neurophysiol* 82: 3417-3422.
- Tseng G-F, Haberly LB. 1988. Characterization of synaptically mediated fast and slow inhibitory processes in piriform cortex in an in vitro slice preparation. *J Neurophysiol* 59: 1352-1376.
- Tseng G-F, Haberly LB. 1989. Deep neurons in piriform cortex. II. Membrane properties that underlie unusual synaptic responses. *J Neurophysiol* 62: 386-400.
- Tsuruo Y, Hokfelt T, Visser TJ. 1988. Thyrotropin-releasing hormone (TRH)-immunoreactive neuron populations in the rat olfactory bulb. *Brain Res* 447: 183-187.
- Tucker K, Fadool DA. 2002. Neurotrophin modulation of voltage-gated potassium channels in rat through TrkB receptors is time and sensory experience dependent. *J Physiol (Lond)* 542.2: 413-429.
- Uchida N, Takahashi YK, Tanifuji M, Mori K. 2000. Odor maps in the mammalian olfactory bulb: Domain organization and odorant structural features. *Nat Neurosci* 3: 1035-1043.
- Unnerstall JR, Kopajtic TA, Kuhar MJ. 1984. Distribution of alpha 2 agonist binding sites in the rat and human central nervous system: Analysis of some functional, anatomic correlates of the pharmacologic effects of clonidine and related adrenergic agents. *Brain Res* 319: 69-101.
- Urban NN, Sakmann B. 2002. Reciprocal intraglomerular excitation and intra- and interglomerular lateral inhibition

- between mouse olfactory bulb mitral cells. *J Physiol (Lond)* 542: 355-367.
- Valverde F. 1965. *Studies on the piriform lobe*. Cambridge, MA: Harvard Univ Press.
- van den Pol AN. 1995. Presynaptic metabotropic glutamate receptors in adult and developing neurons: Autoexcitation in the olfactory bulb. *J Comp Neurol* 359: 253-271.
- van den Pol AN, Gorcs T. 1988. Glycine and glycine receptor immunoreactivity in brain and spinal cord. *J Neurosci* 8: 472-492.
- Van Gehuchten LE, Martin A. 1891. Le bulbe olfactif chez quelques mammifères. *Cellule* 205-237.
- Van Pett K, Viau V, Bittencourt JC, Chan RKW, Li H-Y, et al. 2000. Distribution of mRNAs encoding CRF receptors in the brain and pituitary of rat and mouse. *J Comp Neurol* 428: 191-212.
- Vassar R, Ngai J, Axel R. 1993. Spatial segregation of odorant receptor expression in the mammalian olfactory epithelium. *Cell* 74: 309-318.
- Vassar R, Chao SK, Sitcheran R, Nunez JM, Vosshall LB, et al. 1994. Topographic organization of sensory projections to the olfactory bulb. *Cell* 79: 981-991.
- Vertes RP. 1991. A PHA-L analysis of ascending projections of the dorsal raphe nucleus in the rat. *J Comp Neurol* 313: 643-668.
- Villalba RM, Rodrigo J, Alvarez FJ, Achaval M, Martinez-Murillo R. 1989. Localization of C-PON immunoreactivity in the rat main olfactory bulb. Demonstration that the population of neurons containing endogenous C-PON display NADPH-diaphorase activity. *Neuroscience* 33: 373-382.
- von Baumgarten R, Green JD, Mancina M. 1962. Slow waves in the olfactory bulb and their relation to unitary discharges. *Electroenceph Clin Neurophysiol* 14: 621-634.
- Wachowiak M, Cohen LB. 1999. Presynaptic inhibition of primary olfactory afferents mediated by different mechanisms in lobster and turtle. *J Neurosci* 19: 8808-8817.
- Wachowiak M, Cohen LB. 2001. Representation of odorants by receptor neuron input to the mouse olfactory bulb. *Neuron* 32: 723-735.
- Wachowiak M, Zochowski M, Cohen LB, Falk CX. 2000. The spatial representation of odors by olfactory receptor neuron input to the olfactory bulb is concentration invariant. *Biol Bull* 199: 162-163.
- Wada E, Shigemoto R, Kinoshita A, Ohishi H, Mizuno N. 1998. Metabotropic glutamate receptor subtypes in axon terminals of projection fibers from the main and accessory olfactory bulbs: A light and electron microscopic immunohistochemical study in the rat. *J Comp Neurol* 393: 493-504.
- Wahle P, Sanides-Kohlrausch S, Meyer G, Lubke J. 1990. Substance P- and opioid immunoreactive structures in olfactory centers of the cat: Adult pattern and postnatal development. *J Comp Neurol* 302: 349-369.
- Walsh RR. 1959. Olfactory bulb potentials evoked by electrical stimulation of the contralateral bulb. *Am J Physiol* 196: 327-329.
- Wanaka A, Kiyama H, Murakami T, Matsumoto M, Kamada T, et al. 1989. Immunocytochemical localization of beta-adrenergic receptors in the rat brain. *Brain Res* 485: 125-140.
- Wang F, Nemes A, Mendelsohn M, Axel R. 1998. Odorant receptors govern the formation of a precise topographic map. *Cell* 93: 47-60.
- Warden MK, Young WS. 1988. Distribution of cells containing mRNAs encoding substance P and neurokinin B in the rat central nervous system. *J Comp Neurol* 272: 90-113.
- Watanabe K, Kawana E. 1984. Selective retrograde transport of tritiated D-aspartate from the olfactory bulb to the anterior olfactory nucleus, pyriform cortex and nucleus of the lateral olfactory tract in the rat. *Brain Res* 296: 148-151.
- Watanabe M, Inoue Y, Sakimura K, Mishina M. 1993. Distinct distributions of five N-methyl-D-aspartate receptor channel subunit mRNAs in the forebrain. *J Comp Neurol* 338: 377-390.
- Wellis DP, Kauer JS. 1994. GABAergic and glutamatergic synaptic input to identified granule cells in salamander olfactory bulb. *J Physiol (Lond)* 475.
- Wellis DP, Scott JW. 1990. Intracellular responses of identified rat olfactory bulb interneurons to electrical and odor stimulation. *J Neurophysiol* 64: 932-947.
- Wellis DP, Scott JW, Harrison TA. 1989. Discrimination among odorants by single neurons of the rat olfactory bulb. *J Neurophysiol* 61: 1161-1177.
- Westenbroek RE, Westrum LE, Hendrickson AE, Wu JY. 1987. Immunocytochemical localization of cholecystokinin and glutamic acid decarboxylase during normal development in the prepyriform cortex of rats. *Brain Res* 431: 191-206.
- Whitaker-Azmitia PM, Clarke C, Azmitia EC. 1993. Localization of 5-HT_{1A} receptors to astroglial cells in adult rats: Implications for neuronal-glia interactions and psychoactive drug mechanism of action. *Synapse* 14: 201-205.
- White EL. 1972. Synaptic organization in the olfactory glomerulus of the mouse. *Brain Res* 37: 69-80.
- Whittemore ER, Koerner JF. 1989. An explanation for the purported excitation of piriform cortical neurons by N-acetyl-L-aspartyl-L-glutamic acid (NAAG). *Proc Natl Acad Sci USA* 86: 9602-9605.
- Wichterle H, Turnbull DH, Nery S, Fishell G, Alvareq-Buylla A. 2001. In utero fate mapping reveals distinct migratory pathways and fates of neurons born in the mammalian basal forebrain. *Development* 128: 3759-3771.
- Wilson DA. 1995. NMDA receptors mediate expression of one form of functional plasticity induced by olfactory deprivation. *Brain Res* 677: 238-242.

- Wilson DA. 2001. Scopolamine enhances generalization between odor representations in rat olfactory cortex. *Learn Mem* 8: 279-85.
- Wilson M, Bower JM. 1992. Cortical oscillations and temporal interactions in a computer simulation of piriform cortex. *J Neurophysiol* 67: 981-995.
- Wilson DA, Leon M. 1988. Noradrenergic modulation of olfactory bulb excitability in the postnatal rat. *Dev Brain Res* 42: 69-75.
- Wilson DA, Sullivan RM. 1991. Olfactory associative conditioning in infant rats with brain stimulation as reward: II. Norepinephrine mediates a specific component of the bulb response to reward. *Behav Neurosci* 105: 843-849.
- Wilson DA, Sullivan RM. 1995. The D2 antagonist spiperone mimics the effects of olfactory deprivation on mitral/tufted cell odor response patterns. *J Neurosci* 15: 5574-5581.
- Wilson DA, Wood JG. 1992. Functional consequences of unilateral olfactory deprivation: Time-course and age sensitivity. *Neuroscience* 49: 183-192.
- Winzer-Serhan UH, Raymon HK, Broide RS, Chen Y, Leslie FM. 1997a. Expression of α_2 adrenoreceptors during rat brain development-I. α_{2A} messenger RNA expression. *Neuroscience* 76: 241-260.
- Winzer-Serhan UH, Raymon HK, Broide RS, Chen Y, Leslie FM. 1997b. Expression of α_2 adrenoreceptors during rat brain development-II. α_{2C} messenger RNA expression and [3 H]rauwolscine binding. *Neuroscience* 76: 261-272.
- Wisden W, Seeburg PH. 1993. A complex mosaic of high-affinity kainate receptors in rat brain. *J Neurosci* 13: 3582-3598.
- Wong ST, Trinh K, Hacker B, Chan GC, Lowe G, et al. 2000. Disruption of the type III adenylyl cyclase gene leads to peripheral and behavioral anosmia in transgenic mice. *Neuron* 27: 487-497.
- Woo CC, Leon M. 1995. Distribution and development of beta-adrenergic receptors in the rat olfactory bulb. *J Comp Neurol* 352: 1-10.
- Woolf TB, Shepherd GM, Greer CA. 1991. Serial reconstruction of granule cell spines in the mammalian olfactory bulb. *Synapse* 7: 181-191.
- Wright DE, Seroogy KB, Lundgren KH, Davis BM, Jennes L. 1995. Comparative localization of serotonin1A, 1C, and 2 receptor subtype mRNAs in rat brain. *J Comp Neurol* 351: 357-373.
- Yamamoto C, Yamamoto T, Iwama K. 1963. The inhibitory systems in the olfactory bulb studied by intracellular recording. *J Neurophysiol* 26: 403-415.
- Yang X, Renken R, Hyder F, Siddeek M, Greer CA, et al. 1998. Dynamic mapping at the laminar level of odor-elicited responses in rat olfactory bulb by functional MRI. *Proc Natl Acad Sci USA* 95: 7715-7720.
- Young WS, Kuhar MJ. 1979. Noradrenergic α_1 and α_2 receptors: Autoradiographic visualization. *Eur J Pharmacol* 59: 317-319.
- Young WS, Kuhar MJ. 1980a. Noradrenergic α_1 and α_2 receptors: Light microscopic autoradiographic localization. *Proc Natl Acad Sci USA* 77: 1696-1700.
- Young WS, Kuhar MJ. 1980b. Radiohistochemical localization of benzodiazepine receptors in rat brain. *J Pharm Exp Ther* 212: 337-346.
- Youngentob SL, Pyrski MM, Margolis FL. 2004. Adenoviral vector-mediated rescue of the OMP-null behavioral phenotype: Enhancement of odorant threshold sensitivity. *Behav Neurosci* 118: 636-642.
- Yuan Q, Harley CW, McLean JH. 2003. Mitral cell α_1 and 5-HT_{2A} receptor colocalization and cAMP coregulation: A new model of norepinephrine-induced learning in the olfactory bulb. *Learn Mem* 10: 5-15.
- Zaborszky L, Carlsen J, Brashear HR, Heimer L. 1986. Cholinergic and GABAergic afferents to the olfactory bulb in the rat with special emphasis on the projection neurons in the nucleus of the horizontal limb of the diagonal band. *J Comp Neurol* 243: 488-509.
- Zhang C, Restrepo D. 2002. Expression of connexin 45 in the olfactory system. *Brain Res* 929: 37-47.
- Zhang C, Restrepo D. 2003. Heterogeneous expression of connexin 36 in the olfactory epithelium and glomerular layer of the olfactory bulb. *J Comp Neurol* 459: 426-439.
- Zhang JH, Sato M, Tohyama M. 1991. Different postnatal development profiles of neurons containing distinct GABAA receptor beta subunit mRNAs (beta 1, beta 2, and beta 3) in the rat forebrain. *J Comp Neurol* 308: 586-613.
- Zhao H, Ivic L, Otaki JM, Hashimoto M, Mikoshiba K, et al. 1998. Functional expression of a mammalian odorant receptor. *Science* 279: 237-242.
- Zheng LM, Caldani M, Jourdan F. 1988. Immunocytochemical identification of luteinizing hormone-releasing hormone-positive fibres and terminals in the olfactory system of the rat. *Neurosci* 24: 567-578.
- Zimmer LA, Ennis M, Shipley MT. 1999. Diagonal band stimulation increases piriform cortex pyramidal cell excitability in vivo. *Neuroreport* 10: 2101-2105.
- Zou Z, Horowitz LF, Montmayeur JP, Snapper S, Buck LB. 2001. Genetic tracing reveals a stereotyped sensory map in the olfactory cortex. *Nature* 414: 173-179.
- Zufall F, Firestein S. 1993. Divalent cations block the cyclic nucleotide-gated channel of olfactory receptor neurons. *J Neurophysiol* 69: 1758-1768.
- Zufall F, Munger SD. 2001. From odor and pheromone transduction to the organization of the sense of smell. *Trends Neurosci* 24: 191-193.
- Zufall F, Firestein S, Shepherd GM. 1991. Analysis of single cyclic nucleotide-gated channels in olfactory receptor cells. *J Neurosci* 11: 3573-3580.